

Name of journal: World Journal of Hepatology

ESPS Manuscript NO: 17035

Columns: Original Article

Prospective Study

Burden of nonalcoholic fatty liver disease and advanced fibrosis in a Texas Hispanic community cohort

Jen-Jung Pan, Susan P Fisher-Hoch, Chaoru Chen, Ariel E Feldstein, Joseph B McCormick, Mohammad H Rahbar, Laura Beretta, Michael B Fallon

Abstract

AIM: To investigate the potential burden of nonalcoholic steatohepatitis (NASH) and advanced fibrosis in a Hispanic community.

METHODS: Four hundred and forty two participants with available ultrasonography data from the Cameron County Hispanic Cohort were included in this study. Each participant completed a comprehensive questionnaire regarding basic demographic information, medical history, medication use, and social and family history including alcohol use. Values of the nonalcoholic fatty liver disease (NAFLD) fibrosis score (NFS), FIB4 index, BARD score, and Aspartate aminotransferase to Platelet Ratio Index (APRI) were computed using the blood samples collected within 6 months of liver

Match Overview

1	Internet 109 words crawled on 20-Nov-2014 www.biomedcentral.com	3%
2	CrossCheck 63 words "AASLD Abstracts", <i>Hepatology</i> , 2012.	1%
3	CrossCheck 61 words "Sunday, 3 September 2006", <i>European Heart Journal</i> , 08/02/2006	1%
4	CrossCheck 57 words "Abstracts", <i>Hepatology</i> , 10/2011	1%
5	CrossCheck 41 words "Monday, 4 September 2006", <i>European Heart Journal</i> , 08/02/2006	1%
6	CrossCheck 39 words "AASLD Abstracts", <i>Hepatology</i> , 2013.	1%
7	Internet 27 words crawled on 10-Sep-2014	1%



Burden of NAFLD and advanced fibrosis in a Texas Hispanic community ()



网页 新闻 图片 视频 更多 ▾ 搜索工具

找到约 95,800 条结果 (用时 0.68 秒)

Google 学术: Burden of NAFLD and advanced fibrosis in a Texas Hispanic community cohort

... fatty liver disease (NAFLD) and diagnostic accuracy of ... - Musso - 被引用次数: 327

Chronic liver disease in the Hispanic population of the ... - Carrion - 被引用次数: 21

Chronic Liver Disease in the Hispanic Population of the ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: AF Carrion - 2011 - 被引用次数: 21 - [相关文章](#)

2011年5月12日 - Keywords: Ethnicity, Fibrosis, Race, Latino, Risk, Epidemiology ...

ethnic differences in liver disease burden remain poorly understood but ... For example, Williams et al²³ reported the ethnic distribution of biopsy-proven NAFLD in a cohort ... of advanced fibrosis or cirrhosis only among Hispanic persons.²⁷ ...

Population-based Risk Factors for Elevated Alanine ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: HQ Qu - 2012 - 被引用次数: 2 - [相关文章](#)

2012年9月5日 - From this we obtain an estimate of the potential burden of NAFLD in our

... The Hispanic population in the city of Brownsville, Cameron County, Texas is one of the poorest in the community; the Cameron County Hispanic Cohort (CCHC) (2) it

网页 新闻 图片 视频 更多 ▾ 搜索工具

找到约 53,700 条结果 (用时 0.61 秒)

Google 学术: Burden of nonalcoholic fatty liver disease and advanced fibrosis in a Texas Hispanic community cohort

... : natural history of non-alcoholic fatty liver disease (... - Musso - 被引用次数: 346

Nonalcoholic fatty liver disease and hepatocellular ... - Starley - 被引用次数: 567

Burden of liver disease in the United States: summary ... - Kim - 被引用次数: 432

Gender and racial differences in nonalcoholic fatty liver ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

作者: JJ Pan - 2014 - 被引用次数: 6 - 相关文章

2014年5月27日 - Nonalcoholic fatty liver disease (NAFLD) is highly associated with obesity and insulin ... (NASH) which may progress to advanced fibrosis and cirrhosis[1,2]. ... Among different racial and ethnic populations in the US, Hispanics a higher disease burden in women or, alternatively, sex differences among ...

Chronic Liver Disease in the Hispanic Population of the US ...

www.medscape.com/viewarticle/750777_3 ▾ 翻译此页

Nonalcoholic fatty liver disease (NAFLD) is the most prevalent chronic liver disease in ... of biopsy-proven NAFLD in a cohort of patients with abnormal liver ultrasounds. ... predictors of advanced fibrosis or cirrhosis only among Hispanic persons. ... Outreach by community health workers is one promising approach to tackle ...

[网页](#) [新闻](#) [图片](#) [视频](#) [更多 ▾](#) [搜索工具](#)

找到约 48,000 条结果 (用时 0.66 秒)

Google 学术: Burden of nonalcoholic fatty liver disease and advanced fibrosis in a Texas Hispanic community cohort

... : natural history of non-alcoholic fatty liver disease (... - Musso - 被引用次数: 352

Nonalcoholic fatty liver disease and hepatocellular ... - Starley - 被引用次数: 570

Burden of liver disease in the United States: summary ... - Kim - 被引用次数: 434

Gender and racial differences in nonalcoholic fatty liver ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

作者: JJ Pan - 2014 - 被引用次数: 7 - 相关文章

2014年5月27日 - Nonalcoholic fatty liver disease (NAFLD) is highly associated with obesity and insulin ... (NASH) which may progress to advanced fibrosis and cirrhosis[1,2]. ... Among different racial and ethnic populations in the US, Hispanics ... a higher disease burden in women or, alternatively, sex differences among ...

Chronic Liver Disease in the Hispanic Population of the US ...

www.medscape.com/viewarticle/750777_3 ▾ 翻译此页

Nonalcoholic fatty liver disease (NAFLD) is the most prevalent chronic liver disease in ... of biopsy-proven NAFLD in a cohort of patients with abnormal liver ultrasounds. ... predictors of advanced fibrosis or cirrhosis only among Hispanic persons. ... Outreach by community health workers is one promising approach to tackle ...