

Name of journal: *World Journal of Hepatology*

ESPS Manuscript NO: 21256

Manuscript Type: MINIREVIEWS

Markers of bacterial translocation in end-stage liver disease

Koutsounas I *et al.* Markers of bacterial translocation in end-stage liver disease

Ioannis Koutsounas, Garyfallia Kaltsa, Spyros I. Siakavellas, Giorgos Bamias

Match Overview

1	CrossCheck 210 words Bellot, Pablo, Rubén Francés, and Jose Such. "Pathological bacterial translocation in cirrhosis: pathophysiology..."	4%
2	Internet 76 words crawled on 08-Oct-2010 lib.bioinfo.pl	2%
3	CrossCheck 47 words Rubén Francés. "A sequential study of serum bacterial DNA in patients with advanced cirrhosis and ascites", H	1%
4	Internet 45 words crawled on 06-Jun-2008 ajpgi.physiology.org	1%
5	CrossCheck 38 words Tony Bruns. "Identification of bacterial DNA in neutrocytic and non-neutrocytic cirrhotic ascites by means of a mi..."	1%
6	CrossCheck 33 words Thomas Sersté. "Detection of bacterial DNA in serum and ascitic fluid of asymptomatic outpatients with cirrhosis..."	1%



Markers of bacterial translocation in end-stage liver disease



学术搜索

找到约 8,660 条结果 (用时0.10秒)

我的著作引用情况



文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Human immunodeficiency virus-related microbial translocation and progression of hepatitis C

A Balagopal, FH Philp, J Astemborski, TM Block... - Gastroenterology, 2008 - Elsevier

... Inclusion of more than one marker in the same model of cirrhosis did not improve the ... Microbial translocation in the host with cirrhosis will lead to less clearance of bacterial products and ... reactive IgG is not currently known, given its correlation with the other markers of microbial ...

被引用次数: 167 相关文章 所有 10 个版本 引用 保存

nih.gov 中的 [HTML]

[HTML] ... for end - stage liver disease score and systemic inflammatory response are major prognostic factors in patients with cirrhosis and acute functional renal failure

D Thabut, J Massard, A Gangloff, N Carbonell... - ..., 2007 - Wiley Online Library

... they had SIRS, as defined above, in response to a proven or suspected microbial event ... 10 Severity of liver disease was assessed according to model for end-stage liver disease (MELD)14 ... with new diagnostic criteria,22 renal failure in the setting of ongoing bacterial infection but ...

被引用次数: 145 相关文章 所有 5 个版本 引用 保存

wiley.com 中的 [HTML]

Renal failure in cirrhosis

P Ginès, RW Schrier - New England Journal of Medicine, 2009 - Mass Medical Soc

... compared with those who have cirrhosis but do not have these markers of bacterial ... these cases, the peritonitis is most often caused by gram-negative bacteria due to ... of antibiotic prophylaxis (either norfloxacin or third-generation cephalosporins) to prevent bacterial infections.11 ...

被引用次数: 411 相关文章 所有 24 个版本 引用 保存

pbworks.com 中的 [PDF]

Bacterial translocation and its consequences in patients with cirrhosis

C Guarner, G Soriano - European journal of gastroenterology & ..., 2005 - journals.lww.com

... submitted to laparotomy, especially in those with more severe liver dysfunction [5 ... been observed in cirrhotic rats that not all overgrowing intestinal bacteria translocate to mesenteric ... bacterial



Markers of bacterial translocation in end-stage liver disease



学术搜索

找到约 8,670 条结果 (用时0.08秒)

我的著作引用情况



文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Human immunodeficiency virus-related microbial translocation and progression of hepatitis C

A Balagopal, FH Philp, J Astemborski, TM Block... - Gastroenterology, 2008 - Elsevier

... Inclusion of more than one marker in the same model of cirrhosis did not improve the ... Microbial translocation in the host with cirrhosis will lead to less clearance of bacterial products and ... reactive IgG is not currently known, given its correlation with the other markers of microbial ...

被引用次数: 167 相关文章 所有 10 个版本 引用 保存

nih.gov 中的 [HTML]

[HTML] ... for end - stage liver disease score and systemic inflammatory response are major prognostic factors in patients with cirrhosis and acute functional renal failure

D Thabut, J Massard, A Gangloff, N Carbonell... - ..., 2007 - Wiley Online Library

... they had SIRS, as defined above, in response to a proven or suspected microbial event ... 10 Severity of liver disease was assessed according to model for end-stage liver disease (MELD)14 ... with new diagnostic criteria,22 renal failure in the setting of ongoing bacterial infection but ...

被引用次数: 145 相关文章 所有 5 个版本 引用 保存

wiley.com 中的 [HTML]

Renal failure in cirrhosis

P Ginès, RW Schrier - New England Journal of Medicine, 2009 - Mass Medical Soc

... compared with those who have cirrhosis but do not have these markers of bacterial ... these cases, the peritonitis is most often caused by gram-negative bacteria due to ... of antibiotic prophylaxis (either norfloxacin or third-generation cephalosporins) to prevent bacterial infections.11 ...

被引用次数: 413 相关文章 所有 24 个版本 引用 保存

pbworks.com 中的 [PDF]

Bacterial translocation and its consequences in patients with cirrhosis

C Guarner, G Soriano - European journal of gastroenterology & ..., 2005 - journals.lww.com

... submitted to laparotomy, especially in those with more severe liver dysfunction [5 ... been observed in cirrhotic rats that not all overgrowing intestinal bacteria translocate to mesenteric ... bacterial

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 233,000 条结果 (用时 0.76 秒)

Google 学术: Markers of bacterial translocation in end-stage liver disease

... immunodeficiency virus-related microbial translocation ... - Balagopal - 被引用次数: 166

Model for end-stage liver disease score and systemic ... - Thabut - 被引用次数: 145

Renal failure in cirrhosis - Ginès - 被引用次数: 416

Bacterial translocation and changes in the intestinal ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: AW Yan - 2012 - 被引用次数: 26 - 相关文章

2012年4月27日 - Cirrhosis is an end-stage disease and one of the leading causes of ... as a surrogate marker of bacterial translocation, associated with marked ...

[PDF] Markers of immunity and bacterial translocation in cirrhosis

www.danmedj.dk/.../IPORTAL.wwwpob_page.show?_...Markers... ▾ [翻译此页](#)

of markers of bacterial translocation to immune activation in decompensated ...

Cirrhosis is considered the end-stage of a variety of different chronic liver ...

Complications of Cirrhosis: Evaluation and Management

<https://books.google.com.hk/books?isbn=3319136143> - [翻译此页](#)

Andrew Keaveny, Andrés Cárdenas - 2015 - Medical

Increased translocation of bacteria or bacterial products has been described to be ...

Acute phase proteins are commonly used as clinical markers of infection in ..., levels of acute phase proteins in end-stage liver disease, such as underlying ...