

Neurosurgical procedures in patients with liver cirrhosis: A systemic revi



网页 图片 新闻 视频 购物 更多▼ 搜索工具

找到约 1,090,000 条结果 (用时 0.60 秒)

显示的是以下查询字词的结果: Neurosurgical procedures in patients with liver cirrhosis: A **systematic** review

仍然搜索: Neurosurgical procedures in patients with liver cirrhosis: A systemic review

Google 学术: Neurosurgical procedures in patients with liver cirrhosis: A systematic review

- ... coagulopathy in nonhemophilic neurosurgical patients Park 被引用次数: 129
- ... surgical mortality in patients with cirrhosis Northup 被引用次数: 178
- ... for treatment of severe bleeding: a systematic review Levi 被引用次数: 311

Outcomes of patients with cirrhosis undergoing non-hepatic ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页 作者: F Millwala - 2007 - 被引用次数: 61 - 相关文章

2007年8月14日 - The prevalence of chronic liver disease is increasing in the US and the rest ... The reported mortality rates in patients with cirrhosis undergoing various surgical procedures with significant coagulopathy, neurosurgery usually requires an INR surgery in patients with liver cirrhosis: a systematic review.

Model for End-Stage Liver Disease (MELD) Predicts ...

www.ncbi,nlm.nih.gov > ... > PubMed Central (PMC) ▼翻译此页

作者: PG Northup - 2005 - 被引用次数: 177 - 相关文章

Adult patients with a diagnosis of cirrhosis undergoing nontransplant ... patients will require surgical procedures, often when their liver disease is poorly compensated. ... the

18477-Review BY CHING-CHANG CHEN

Quotes Excluded Bibliography Excluded

8% SIMILAR

Text-Only Report

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 18477

Manuscript Type: REVIEW

Neurosurgical procedures in patients with liver cirrhosis: A review

Ching-Chang Chen, Yin-Cheng Huang, Chun-Nan Yeh

Abstract

Liver cirrhosis, a devastating liver fibrosis caused by hepatitis/inflammation or tumors, is a major comorbid factor in known surgery fields, such as cardiovascular and abdominal surgeries. It is important to review possible

Match Overview			
4		10	
1	Internet 112 words crawled on 04-Dec-2012 www.neurocirurglabr.com	3%	
2	CrossCheck 82 words Liao, Jen-Chung, Wen-Jer Chen, Lih-Hui Chen, Chi-Chien Niu, Tsai-Shen Fu, Po-Liang Lai, and Tsung-Ting Tsai, "Co	2%	
3	CrossCheck 57 words Huang, H.H., "Spontaneous intracranial hemorrhage in cl hotic patients", Clinical Neurology and Neurosurgery, 2008	1%	
4	CrossCheck 39 words Ching-Chang Chen, "Brain surgery in patients with liver combosis: Clinical article", Journal of Neurosurgery, 05/25/201	1%	
5	Internet 10 words crawled on 27-Jan-2015 doaj.org	<1%	





Neurosurgical procedures in patients with liver cirrhosis: A review



Q

网页 图片 新闻 视频 更多▼ 搜索工具

找到约 9,710,000 条结果 (用时 0.60 秒)

Google 学术: Neurosurgical procedures in patients with liver cirrhosis: A review

- ... surgical mortality in patients with cirrhosis Northup 被引用次数: 175
- ... coagulopathy in nonhemophilic neurosurgical patients Park 被引用次数: 128
- ... factor VII for coagulopathy in fulminant hepatic failure ... Shami 被引用次数: 172

Surgery in the Patient with Liver Disease

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▼ 翻译此页

作者: LS Friedman - 2010 - 被引用次数: 51 - 相关文章

Most surgical procedures, whether performed under general, spinal or epidural ... In a patient with liver disease, surgical risk depends on the severity of liver disease, nature of Boyer, New Haven: Thank you, Larry, for this very nice review.

Model for End-Stage Liver Disease (MELD) Predicts ...

www.ncbi.nlm.nih.gov>...> PubMed Central (PMC) ▼ 翻译此页

作者: PG Northup - 2005 - 被引用次数: 175 - 相关文章

We sought to determine the ability of the Model for End-Stage Liver Disease (MELD) In patients that had more than one surgical procedure, a preoperative MELD ... 30 days after the procedure, their status was verified by review of electronic

Management of thrombocytopenia due to liver cirrhosis: A ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▼ 翻译此页

作者: H Hayashi - 2014 - 被引用次数: 10 - 相关文章

2014年3月14日 - In this review, we describe the current management of ... However, the interventional and/or surgical treatments aimed at reversing portal In patients with



Scholar

About 24,000 results (0.11 sec)

Articles

Case law

My library

Any time

Since 2015

Since 2014

Since 2011

Custom range...

Sort by relevance

Sort by date

✓ include patents

✓ include citations

Create alert

[HTML] Model for End-Stage Liver Disease (MELD) predicts nontransplant surgical mortality in [HTML] from nih.gov patients with cirrhosis

PG Northup, RC Wanamaker, VD Lee... - Annals of ..., 2005 - ncbi.nlm,nih.gov

... The remainder of the **procedures** were thoracic, **neurosurgical**, or urologic. ... be a useful tool in counseling **cirrhotic patients** regarding their risk for a surgical **procedure**. ... a postoperative mortality of 5%. Thus, strong consideration should be given to performing **procedures** on this ... Cited by 178 Related articles All 7 versions Cite Save

Recombinant activated factor VII for the rapid correction of coagulopathy in nonhemophilic neurosurgical patients

P Park, ME Fewel, HJ Garton, BG Thompson... - Neurosurgery, 2003 - journals.lww.com

... OBJECTIVE: Coagulopathy is a significant contraindication for **neurosurgery**. ... Published experience with spinal or intracranial **procedures** using rFVIIa in nonhemophilic **patients** is limited. ... There is only one case report in the **neurosurgical** literature describing the use of rFVIIa ...

Cited by 128 Related articles All 5 versions Cite Save

Recombinant activated factor VII for coagulopathy in fulminant hepatic failure compared with [PDF] from wiley.com conventional therapy

VM Shami, SH Caldwell, EE Hespenheide... - Liver ..., 2003 - Wiley Online Library

... also is enhanced.7,18 Performance of invasive, but potentially life-saving, **procedures** in **patients** ... team, and in the United States, these devices are most often placed by **neurosurgeons**. ... placement of an ICP monitor, but it is our experience that **neurosurgical** consultants always ...

Cited by 174 Related articles All 8 versions Cite Save

Placebo surgery in trials of therapy for Parkinson's disease

WJ Weiner, J Burris, C Freed - N Engl J Med, 2000 - Mass Medical Soc

... associated with pallidotomy is related to the varying skills of the operating **neurosurgeon**. ... is the only solution for proving the validity of the **neurosurgical procedures** discussed in ... community, including Dr, Freeman's coauthor Dr. Olanow.1 Because no **neuro-surgical** trial had ... Cited by 221 Related articles All 13 versions Cite Save

Primary sclerosing cholangitis

NF LaRusso, RH Wiesner, J Ludwig... - New England Journal ..., 1984 - Mass Medical Soc

... This process leads to the formation of fibrous septa (stage 3) and biliary cirrhosis ... ulcerative colitis

[PDF] from academia.edu