

27

Name of Journal: World Journal of Hepatology

ESPS Manuscript NO: 18324

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Histopathological differences utilizing the NAFLD Activity score criteria ⁸ in diabetic (T2DM) and non-diabetic patients with NAFLD

Bharat K Puchakayala, Siddharth Verma, Pushpjeet Kanwar, John Hart, Raghavendra R Sanivarapu, Smruti R Mohanty

Abstract

AIM: To study clinical and histopathological features of nonalcoholic fatty liver disease (NAFLD) in patients with and without type 2 diabetes mellitus (T2DM) using updated nonalcoholic steatohepatitis clinical research network (NASH-CRN) grading system.

METHODS: We retrospectively analyzed data of 235 patients with biopsy proven NAFLD with and without T2DM. This database was utilized in the previously published study comparing ethnicity outcomes in NAFLD by the same corresponding

Match Overview

1	Internet 141 words crawled on 05-Feb-2015 www.endo-gastrosantafe.com.ar	2%
2	CrossCheck 120 words "AASLD Abstracts"; Hepatology, 2013.	2%
3	Internet 93 words crawled on 22-Apr-2015 www.ncbi.nlm.nih.gov	1%
4	CrossCheck 60 words Mohanty, S.R.. "Influence of ethnicity on histological differences in non-alcoholic fatty liver disease", Journal of Hepato	1%
5	CrossCheck 53 words "Abstracts of 51st EASD Annual Meeting", Diabetologia, 2015.	1%
6	CrossCheck 49 words "Abstracts", Hepatology, 10/2011	1%
7	CrossCheck 46 words Younossi, Z.M.. "Nonalcoholic fatty liver disease in patients with type 2 diabetes", Clinical Gastroenterology and Hepat	1%
	Internet 40 words	10%



and non-diabetic patients with Nonalcoholic Fatty Liver Disease (NAFLD)



网页 新闻 图片 视频 更多 搜索工具

找到约 26,500 条结果 (用时 0.92 秒)

您是不是要找: Histopathological differences utilizing the NAS criteria in **diabetes** (T2DM) and non-diabetic patients with Nonalcoholic Fatty Liver Disease (NAFLD)

Google 学术: Histopathological differences utilizing the NAS criteria in **diabetic** (T2DM) and non-diabetic patients with Nonalcoholic Fatty Liver Disease (NAFLD)

... in patients with non-alcoholic fatty liver disease: a ... - Haukeland - 被引用次数: 133

... and management of non-alcoholic fatty liver disease: ... - Chalasani - 被引用次数: 747

... non-diabetic patients with non-alcoholic steatohepatitis ... - Chalasani - 被引用次数: 79

Clinical, laboratory and histological associations in adults ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) - 翻译此页

作者: NM Bass - 2010 - 被引用次数: 174 - 相关文章

Nonalcoholic fatty liver disease (NAFLD) affects 10–30% of the general U.S. ... for the Treatment of Nondiabetic Patients with Nonalcoholic Steatohepatitis” ... using the previously published NASH CRN NAFLD Activity Score (NAS) and fibrosis score. ... of co-morbid conditions including hypertension and type 2 diabetes.

Non-alcoholic fatty liver disease and diabetes: From ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

作者: NC Leite - 2014 - 被引用次数: 4 - 相关文章



Histopathological differences utilizing the NAFLD Activity score criteria in dia  

网页 图片 新闻 视频 更多 搜索工具

找到约 79,200 条结果 (用时 0.83 秒)

Clinical spectrum of non-alcoholic fatty liver disease in ...
www.sciencedirect.com/science/article/.../S221464741400018X  翻译此页
作者: GBB Goh - 2015 - 相关文章

The NAFLD fibrosis score may be less accurate in non-diabetics. ... Patients with type 2 diabetes mellitus (DM) have an increased risk of developing ... were read using standardised well defined histological criteria [18] established by the ... Baseline characteristics and differences in demographic, clinical, histological and ...

Non-alcoholic fatty liver disease and type 2 diabetes mellitus
www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC)  翻译此页

作者: G Firneisz - 2014 - 被引用次数: 20 - 相关文章
2014年7月21日 - Type 2 diabetes mellitus (T2DM), insulin resistance (IR), obesity, metabolic syndrome and ... The differences in the prevalence and clinical-pathological ... In most patients with NAFLD, a non-invasive score proposed by Angulo et ... The fatty liver index (FLI) -measured using ¹HMRS- in women with previous ...

Clinical, laboratory and histological associations in adults ...
www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC)  翻译此页

作者: NM Bass - 2010 - 被引用次数: 200 - 相关文章
Nonalcoholic fatty liver disease (NAFLD) affects 10–30% of the general U.S. ... The PIVENS trial was limited to patients without diabetes or a history of ... using the previously published NASH CRN NAFLD Activity Score (NAS) and fibrosis score. ... It is worth noting that 16% of biopsies did not meet NASH criteria yet had a ...

[PDF] The Diagnosis and Management of Non-Alcoholic ... - AA...
<https://www.aasld.org/.../NonalcoholicFattyLiverDisease2012.p...>  翻译此页



[网页](#) [图片](#) [新闻](#) [视频](#) [更多 ▾](#) [搜索工具](#)

找到约 74,200 条结果 (用时 0.81 秒)

Clinical spectrum of non-alcoholic fatty liver disease in ...

www.sciencedirect.com/science/article/.../S221464741400018X ▾ [翻译此页](#)

作者: GBB Goh - 2015 - [相关文章](#)

The NAFLD fibrosis score may be less accurate in non-diabetics. ... Patients with type 2 diabetes mellitus (DM) have an increased risk of developing ... were read using standardised well defined histological criteria [18] established by the ... Baseline characteristics and differences in demographic, clinical, histological and ...

Non-alcoholic fatty liver disease and type 2 diabetes mellitus

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: G Firneisz - 2014 - 被引用次数: 21 - [相关文章](#)

2014年7月21日 - Type 2 diabetes mellitus (T2DM), insulin resistance (IR), obesity, metabolic syndrome and ... The differences in the prevalence and clinical-pathological ... In most patients with NAFLD, a non-invasive score proposed by Angulo et ... The fatty liver index (FLI) -measured using ¹HMRs- in women with previous ...

^[PDF] The Diagnosis and Management of Non-Alcoholic ... - AA...

<https://www.aasld.org/.../NonalcoholicFattyLiverDisease2012.p...> ▾ [翻译此页](#)

作者: N Chalasani - 2012 - 被引用次数: 971 - [相关文章](#)

Fatty Liver Disease in Children; NAS, NAFLD Activity Score; CK18; Cytokeratin 18 Fragments; ELF, ... study of patients with T2DM showed a 69% prevalence.

Diagnosis and Evaluation of Nonalcoholic Fatty Liver Disease

www.hindawi.com/journals/jdr/2012/145754/ ▾ [翻译此页](#)

作者: M Obika - 2011 - 被引用次数: 72 - [相关文章](#)

2011年8月31日 - NAFLD and type 2 diabetes mellitus (T2DM) frequently coexist ... or not liver biopsy is required to confirm a diagnosis of NAFLD [8]. A significant elevation of the serum TRX levels was demonstrated in patients with NASH in comparison to The NAFLD Activity Score (NAC) is used for the histological