

Name of Journal: *World Journal of Hepatology*

ESPS Manuscript NO: 21461

Manuscript Type: TOPIC HIGHLIGHTS

## Bile acid receptors and Nonalcoholic Fatty Liver Disease

Liyun Yuan, Kiran Bambha

### Abstract

With the high prevalence of obesity, diabetes, and other features of the metabolic

syndrome in United States, nonalcoholic fatty liver disease (NAFLD) has inevitably

become a very prevalent chronic liver disease and is now emerging as one of the

### Match Overview

- |   |   |    |
|---|---|----|
| 1 | Internet 72 words<br>crawled on 18-Oct-2014<br><a href="http://www.science.gov">www.science.gov</a>   | 2% |
| 2 | Internet 49 words<br>crawled on 04-Jan-2010<br><a href="http://www.jci.org">www.jci.org</a>   | 2% |
| 3 | CrossCheck 38 words<br>Dufour, Jean-François. "Hepatology Highlights", Hepatology, 2014.  | 1% |
| 4 | CrossCheck 24 words<br>"AASLD Abstracts (pp. 179A–281A)", Hepatology, 10/2003   | 1% |
| 5 | CrossCheck 23 words<br>Zhou, Huiping, and Phillip B. Hylemon. "Bile Acids are Nutrient-Signaling Hormones", Steroids, 2014.                   | 1% |
| 6 | CrossCheck 18 words<br>William Insull. "Clinical Utility of Bile Acid Sequestrants in the Treatment of Dyslipidemia: A Scientific Review", So | 1% |
| 7 | CrossCheck 18 words<br>Tiangang Li. "Regulation of Bile Acid and Cholesterol M  | 1% |



## 学术搜索

找到约 13,500 条结果 (用时 0.04 秒)



文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

 包括专利 包含引用 创建快讯

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

## Bile acid receptors in non-alcoholic fatty liver disease

Y Li, K Jadhav, Y Zhang - Biochemical pharmacology, 2013 - Elsevier

Abstract Accumulating data have shown that bile acids are important cell signaling molecules, which may activate several signaling pathways to regulate biological processes.

Bile acids are endogenous ligands for the farnesoid X receptor (FXR) and TGR5, a G-

被引用次数: 25 相关文章 所有 5 个版本 引用 保存

europepmc.org 中的 [HTML]

## Efficacy and safety of the farnesoid X receptor agonist obeticholic acid in patients with type 2 diabetes and nonalcoholic fatty liver disease

S Mudaliar, RR Henry, AJ Sanyal, L Morrow... - Gastroenterology, 2013 - Elsevier

... Bile acids (BAs), classically recognized as detergent-like compounds involved in lipid absorption and ... 26 OCA shows 100-fold greater FXR agonistic activity than chenodeoxycholic acid, the natural ...

27 OCA does not activate other nuclear receptors, 28 and its activities appear to ...

被引用次数: 151 相关文章 所有 7 个版本 引用 保存

gastrojournal.org 中的 [HTML]

## Nonalcoholic fatty liver disease

P Angulo - New England Journal of Medicine, 2002 - Mass Medical Soc

... This pathway of fatty-acid metabolism is closely related to mitochondrial  $\beta$ -oxidation and ... Increased intrahepatic levels of fatty acids provide a source of oxidative stress, which may in ... infiltration, rapid weight loss may promote necroinflammation, portal fibrosis, and bile stasis. 26 ...

被引用次数: 4114 相关文章 所有 21 个版本 引用 保存

gastrohep.com 中的 [PDF]

## [HTML] Lipid metabolism and liver inflammation. II. Fatty liver disease and fatty acid oxidation

JK Reddy, MS Rao - ... of Physiology-Gastrointestinal and Liver ..., 2006 - Am Physiological Soc

... steatohepatitis; peroxisome proliferator-activated receptors; nonalcoholic fatty liver disease; obesity active very-long-chain fatty acids (containing 20 or more carbon atoms); 2-methylbranched

physiology.org 中的 [HTML]



## 学术搜索

## 文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

 包括专利 包含引用

## Bile acid receptors in non-alcoholic fatty liver disease

Y Li, K Jadhav, Y Zhang - Biochemical pharmacology, 2013 - Elsevier

Abstract Accumulating data have shown that bile acids are important cell signaling molecules, which may activate several signaling pathways to regulate biological processes. Bile acids are endogenous ligands for the farnesoid X receptor (FXR) and TGR5, a G- ...

被引用次数: 30 相关文章 所有 5 个版本 引用 保存

以上显示的是最相近的搜索结果。 [查看全部搜索结果](#)

网页 图片 新闻 视频 更多 ▾ 搜索工具

找到约 97,000 条结果 (用时 0.42 秒)

### Google 学术: Bile acid receptors and nonalcoholic fatty liver disease

Bile acid receptors in non-alcoholic fatty liver disease - Li - 被引用次数: 30

... type 2 diabetes and nonalcoholic fatty liver disease - Mudaliar - 被引用次数: 174

Nonalcoholic fatty liver disease - Angulo - 被引用次数: 4263

### Bile acid receptors in non-alcoholic fatty liver disease.

[www.ncbi.nlm.nih.gov/pubmed/23988487](http://www.ncbi.nlm.nih.gov/pubmed/23988487) ▾ 翻译此页

作者: Y Li - 2013 - 被引用次数: 29 - 相关文章

Biochem Pharmacol. 2013 Dec 1;86(11):1517-24. doi: 10.1016/j.bcp.2013.08.015.

Epub 2013 Aug 26. Bile acid receptors in non-alcoholic fatty liver disease.

### Bile acid receptor activation modulates hepatic monocyte ...

[www.ncbi.nlm.nih.gov/pubmed/23460643](http://www.ncbi.nlm.nih.gov/pubmed/23460643) ▾ 翻译此页

作者: RH McMahan - 2013 - 被引用次数: 42 - 相关文章

2013年3月4日 - Bile acid receptor activation modulates hepatic monocyte activity and improves nonalcoholic fatty liver disease. McMahan RH(1), Wang XX, ...

### Non-Alcoholic Fatty Liver Disease: The Bile Acid-Activated ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > PubMed Central (PMC) ▾ 翻译此页

作者: M Fuchs - 2012 - 被引用次数: 54 - 相关文章

2011年12月7日 - Non-Alcoholic Fatty Liver Disease: The Bile Acid-Activated Farnesoid X receptor (FXR) has been shown to play a role not only in bile acid but ...