

Name of journal: World Journal of Hepatology

ESPS Manuscript NO: 13640

Columns: REVIEW

Composite prognostic models across the NAFLD spectrum: clinical application in developing countries

Hilmar K Lückhoff, Frederik C Kruger, Maritha J Kotze

Abstract

The burgeoning prevalence of obesity and the metabolic syndrome in the developing world has seen non-alcoholic fatty liver disease (NAFLD) emerge as a growing health concern and cause of significant financial burden. NAFLD defines a spectrum of hepatic abnormalities ranging from simple steatosis through steatohepatitis (NASH), progressive hepatic fibrosis and finally cirrhosis along with its associated complications. This characteristic heterogeneity in pathological severity and clinical outcomes necessitates the development of scientifically sound classification schemes in order to stratify patients into meaningful prognostic subgroups with the goal of

Match Overview

1	CrossCheck 67 words "Abstracts", Hepatology, 10/2011	1%
2	CrossCheck 53 words de Lange, Naydene, and Claudia Mitchell. "Building a future without gender violence: rural teachers and youth in rural ...	1%
3	CrossCheck 25 words Mikako Obika. "Diagnosis and Evaluation of Nonalcoholic Fatty Liver Disease", Experimental Diabetes Research, 2012	1%
4	CrossCheck 12 words "Abstracts of the EASD, Vienna 2009", Diabetologia, 09/2009	<1%
5	Internet 11 words crawled on 17-Dec-2012 eprints.qut.edu.au	<1%
6	Internet 10 words crawled on 25-Sep-2010 www.sou.strefa.pl	<1%
7	Internet 9 words crawled on 11-Oct-2008 intl-jcem.endojournals.org	<1%
8	Internet 8 words crawled on 22-Apr-2010 www.wjgnet.com	<1%
9	CrossCheck 6 words Dina S. El-Agamy. "Comparative effects of curcumin and silymarin on aflatoxin B1-induced liver injury in rats", Archiv	<1%



is across the NAFLD spectrum: Clinical application in developing countries



网页 图片 新闻 视频 更多 搜索工具

找到约 221,000 条结果 (用时 0.51 秒)

Google 学术 : Composite prognostic models across the NAFLD spectrum: Clinical application in developing countries

NAFLD fibrosis score: a prognostic predictor for ... - Treeprasertsuk - 被引用次数 : 24

... of Human and Mouse Model Serum Reveals Putative ... - Barr - 被引用次数 : 66

Nonalcoholic steatohepatitis: risk factors and diagnosis - Adams - 被引用次数 : 37

NAFLD fibrosis score: A prognostic predictor for mortality ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) > 翻译此页

作者 : S Treeprasertsuk - 2013 - 被引用次数 : 24 - 相关文章

2013年2月28日 - The NAFLD fibrosis score (NFS) was used to separate NAFLD patients ... of the most common causes of chronic liver disease in Western countries [1]. ... Thus, a benefit of the current study is an extension of the clinical use of the ... with NAFLD are not uniform across the spectrum of the disease[3,22,23].

Systematic review: the diagnosis and staging of non ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) > 翻译此页

作者 : JK Dowman - 2011 - 被引用次数 : 147 - 相关文章

2010年12月29日 - The development of non-alcoholic steatohepatitis (NASH) and fibrosis ... The identification and management of this at-risk group remains a clinical challenge. ... Non-alcoholic fatty liver disease (NAFLD) encompasses a spectrum of ... The AUROC was 0.84, and application of this model to the study ...

Noninvasive Diagnosis of NASH and Liver Fibrosis Within ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) > 翻译此页

作者 : N Alkhouri - 2012 - 被引用次数 : 4 - 相关文章

The histologic spectrum of NAFLD ranges from simple steatosis to nonalcoholic ... Due to the high prevalence of NAFLD, there has been an urgent need to develop reliable ... in association with NASH, are important factors in the prognosis of NAFLD.

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3689658/ at use a combination of clinical and ...

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 197,000 条结果 (用时 0.72 秒)

Google 学术: Composite prognostic models across the NAFLD spectrum: Clinical application in developing countries

NAFLD fibrosis score: a prognostic predictor for ... - Treeprasertsuk - 被引用次数: 25

... of Human and Mouse Model Serum Reveals Putative ... - Barr - 被引用次数: 66

Nonalcoholic steatohepatitis: risk factors and diagnosis - Adams - 被引用次数: 37

NAFLD fibrosis score: A prognostic predictor for mortality ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: S Treeprasertsuk - 2013 - 被引用次数: 25 - [相关文章](#)

2013年2月28日 - The NAFLD fibrosis score (NFS) was used to separate NAFLD patients ... of the most common causes of chronic liver disease in Western countries[1].

..... Thus, a benefit of the current study is an extension of the clinical use of the ... with NAFLD are not uniform across the spectrum of the disease[3,22,23].

Noninvasive Diagnosis of NASH and Liver Fibrosis Within ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: N Alkhouri - 2012 - 被引用次数: 4 - [相关文章](#)

The histologic spectrum of NAFLD ranges from simple steatosis to nonalcoholic ... Due to the high prevalence of NAFLD, there has been an urgent need to develop reliable ... in association with NASH, are important factors in the prognosis of NAFLD. Examples of predictive models that use a combination of clinical and