

Match Overview  

There are no matching sources for this report.

Name of journal: World Journal of Hepatology

ESPS Manuscript NO: 24851

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

HBsAg clearance in inactive HBsAg carriers treated by peginterferon alfa-2a

Ming-Hui Li, Yao Xie, Lu Zhang, Yao Lu, Ge Shen, Shu-Ling Wu, Min Chang, Cai-Qin

Mu, Lei-Ping Hu, Wen-Hao Hua, Shu-Jing Song, Shu-Feng Zhang, Jun Cheng,

Dao-Zhen Xu

HBsAg clearance in inactive HBsAg carriers treated by peginterferon alfa-2a

全部 图片 视频 新闻 购物 地图 图书

找到约 17,200 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[HBsAg Clearance in Inactive Chronic HBsAg Carriers After ...](https://clinicaltrials.gov/ct2/show/NCT01471535)

<https://clinicaltrials.gov/ct2/show/NCT01471535>

31 Oct 2011 ... Drug: Pegylated interferon alfa-2a ... the inactive chronic HBsAg carriers were treated with peginterferon alpha 2a for 72 weeks and followed for ...

[Clinical Implications of HBsAg Quantification in Patients with ...](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3326981/)

www.ncbi.nlm.nih.gov/pmc/articles/PMC3326981/

During antiviral treatment, HBsAg levels decline more rapidly in patients under ... Given the low rates of HBsAg clearance on NUC therapy, new studies to test whether ... for HBeAg and those who remain persistently inactive carriers of HBsAg. A response guided approach to peg-interferon alfa-2a at weeks 12 and 24 ...

[\[PDF\] 308 KB - International Medical Press](http://www.intmedpress.com/serveFile.cfm?sUID=316234ae-a6a7-41d1-b4db...)

www.intmedpress.com/serveFile.cfm?sUID=316234ae-a6a7-41d1-b4db...

tive hepatitis B surface antigen (HBsAg) carriers might encounter HBV ... negative chronic hepatitis, most inactive carriers remain ... clearance during the natural course of infection or after antiviral treatment with peginterferon alpha-2a.

[Micropathology: News](http://www.micropathology.com/0414-hepb.php)

www.micropathology.com/0414-hepb.php

HBsAg clearance in inactive HBsAg carriers treated by peginterferon alfa-2a

全部 图片 视频 新闻 购物 地图 图书

找到约 17,200 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[HBsAg Clearance in Inactive Chronic HBsAg Carriers After ...](https://clinicaltrials.gov/ct2/show/NCT01471535)

<https://clinicaltrials.gov/ct2/show/NCT01471535>

31 Oct 2011 ... Drug: Pegylated interferon alfa-2a ... the inactive chronic HBsAg carriers were treated with peginterferon alpha 2a for 72 weeks and followed for ...

[Clinical Implications of HBsAg Quantification in Patients with ... - NCBI](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3326981/)

www.ncbi.nlm.nih.gov/pmc/articles/PMC3326981/

During antiviral treatment, HBsAg levels decline more rapidly in patients under ... Given the low rates of HBsAg clearance on NUC therapy, new studies to test whether Peg-IFN ... for HBeAg and those who remain persistently inactive carriers of HBsAg ... Early on-treatment prediction of response to peginterferon alfa-2a for ...

[\[PDF\] 308 KB - International Medical Press](http://www.intmedpress.com/serveFile.cfm?sUID=316234ae-a6a7-41d1-b4db...)

www.intmedpress.com/serveFile.cfm?sUID=316234ae-a6a7-41d1-b4db...

tive hepatitis B surface antigen (HBsAg) carriers might encounter HBV ... negative chronic hepatitis, most inactive carriers remain ... clearance during the natural course of infection or after antiviral ... treatment with peginterferon alpha-2a.

[Micropathology: News](http://www.micropathology.com/0414-hepb.php)

www.micropathology.com/0414-hepb.php



全部 新闻 图片 视频 更多 ▾ 搜索工具

找到约 19,400 条结果 (用时 0.61 秒)

[PDF] Review Hepatitis B surface antigen seroclearance during c...

www.intmedpress.com/serveFile.cfm?sUID=316234ae-a6a7... ▾ 翻译此页

ive hepatitis B surface antigen (HBsAg) carriers might encounter HBV ... negative chronic hepatitis, most inactive carriers remain ... clearance during the natural course of infection or after antiviral treatment with peginterferon alpha-2a.

Hepatitis B surface antigen quantification: Why and how to us...

www.sciencedirect.com/science/article/.../S0168827811004971 ▾ 翻译此页

作者: HLY Chan - 2011 - 被引用次数: 222 - 相关文章

Since the discovery of hepatitis B surface antigen (HBsAg) by Blumberg in 1965, it has ... It is followed by an immune clearance phase, which may lead to HBeAg Among inactive carriers, HBsAg serum levels were lower in 31 asymptomatic from a small cohort of patients treated with peginterferon alfa-2a and adefovir ...

Changing serum levels of quantitative hepatitis B surface anti...

www.sciencedirect.com/science/article/.../S1607551X14002253 ▾ 翻译此页

作者: YH Kuo - 2015 - 被引用次数: 4 - 相关文章

Lower levels of both HBsAg and HBV DNA might represent an inactive HBV infection. ... The serum concentration of HBV surface antigen (HBsAg) is proposed to reflect the ... HBV carriers to evaluate the association of HBV infection and treatment. response to peginterferon alfa-2a in HBeAg-negative chronic hepatitis B.

HBsAg Clearance in Inactive Chronic HBsAg Carriers After In...

<https://clinicaltrials.gov/ct2/show/NCT01471535> ▾ 翻译此页

2011年10月31日 - Hepatitis B surface antigen loss/seroconversion, considered to be the ideal ... Chronic Hepatitis B Carriers Treated With Peginterferon Alpha-2a ...