

Name of Journal: World Journal of Hepatology

ESPS Manuscript NO: 24933

Manuscript Type: Retrospective study

Clinical characteristics and progression of liver abscess caused by toxocara

Kyung Ho Ha¹, Jung Eun Song², Byung Seok Kim², Chang Hyeong Lee²

Match Overview

1	Internet 29 words crawled on 20-Mar-2016 bmj.fmed.uniba.sk	1%
2	Internet 25 words crawled on 22-Dec-2013 www.hill-dale.com	1%
3	Internet 24 words crawled on 16-Sep-2013 parasitol.or.kr	1%
4	Internet 10 words crawled on 22-Sep-2009 www.pubmedcentral.nih.gov	<1%
5	CrossCheck 10 words Mazur-Melewska, Katarzyna, Magdalena Figlerowicz, Agnieszka Cwalińska, Hanna Mikoś, Katarzyna Jończyk-Poto	<1%
6	Internet 10 words crawled on 11-Mar-2016 ateneo.unmsm.edu.pe	<1%

学术搜索

找到约 1,180 条结果 (用时0.08秒)

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

时间不限
2016以来
2015以来
2012以来
自定义范围...

Toxocariasis: clinical aspects, epidemiology, medical ecology, and molecular aspects
D Despommier - Clinical microbiology reviews, 2003 - Am Soc Microbiol
... 5), and brain (Fig. 6), produce symptoms characteristic of VLM. ... OLM is diagnosed primarily on the basis of clinical criteria during an ophthalmologic examination. ... In one study, only 45% of patients with clinically diagnosed OLM had titers higher than 1:32 (51). ...
被引用次数: 539 相关文章 所有 12 个版本 引用

asm.org 中的 [HTML]

按相关性排序
按日期排序

Schistosoma mansoni: assessment of morbidity before and after control
JR Lambertucci, JC Serufo, R Gerspacher-Lara... - Acta tropica, 2000 - Elsevier
... persistent infection and an evolving pathological process, hepatic enlargement is clinically recognized in ... The most common characteristics of the clinical syndrome are: (i) an indolent febrile ...
Clinical and radiological findings are similar to those associated with other causes of ...
被引用次数: 142 相关文章 所有 7 个版本 引用

搜索所有网页
中文网页
简体中文网页

☒ 包括专利
☒ 包含引用

[HTML] Pyogenic abscesses and parasitic diseases
JR Lambertucci, AA Rayes, JC Serufo... - Revista do Instituto de ..., 2001 - SciELO Brasil
... Clinically, pyomyositis is characterized by fever, localized muscle pain and stiffness, swelling, and ... Demographic characteristics of the control group were similar to those of the study group: ...
Our clinical and experimental data suggest an association between T. canis infection and ...
被引用次数: 47 相关文章 所有 9 个版本 引用

scielo.br 中的 [HTML]

Visceral larva migrans syndrome complicated by liver abscess
A Rayes, D Teixeira, V Nobre, JC Serufo... - ... journal of infectious ..., 1999 - Taylor & Francis
... hepatitis (4), my- ocarditis (5) and pleural effusion (6). Anaemia, leukocytosis, eosinophilia and hypergamaglob- ulinaemia are common laboratory features of VLM (2). In our patient, the initial symptoms were suggestive of ... Clinical and laboratorial features of visceral ...
被引用次数: 18 相关文章 所有 5 个版本 引用

Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm

researchgate.net



Clinical Characteristics and Progression of Liver Abscess Caused by Toxocara



Scholar

About 707 results (0.04 sec)

My Citations

Articles

Eosinophilic meningoencephalitis due to Toxocara canis: case report and review of the literature[\[HTML\] from ajtmh.org](#)

Case law

JE Vidal, J Sztajn bok, [AC Seguro](#) - ... [journal of tropical medicine and hygiene](#), 2003 - [ASTMH](#)
 ... Although visceral larva migrans is a **characteristic** disease in children, CNS **toxocariasis** was seen ... **due to** T. canis is suggested by the presence of **clinical features**, peripheral eosinophilia ...
Clinical findings associated with eosinophilic pleocytosis and CSF reactivity on an ELISA ...
 Cited by 128 [Related articles](#) [All 9 versions](#) [Cite](#) [Save](#)

My library

Any time

Since 2016

Since 2015

Since 2012

Custom range...

Schistosoma mansoni: assessment of morbidity before and after control

JR Lambertucci, JC Serufo, R Gerspacher-Lara... - [Acta tropica](#), 2000 - [Elsevier](#)
 ... persistent infection and an evolving pathological process, **hepatic** enlargement is **clinically** recognized in ... The most common **characteristics** of the **clinical** syndrome are: (i) an indolent febrile ...
Clinical and radiological findings are similar to those associated with other **causes** of ...
 Cited by 143 [Related articles](#) [All 7 versions](#) [Cite](#) [Save](#)

Sort by relevance

Sort by date

Hepatic granulomas due to visceral larva migrans in adults: appearance on US and MRI

R Jain, S Sawhney, DK Bhargava, [SK Panda](#)... - [Abdominal ...](#), 1994 - [Springer](#)
 ... **Characteristic** sonographic and magnetic resonance imaging (MRI) **features** which have not previ ... The essential **features** of VLM are prolonged hyperecho- sinophilia (>30%) in children who eat ... 7-9]. Diagnosis is usually made on the basis of **clinical presentation**, eosinophilia, im ...
 Cited by 22 [Related articles](#) [All 5 versions](#) [Cite](#) [Save](#)

☒ include patents☒ include citations

全部

图片

新闻

视频

更多 ▾

搜索工具

找到约 752,000 条结果 (用时 0.91 秒)

Google 学术: Clinical characteristics and progression of liver abscess caused by toxocara

Toxocariasis: clinical aspects, epidemiology, medical ... - Despommier - 被引用次数: 563

Schistosoma mansoni: assessment of morbidity before ... - Lambertucci - 被引用次数: 144

Pyogenic abscesses and parasitic diseases - Lambertucci - 被引用次数: 48

Toxocariasis: Clinical Aspects, Epidemiology, Medical Ecology, and ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: D Despommier - 2003 - 被引用次数: 559 - 相关文章

Toxocariasis is caused by a series of related nematode species (ascarids) that routinely ... Infective larvae hatch after ingestion of eggs, but the juvenile stages fail to develop to ... The dominant clinical manifestations associated with toxocariasis are Human toxocariasis and pyogenic liver abscess: a possible association.

Eosinophilic Liver Abscess Caused by Toxocara Canis - ResearchGate

https://www.researchgate.net/.../51759769_Eosinophilic_Liver_Abscess_Cau... - 翻译此页

Eosinophilic Liver Abscess Caused by Toxocara Canis on ResearchGate, the ... Article: Clinical Parasitology ... Article: Eosinophilic granuloma of the liver: a characteristic lesion with ... Infection in humans, in contrast to their definitive hosts, remains occult, often resulting in disease caused by the migrating larval stages.

[Eosinophilic liver abscess caused by Toxocara canis]. - PubPDF

www.pubpdf.com/pub/.../Eosinophilic-liver-abscess-caused-by-Toxocara-ca... ▾ 翻译此页

Toxocariasis shows variable clinical manifestations including hepatomegaly, ... However a hepatic abscess caused by toxocara infection in adults is rare.