

Name of Journal: *World Journal of Hepatology*

ESPS Manuscript NO: 25063

Manuscript Type: Review

Multimodal brain monitoring in fulminant hepatic failure

Paschoal FM Jr *et al* Fulminant hepatic failure

Fernando Mendes Paschoal Jr, Ricardo de Carvalho Nogueira, Karla de Almeida Lins Ronconi, Marcelo de Lima Oliveira, Manoel Jacobsen Teixeira, Edson Bor-Seng-Shu

Abstract

Acute liver failure, also known as fulminant hepatic failure (FHF), embraces a spectrum of clinical entities characterized by acute liver injury, severe

Match Overview

| | | |
|---|---|----|
| 1 | CrossCheck 82 words Bor-Seng-Shu, Edson, Manoel Jacobsen Teixeira, Roberto Hirsch, Almir Ferreira de Andrade, and Raul Marino J | 2% |
| 2 | Internet 54 words crawled on 01-Apr-2016 www.omicsonline.org | 1% |
| 3 | CrossCheck 49 words Bo-Michael Bellander, "Consensus Meeting on Microdialysis in Neurointensive Care", Intensive Care Medicine, 1 | 1% |
| 4 | Internet 42 words crawled on 15-Sep-2015 www.ncbi.nlm.nih.gov | 1% |
| 5 | CrossCheck 32 words Katja Elfriede Wartenberg, "Future of brain support: multimodality monitoring", Future Neurology, 07/2006 | 1% |
| 6 | CrossCheck 30 words Textbook of Neurointensive Care, 2013. | 1% |
| 7 | Internet 27 words crawled on 25-Oct-2002 www.elsevier.com | 1% |

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 53,100 条结果 (用时 0.75 秒)

Google 学术: Multimodal Brain Monitoring in Fulminant Hepatic Failure

... and oxygen transport in fulminant hepatic failure - Harrison - 被引用次数: 490

Treatment of brain edema in acute liver failure - Rabinstein - 被引用次数: 22

... of brain edema in patients with acute liver failure - Larsen - 被引用次数: 34

Neuromonitoring in a porcine model of acute hepatic failure.

www.ncbi.nlm.nih.gov/pubmed/21508115 - 翻译此页

作者: NR Frühauf - 2011 - 被引用次数: 4 - 相关文章

2011年4月20日 - Therefore, in the present study, multimodal neuromonitoring was evaluated in an ... Acute liver failure was surgically induced in swine by complete hepatectomy (n = 8). Intracranial pressure monitoring via a ventricular drainage system, ... At autopsy, brain tissue of the animals was found to be swollen ...

Complications of intracranial pressure monitoring in ...

www.ncbi.nlm.nih.gov/pubmed/8093756 ▾ 翻译此页

作者: AT Blei - 1993 - 被引用次数: 208 - 相关文章

In patients with fulminant hepatic failure, brain oedema and the resulting intracranial hypertension often lead to death; intracranial pressure (ICP) monitoring may ...
缺少字词: multimodal

Oxford Textbook of Neurocritical Care - 第 433 页 - Google 图...

<https://books.google.com/books?isbn=0198739559> - 翻译此页

学术搜索

找到约 696 条结果 (用时0.05秒)

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☒ 包含引用

Improvement by acetylcysteine of hemodynamics and oxygen transport in fulminant hepatic failure

PM Harrison, JA Wendon, AES Gimson... - ... England Journal of ... , 1991 - Mass Medical Soc

... In all the patients a pulmonary-artery flotation catheter (7 French, Edwards Laboratories, Irvine, Calif.) and a radial arterial line had been inserted as part of the routine cardiovascular monitoring of patients with ... (2013) Fulminant Viral Hepatitis ... Metabolic Brain Disease 28, 7-10. 24. ...

被引用次数: 494 相关文章 所有 4 个版本 引用

Treatment of brain edema in acute liver failure

AA Rabinstein - Current treatment options in neurology, 2010 - Springer

... Multimodality brain monitoring can be achieved by adding the use of 1) jugular bulb catheters ... This therapy demands continuous electroencephalographic monitoring because the rate of barbiturate administration ... Detry O, De Roover A, Honore P, Meurisse M: Brain edema and ...

被引用次数: 22 相关文章 所有 7 个版本 引用

[HTML] Prevention and management of brain edema in patients with acute liver failure

FS Larsen, J Wendon - Liver transplantation, 2008 - Wiley Online Library

... Through the use of such multimodality monitoring of the brain, more accurate interventions can ... plasma tonicity; the future development of treatments that specifically antagonize brain edema will ... PJ, Silk DB, Strunin L, McCabe JJ, Williams R. Clinical monitoring of intracranial ...

被引用次数: 35 相关文章 所有 6 个版本 引用

wiley.com 中的 [HTML]

Subclinical seizure activity and prophylactic phenytoin infusion in acute liver failure: a controlled clinical trial

AJ Ellis, JA Wendon, R Williams - Hepatology, 2000 - Wiley Online Library

... a vicious circle of glutamate secretion, sei- zure activity, followed up by brain swelling with ... prophylactic anticon- vulsive therapy and the routine use of the CFAM in monitor- ing, it ... The measurement of JVO2 satn adds a further dimension to the multimodality monitoring that is so ...

被引用次数: 113 相关文章 所有 7 个版本 引用

wiley.com 中的 [PDF]



全部 图片 新闻 视频 更多 ▾ 搜索工具

找到约 15,400 条结果 (用时 0.45 秒)

Google 学术: Multimodal brain monitoring in fulminant hepatic failure

... and oxygen transport in fulminant hepatic failure - Harrison - 被引用次数: 494

Treatment of brain edema in acute liver failure - Rabinstein - 被引用次数: 22

... of brain edema in patients with acute liver failure - Larsen - 被引用次数: 35

Intracranial hypertension in acute liver failure

ndt.oxfordjournals.org/content/17/1/23.full ▾ 翻译此页

作者: D Richardson - 2002 - 被引用次数: 15 - 相关文章

Intracranial hypertension secondary to cerebral oedema reduces the cerebral ... Monitoring confers an advantage in acute liver failure in allowing early ..., the use of multimodality monitoring, elective ventilation, continuous ultrafiltration using ...

Neuromonitoring in a porcine model of acute hepatic failure.

www.ncbi.nlm.nih.gov/pubmed/21508115 - 翻译此页

作者: NR Frühauf - 2011 - 被引用次数: 5 - 相关文章

2011年4月20日 - Cerebral oedema has been noted to occur frequently in patients dying of fulminant hepatic failure. Therefore, in the present study, multimodal neuromonitoring was ... Acute liver failure was surgically induced in swine by complete ... Intracranial pressure monitoring via a ventricular drainage system, ...

Intracranial pressure monitoring and liver transplantation for fulminant ...

www.ncbi.nlm.nih.gov/pubmed/1618463 ▾ 翻译此页

作者: SD Lidofsky - 1992 - 被引用次数: 215 - 相关文章

Cerebral edema and intracranial hypertension, commonly present in ... pressure monitoring in comatose patients with fulminant hepatic failure as part of their ...