

Name of journal: *World Journal of Hepatology*

ESPS Manuscript NO: 25480

Manuscript type: ORIGINAL ARTICLE

Observational Study

Diagnostic non-invasive model of large risky esophageal varices in cirrhotic hepatitis C virus patients

Elalfy H *et al.* Diagnostic model and large esophageal varices

Hatem Elalfy, Walid Elsherbiny, Ashraf Abdel Rahman, Dina Elhammady, Shaker Wagih Shaltout, Ayman Z Elsamanoudy, Bassem El Deek

Abstract

AIM: To build a diagnostic non-invasive model for screening of large varices in cirrhotic hepatitis C virus (HCV) patients.

Match Overview

1	Internet 65 words crawled on 30-Sep-2015 jemds.com	1%
2	CrossCheck 23 words "The 20th Conference of the Asian Pacific Association ... r the Study of the Liver Poster Presentation", <i>Hepatology</i>	1%
3	Internet 22 words crawled on 04-Feb-2015 www.clinicalvirology.org	<1%
4	CrossCheck 20 words Zhang, Wei, Liqiong Wang, Lei Wang, Gang Li, Aoshuan g Huang, Ping Yin, Zhenhua Yang, Changquan Ling, an	<1%
5	CrossCheck 20 words Sun, Hye Young, Jeong Min Lee, Joon Koo Han, and Byu ng Ihn Choi. "Usefulness of MR elastography for predicti	<1%
6	Internet 19 words crawled on 12-Apr-2012 www.saudi gastro.com	<1%
7	Internet 15 words crawled on 29-Dec-2010 www.saudi gastro.com	<1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 13,000 条结果 (用时 1.00 秒)

Non-invasive Prediction of Esophageal Varices: Is It Possible?...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: KC Thomopoulos - 2011 - 被引用次数: 7 - [相关文章](#)

Also a substantial number of **patients** will not develop **large** varices during screening ... of **cirrhosis** was alcohol consumption (42.4%) followed by B or C **viral hepatitis** (25.3%). ... to endoscopy only in high-risk **patients** according to a predictive **model**. ... **Non-invasive diagnosis of esophageal varices** in chronic liver diseases.

Noninvasive Prediction of Large Esophageal Varices in Chroni...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: A Sarangapani - 2010 - 被引用次数: 55 - [相关文章](#)

Keywords: **Esophageal varices**, **non invasive** predictor of varices, platelet spleen ratio ...

The prevalence of varices in **patients** with **cirrhosis** is approximately 60-80% and the risk of ... [5] **Diagnosis of cirrhosis** was based on clinical, biochemical, and ... and antibodies to **hepatitis C virus** to determine the cause of liver **cirrhosis**.

[PDF] Detection of Risky Esophageal Varices by Two-Dimension...

[www.amjmedsci.com/article/S0002-9629\(15\)30433-X/pdf](http://www.amjmedsci.com/article/S0002-9629(15)30433-X/pdf) ▾ [翻译此页](#)

作者: H Mahfouz - 2014 - 被引用次数: 12 - [相关文章](#)

cirrhotic patients undergo screening endoscopy at **diagnosis** to identify **patients** with ...

Several studies have evaluated possible **noninvasive** predictors of esoph-

splenomegaly to a **huge** spleen with marked portal hypertension. ... **hepatitis C virus** (HCV) or hepatitis B virus (HBV) or a mixed Several predictive **models**..

学术搜索

找到约 115 条结果 (用时0.08秒)

文章

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

我的图书馆

[HTML] Prevalence and predictors of esophageal varices in patients with primary sclerosing cholangitis

wiley.com 中的 [HTML]

CO Zein, KD Lindor, P Angulo - Hepatology, 2004 - Wiley Online Library

... is not necessary in the standard diagnostic workup of patients with a diagnosis of PSC ... an endoscopy of the upper gastrointestinal tract will yield the most diagnostic information. ... moderate to large esophageal varices might improve the reliability of our proposed predictive model. ...

被引用次数: 92 相关文章 所有 5 个版本 引用 保存

时间不限

2016以来

2015以来

2012以来

自定义范围...

Assessment of the presence and severity of esophagogastric varices by splenic index in patients with liver cirrhosis

S Watanabe, N Hosomi, Y Kitade... - Journal of computer ..., 2000 - journals.lww.com

... In 53 patients (48.2%), the diagnosis of cirrhosis was based on the results of CT ... albumin, total bilirubin, serum cholinesterase, platelet count, γ -globulin, prothrombin time (DADE Diagnostics of Puerto ... The endoscopes used were models XP-10, GIF-20, GIF XQ-230, and GIF XQ ...

被引用次数: 29 相关文章 所有 5 个版本 引用 保存

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

... fibrosis scan for liver stiffness score measurement: a useful preendoscopic screening test for the detection of varices in postoperative patients with biliary atresia

researchgate.net 中的 [PDF]

HK Chang, YJ Park, H Koh, SM Kim... - Journal of pediatric ..., 2009 - journals.lww.com

... a body weight less than 5 kg were excluded from the endoscopic examination because of the large diameter (7 ... Diagnostic Accuracy of Liver Stiffness Measurement for Prediction of Varices. ...

We analyzed the ROC curves using the liver stiffness scores for the diagnosis of varices. ...

被引用次数: 54 相关文章 所有 8 个版本 引用 保存

☒ 包括专利☒ 包含引用☐ 创建快讯

[引用] The diagnosis and management of non-alcoholic fatty liver disease: practice guideline by the American Gastroenterological Association, American Association ...

N Chalasani, Z Younossi, JE Lavine, AM Diehl... - ..., 2012 - WB Saunders

被引用次数: 619 相关文章 所有 85 个版本 引用 保存

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 13,200 条结果 (用时 0.64 秒)

Non-invasive Prediction of Esophageal Varices: Is It Possible? - NCBI

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: KC Thomopoulos - 2011 - 被引用次数: 7 - [相关文章](#)

Also a substantial number of **patients** will not develop **large** varices during screening ... of **cirrhosis** was alcohol consumption (42.4%) followed by B or C **viral hepatitis** (25.3%). ... to endoscopy only in high-risk **patients** according to a predictive **model**. ... **Non-invasive diagnosis** of **esophageal varices** in chronic liver diseases.

Noninvasive Prediction of Large Esophageal Varices in Chronic Liver ...

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: A Sarangapani - 2010 - 被引用次数: 57 - [相关文章](#)

Keywords: **Esophageal varices**, **non invasive** predictor of varices, platelet spleen ratio ... The prevalence of varices in **patients** with **cirrhosis** is approximately 60-80% and the risk of ... [5] **Diagnosis** of **cirrhosis** was based on clinical, biochemical, and ... and antibodies to **hepatitis C virus** to determine the cause of liver **cirrhosis**.

HIV/Hepatitis C Virus–Coinfected Patients and Cirrhosis: How to ...

cid.oxfordjournals.org/content/58/6/840.full.pdf - [翻译此页](#)

作者: V Martel-Laferrière - 2014 - 被引用次数: 4 - [相关文章](#)

The **diagnosis** of early **cirrhosis** in HIV/**hepatitis C virus** (HCV)–coinfected pa- ... The development of **noninvasive** methods for fibrosis assessment empowers the in- fectious ... **patients** in screening programs for **esophageal varices** and hepatocellular ratio; Ln, Natural logarithm; MELD, **Model** for End-Stage Liver Disease.

A comparative study of Duplex Doppler ultrasound and blood indices ...

www.tdj.eg.net/article.asp?issn=1110-1415;year=2014;volume... ▾ [翻译此页](#)

作者: M Shehata - 2014 - [相关文章](#)

Endoscopic surveillance of **oesophageal varices** (OV) in **cirrhotic patients** is ... Therefore, there is a