

Name of journal: *World Journal of Hepatology*

ESPS Manuscript NO: 28178

Manuscript Type: Original Article

Case Control Study

Pancreatic hyperechogenicity associated with hypoadiponectinemia and insulin resistance: a Japanese population study

Naohiko Makino, Nakao Shirahata, Teiichiro Honda, Yoshiaki Ando, Akiko

Matsuda, Yushi Ikeda, Miho Ito, Yuko Nishise, Takafumi Saito, Yoshiyuki Ueno and

Sumio Kawata

Match Overview

1	Crossref 47 words Matsuda, Akiko, Naohiko Makino, Tomohiro Tozawa, Nakao Shirahata, Teiichiro Honda, Yushi Ikeda, Hideyuki Sat.	2%
2	Internet 32 words crawled on 02-Mar-2016 www.wjgnet.com	1%
3	Crossref 30 words Solomon, Sunil S., Aylur K. Srikrishnan, Allison M. McFall, M. Suresh Kumar, Shanmugam Saravanan, Pachamut	1%
4	Internet 26 words crawled on 19-Apr-2016 spandidos-publications.com	1%
5	Internet 25 words crawled on 12-Oct-2010 lib.bioinfo.pl	1%
6	Crossref 19 words Soro - Paavonen, Aino, Jukka Westerbacka, Christian Ehnholm, and Marja - Riitta Taskinen. "Metabolic syndro	1%

Pancreatic hyperechogenicity associated with hypoadiponectinemia and insulin r



全部 图片 视频 新闻 购物 地图 图书

找到约 316 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

Google 学术: [Pancreatic hyperechogenicity associated with hypoadiponectinemia and insulin resistance: a Japanese population study](#)

[Nonalcoholic fatty liver disease a feature of the ...](#) - Marchesini - 被引用次数: 2227

[... steatohepatitis? A preliminary human study](#) - Vuppalanchi - 被引用次数: 103

[Pediatric nonalcoholic fatty liver disease: a critical ...](#) - Patton - 被引用次数: 230

[Hypoadiponectinemia is associated with visceral fat accumulation ...](#)

www.ncbi.nlm.nih.gov/pubmed/14564678

... is associated with visceral fat accumulation and insulin resistance in Japanese men ... The aim of the present study was to investigate the association of serum ...

[Incidence of insulin resistance in obese subjects in a rural Japanese ...](#)

www.ncbi.nlm.nih.gov/pubmed/15642079

... in obese subjects in a rural Japanese population: the Tanno and Sobetsu study. ... Although it is well known that obesity is closely related to insulin resistance, ... In this study, we investigated the incidence of insulin resistance in citizens of ...

[Hypoadiponectinemia in obesity and type 2 diabetes: close ... - NCBI](#)

www.ncbi.nlm.nih.gov/pubmed/11344187

Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. ... were found to be decreased in Japanese individuals with obesity, type

Pancreatic hyperechogenicity associated with hypoadiponectinemia and insulin r



全部 图片 视频 新闻 购物 地图 图书

找到约 316 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

Google 学术: [Pancreatic hyperechogenicity associated with hypoadiponectinemia and insulin resistance: a Japanese population study](#)

[Nonalcoholic fatty liver disease a feature of the ...](#) - Marchesini - 被引用次数: 2227

[... steatohepatitis? A preliminary human study](#) - Vuppalanchi - 被引用次数: 103

[Pediatric nonalcoholic fatty liver disease: a critical ...](#) - Patton - 被引用次数: 230

[Hypoadiponectinemia is associated with visceral fat accumulation ...](#)

www.ncbi.nlm.nih.gov/pubmed/14564678

... is associated with visceral fat accumulation and insulin resistance in Japanese men ... The aim of the present study was to investigate the association of serum ...

[Incidence of insulin resistance in obese subjects in a rural Japanese ...](#)

www.ncbi.nlm.nih.gov/pubmed/15642079

... in obese subjects in a rural Japanese population: the Tanno and Sobetsu study. ... Although it is well known that obesity is closely related to insulin resistance, ... In this study, we investigated the incidence of insulin resistance in citizens of ...

[Hypoadiponectinemia in obesity and type 2 diabetes: close ... - NCBI](#)

www.ncbi.nlm.nih.gov/pubmed/11344187

Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. ... were found to be decreased in Japanese individuals with obesity, type

[全部](#)[新闻](#)[图片](#)[地图](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 277 条结果 (用时 0.59 秒)

Fatty pancreas, insulin resistance, and β -cell function: a population ...

www.ncbi.nlm.nih.gov/pubmed/24492753 ▾ [翻译此页](#)

作者: VW Wong - 2014 - 被引用次数: 31 - [相关文章](#)

2014年2月4日 - Fatty pancreas, insulin resistance, and β -cell function: a population study using ... Studies, The Chinese University of Hong Kong, Hong Kong, China. ... range of pancreatic fat and identify factors associated with fatty pancreas.

缺少字词: hyperechogenicity hypoadiponectinemia japanese

Association of birthweight with diabetes and insulin sensitivity or ...

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: J Oya - 2015 - 被引用次数: 2 - [相关文章](#)

2015年1月29日 - The aim of the present study was to assess the association between birth ... with diabetes and insulin resistance in the Japanese general population. was mediated through programming of the development of the pancreas, ...

Incidence of insulin resistance in obese subjects in a rural Japanese ...

www.ncbi.nlm.nih.gov/pubmed/15642079 ▾ [翻译此页](#)

作者: H Ohnishi - 2005 - 被引用次数: 35 - [相关文章](#)

... in obese subjects in a rural Japanese population: the Tanno and Sobetsu study. ... Although it is well known that obesity is closely related to insulin resistance, ... In this study, we investigated the incidence of insulin resistance in citizens of ...

缺少字词: pancreatic hyperechogenicity hypoadiponectinemia

Association of hypoadiponectinemia with non-alcoholic fatty liver ...

https://www.researchgate.net/.../46288547_Association_of_hypoadiponectin... - [翻译此页](#)

Studies in Indian population are lacking. ... Insulin resistance was calculated using Homeostasis Assessment model (HOMA-IR). Serum ... the association of hypoadiponectinemia with NAFLD in Asian Indians ... increased liver echogenicity with normal vessels and liver muscle and pancreatic beta