

Match Overview

- | Rank | Match Type | Words | Similarity |
|------|------------|---|------------|
| 1 | Crossref | 30 words
Udomsinprasert, Wanvisa, Sittisak Honsawek, Wilai Anom asiri, Voranush Chongsrisawat, Paisarn Vejchapipat, and | 1% |
| 2 | Crossref | 27 words
Sittisak Honsawek, "Serum adiponectin and transient elastography as non-invasive markers for postoperative biliary a | 1% |
| 3 | Crossref | 13 words
S. Honsawek, "Elevation of Serum Galectin-3 and Liver Stiffness Measured by Transient Elastography in Biliary Atresia | 1% |
| 4 | Crossref | 13 words
Honsawek, Sittisak, Pongsak Yuktanandana, Aree Tanaval ee, Chintana Chirathaworn, Wilai Anomasiri, Wanvisa Udo | 1% |
| 5 | Crossref | 13 words
Honsawek, Sittisak, Paisarn Vejchapipat, Sunchai Payungporn, Apiradee Theamboonlers, Voranush Chongsrisawat, | 1% |

Name of journal: *World Journal of Hepatology*

ESPS Manuscript NO: 27576

Manuscript Type: Original Article

Prospective Study

Elevation of serum urokinase plasminogen activator receptor and liver stiffness in postoperative biliary atresia

Wanvisa Udomsinprasert, Sittisak Honsawek, Napaphat Jirathanathornnukul, Voranush Chongsrisawat, Yong Poovorawan

Elevation of serum urokinase plasminogen activator receptor and liver stiffness

全部

图片

视频

新闻

购物

地图

图书

找到约 5,090 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Related Articles - PubMed Central Canada](#)

[pubmedcentralcanada.ca/pmc/solr/mlt.jsessionid...](#)

Results 1 - 25 ... Circulating soluble **urokinase plasminogen activator receptor** is stably **elevated** ... High suPAR **serum** concentrations are suggested to reflect the activation of the ... **Liver stiffness** measurement (LSM) by transient elastography (TE, and **liver stiffness** using transient elastography in **postoperative biliary atresia** ...

[Related Articles - PubMed Central Canada](#)

[pubmedcentralcanada.ca/pmc/solr/mlt.jsessionid...](#)

Results 1 - 25 ... Extrahepatic **biliary atresia**: Correlation of histopathology and liver function in the need for liver transplant in **postoperative** patients treated with a variety of the relationship of adiponectin with clinical parameters and **liver stiffness** scores. in the livers of **urokinase-type plasminogen activator**-NOG mice.

[Soluble receptor for advanced glycation end products and liver ...](#)

[europepmc.org/abstract/MED/23195134](#)

Abstract: OBJECTIVES: Soluble **receptor** for advanced glycation end products ... for advanced glycation end products and **liver stiffness in postoperative biliary atresia**. ... Serum sRAGE and **liver stiffness** values were significantly **elevated** in BA....

[Circulating soluble urokinase plasminogen activator is elevated in ...](#)

[www.ncbi.nlm.nih.gov/pubmed/22098627](#)

Elevation of serum urokinase plasminogen activator receptor and liver stiffness ir

全部

图片

视频

新闻

购物

地图

图书

找到约 5,180 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Elevation of serum urokinase plasminogen activator receptor and liver stiffness in postoperative biliary atresia](#)

[Liver fibrosis in biliary atresia](#) - Haafiz - 被引用次数: 38

[The matrix metalloproteinases and their inhibitors](#) - Murphy - 被引用次数: 650

[Hyaluronic acid as a non-invasive biomarker of liver ...](#) - Neuman - 被引用次数: 5

[Related Articles - PubMed Central Canada](#)

[pubmedcentralcanada.ca/pmcc/solr/mlt.jsessionid...](#)

Results 1 - 25 ... Circulating soluble urokinase plasminogen activator receptor is stably elevated ... High suPAR serum concentrations are suggested to reflect the activation of the ... Liver stiffness measurement (LSM) by transient elastography (TE, liver stiffness using transient elastography in postoperative biliary atresia ...

[Soluble urokinase plasminogen activator receptor is associated with...](#)

[www.ncbi.nlm.nih.gov/pubmed/21934527](#)

Soluble urokinase plasminogen activator receptor is associated with progressive liver ... BACKGROUND: Progressive liver fibrosis is the main predictor of disease ... of serum levels of the fibrinolysis-associated soluble urokinase plasminogen ...

[\[The plasma levels of urokinase plasminogen activator and ... - NCBI\]](#)

[www.ncbi.nlm.nih.gov/pubmed/14980106](#)

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 4,600 条结果 (用时 0.84 秒)

Google 学术: Elevation of serum urokinase plasminogen activator receptor and liver stiffness in postoperative biliary atresia

Liver fibrosis in biliary atresia - [Haafiz](#) - 被引用次数: 38

The matrix metalloproteinases and their inhibitors - [Murphy](#) - 被引用次数: 654

Hyaluronic acid as a non-invasive biomarker of liver ... - [Neuman](#) - 被引用次数: 5

Serum soluble urokinase plasminogen activator receptor as a ... - NCBI

www.ncbi.nlm.nih.gov/pubmed/25348952 ▾ 翻译此页

作者: A Chounta - 2015 - 被引用次数: 4 - 相关文章

Liver Int. 2015 Feb;35(2):601-7. doi: 10.1111/liv.12705. Epub 2014 Nov 20. Serum soluble urokinase plasminogen activator receptor as a screening test for the ...

缺少字词: elevation stiffness postoperative biliary atresia

Serum autotaxin levels correlate with hepatic dysfunction and severity ...

www.ncbi.nlm.nih.gov/pubmed/25536867 - 翻译此页

作者: W Udomsinprasert - 2015 - 被引用次数: 2 - 相关文章

2014年12月24日 - METHODS: Eighty postoperative BA patients and 15 controls were recruited. Serum ... Serum autotaxin and liver stiffness were markedly elevated in BA ... Biliary Atresia/blood*; Biliary Atresia/pathology; Biliary Atresia/surgery ...

缺少字词: urokinase plasminogen activator receptor

Serum adiponectin and transient elastography as non-invasive ...

bmcgastroenterol.biomedcentral.com/articles/10.1186/1471-230X-11-16 ▾ 翻译此页

作者: S Honsawek - 2011 - 被引用次数: 22 - 相关文章

Biliary atresia (BA) is a progressive inflammatory disorder of the extrahepatic bile ducts leading ...

Serum adiponectin and liver stiffness values were higher in BA patients. The elevated serum