

Name of Journal: *World Journal of Hepatology*

ESPS Manuscript NO: 27917

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Fractionation of gamma-glutamyltransferase in patients with nonalcoholic fatty liver disease and alcoholic liver disease

Sueyoshi S *et al.* GGT fractionation in liver diseases

Shigeo Sueyoshi, Setsu Sawai, Mamoru Satoh, Masanori Seimiya, Kazuyuki

Sogawa, Atsushi Fukumura, Mikihiro Tsutsumi, Fumio Nomura

Abstract

AIM

To assess how serum gamma-glutamyltransferase (GGT) fractions vary in patients with alcoholic liver disease (ALD) and non-alcoholic fatty liver disease

Match Overview

1	Internet 58 words crawled on 30-May-2016 etd.adm.unipi.it	2%
2	Crossref 27 words Immacolata Castellano . "γ-Glutamyltranspeptidases: sequence, structure, biochemical properties, and biotechnolo ...	1%
3	Internet 23 words crawled on 30-Apr-2016 spandidos-publications.com	1%
4	Internet 17 words crawled on 27-Apr-2016 www.clinchem.org	1%
5	Internet 16 words crawled on 23-Sep-2015 brenda-enzymes.org	1%
6	Crossref 13 words Franzini, M. . "A high performance gel filtration chromatogr ... phy method for @c-glutamyltransferase fraction analysis",	<1%

学术搜索

找到约 395 条结果 (用时0.07秒)

我的著作引用情况

文章

您是不是要找: Fractionation of gamma-*glutamyl transferase* in patients with nonalcoholic fatty liver disease and alcoholic liver disease

我的图书馆

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

时间不限

2016以来

2015以来

2012以来

自定义范围...

Fatty infiltration of liver in hyperlipidemic patients

N Assy, K Kaita, D Mymin, C Levy, B Rosser... - Digestive diseases and ..., 2000 - Springer
... A model that indepen- dently incorporated the different lipid fractions and then added ... alanine
aminotransferase (ALT) values were increased in 45 (47%), gamma-glutamyltransferase (GGT)
in ... CE: Relationship between 505 paired liver tests and biopsies in 242 obese patients. ...

被引用次数: 393 相关文章 所有 10 个版本 引用 保存

[PDF] academia.edu

按相关性排序

按日期排序

[HTML] Non-alcoholic fatty liver disease (NAFLD) and its connection with insulin
resistance, dyslipidemia, atherosclerosis and coronary heart disease

M Gaggini, M Morelli, E Buzzigoli, RA DeFronzo... - Nutrients, 2013 - mdpi.com

... profile, ie, increased small dense LDL fractions and reduced in HDL fractions [7,44 ... Risk of severe
liver disease in nonalcoholic fatty liver disease with normal aminotransferase levels: A role for ...
Yki-Jarvinen, H. Liver fat is increased in type 2 diabetic patients and underestimated ...

被引用次数: 186 相关文章 所有 14 个版本 引用 保存 更多

[HTML] mdpi.com

搜索所有网页

中文网页

简体中文网页

Hepatic steatosis and insulin resistance: does etiology make a difference?

A Lonardo, S Lombardini, F Scaglioni, L Carulli... - Journal of ..., 2006 - Elsevier

... following laboratory tests were performed: blood cells count, creatinine, serum total and
fractionated proteins, ALT ... 2. Histological samples: (a) FHBL (b) HCV (c) NAFLD patient. First,
considering our whole series of patients studied, liver steatosis, rather than histological grading ...

被引用次数: 126 相关文章 所有 8 个版本 引用 保存

☒ 包括专利☒ 包含引用

创建快讯

How different is the dietary pattern in non-alcoholic steatohepatitis patients?

H Cortez-Pinto, L Jesus, H Barros, C Lopes, MC Moura... - Clinical Nutrition, 2006 - Elsevier
... insulin ($\mu\text{U/L}$)/22.5; AST, aspartate aminotransferase; ALT, alanine aminotransferase; GGT,
gamma glutamyltransferase. was low to recommendations in 82%; 80.0% of patients exceeded
the ... Although no patient exceeded the recommended ingestion of polyunsaturated fat, our ...

被引用次数: 204 相关文章 所有 7 个版本 引用 保存

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 16,500 条结果 (用时 0.90 秒)

Google 学术: **Fractionation of gamma-glutamyltransferase in patients with nonalcoholic fatty liver disease and alcoholic liver disease**

Fatty infiltration of liver in hyperlipidemic patients - Assy - 被引用次数: 393

Non-alcoholic fatty liver disease (NAFLD) and its ... - Gaggini - 被引用次数: 186

Hepatic steatosis and insulin resistance: does etiology ... - Lonardo - 被引用次数: 126

Accuracy of b-GGT fraction for the diagnosis of non-alcoholic fatty liver ...

www.ncbi.nlm.nih.gov/pubmed/22098947 ▾ 翻译此页

作者: M Franzini - 2012 - 被引用次数: 28 - 相关文章

2011年11月4日 - Liver Int. 2012 Apr;32(4):629-34. doi: ... Accuracy of b-GGT fraction for the diagnosis of non-alcoholic fatty liver disease. ... METHODS: Gamma-glutamyltransferase fractions were determined in patients with NAFLD (n = 90), ...

A Study of Carotid Atherosclerosis in Patients with Non-alcoholic Fatty ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: S Mishra - 2013 - 被引用次数: 12 - 相关文章

2012年12月29日 - Non-alcoholic fatty liver disease shares many features of metabolic syndrome and its presence ... cholesterol and its fractions, insulin, alanine and aspartate transaminases, gamma glutamyl transferase and free fatty acids.

Gamma-Glutamyltransferase Fractions in Human Plasma and Bile ...

journals.plos.org/plosone/article?id=10.1371/journal.pone.0088532 ▾ 翻译此页

作者: L E ... - 2014 - 被引用次数: 8 - 相关文章

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 15,200 条结果 (用时 1.05 秒)

您是不是要找: **Fractionation of gamma-glutamyl transferase in patients with nonalcoholic fatty liver disease and alcoholic liver disease**

Google 学术: **Fractionation of gamma-glutamyltransferase in patients with nonalcoholic fatty liver disease and alcoholic liver disease**

Fatty infiltration of liver in hyperlipidemic patients - Assy - 被引用次数: 395

Non-alcoholic fatty liver disease (NAFLD) and its ... - Gaggini - 被引用次数: 193

Hepatic steatosis and insulin resistance: does etiology ... - Lonardo - 被引用次数: 126

Accuracy of b-GGT fraction for the diagnosis of non-alcoholic fatty liver ...

<https://www.ncbi.nlm.nih.gov/pubmed/22098947> - 翻译此页

作者: M Franzini - 2012 - 被引用次数: 28 - 相关文章

2011年11月4日 - Liver Int. 2012 Apr;32(4):629-34. doi: ... Accuracy of b-GGT fraction for the diagnosis of non-alcoholic fatty liver disease. ... METHODS: Gamma-glutamyltransferase fractions were determined in patients with NAFLD (n = 90), ...

Separation and identification of serum gamma-glutamyl ... - NCBI

www.ncbi.nlm.nih.gov/pubmed/7907377 - 翻译此页

作者: K Okuyama - 1993 - 被引用次数: 12 - 相关文章

Serum gamma-glutamyl transpeptidase (gamma-GTP) was separated into four ... those in patients with non-alcoholic liver cirrhosis (LC) and chronic hepatitis (CH). ... In alcoholic liver disease (ALD), no remarkable differences of the rate of each ... among alcoholic LC, alcoholic hepatitis, alcoholic CH and fatty liver patients.