

Name of Journal: *World Journal of Hepatology*

ESPS Manuscript NO: 28239

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Spontaneous bacterial peritonitis prevalence in pre-transplant patients and its effect on survival and graft loss post-transplant

Neeral L Shah, Nicolas M Intagliata, Zachary H Henry, Curtis K Argo, Patrick G Northup

Abstract

Aim: Investigate the incidence of SBP in pre-transplant patients and its effect on post-transplant mortality and graft failure.

Methods: We conducted a retrospective cohort study of patient records from the OPTN

data set. Patients were identified by the presence of spontaneous bacterial peritonitis

(SBP) pre-transplant. Univariate post-transplant survival models were constructed

using the Kaplan-Meier technique and multivariate models were constructed using the

Cox proportional hazards model. Variables that affected post-transplant graft survival

were identified in the SBP population.

Match Overview

Rank	Source	Words	Similarity
1	CrossCheck "Abstracts 416-623", <i>Hepatology</i> , 09/23/2009	123 words	4%
2	CrossCheck "AASLD Abstracts", <i>Hepatology</i> , 2013	88 words	3%
3	CrossCheck Jaillette, Emmanuelle, Christophe Girault, Guillaume Brunin, Farid Zerimech, Arnaud Chiche, Céline Brouqsault-De	45 words	2%
4	CrossCheck "WORLD TRANSPLANT CONGRESS 2006 POSTER ABSTRACTS", <i>American Journal of Transplantation</i> , 8/2006	31 words	1%
5	CrossCheck "American Transplant Congress 2007 Executive and Program Planning Committees and Abstract Review Committee	27 words	1%
6	CrossCheck "Abstracts Poster", <i>American Journal of Transplantation</i> , 5/2008	24 words	1%



全部 新闻 图片 视频 更多 ▾ 搜索工具

找到约 109,000 条结果 (用时 0.73 秒)

Etiology and mortality of spontaneous bacterial peritonitis in liver ...

onlinelibrary.wiley.com/doi/10.1002/lt.23889/pdf - 翻译此页

作者: C Pérez - Cameo - 2014 - 被引用次数: 2 - 相关文章

2014年4月3日 - Spontaneous bacterial peritonitis (SBP) in liver transplantation (LT) recipients who ... complications are common causes of late graft loss and have an impact on morbidity and mortality.1-4 ... risk factors, but data are lacking on the epidemiology of SBP in ... with 3CP in patients who develop cirrhosis after LT.

Peritonitis after liver transplantation: Incidence, risk factors ...

onlinelibrary.wiley.com/doi/10.1002/lt.20801/pdf ▾ 翻译此页

作者: S Pungpapong - 2006 - 被引用次数: 24 - 相关文章

950 cadaveric liver transplantation (LT) procedures in 837 patients, followed for a mean of ... PLT after their first LT had a significantly greater risk of graft loss or mortality ... intra-abdominal pathology and reduced recipient and graft survival. ... Abbreviations: SBP, spontaneous bacterial peritonitis; PLT, peritonitis after liver ...

Pre-transplant portal vein thrombosis is an independent risk factor for ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: JG Stine - 2016 - 相关文章

2015年12月10日 - Pre-transplant portal vein thrombosis is an independent risk factor for graft loss due to hepatic artery thrombosis in liver transplant recipients ... network; PVT: portal vein thrombosis; SBP: spontaneous bacterial peritonitis ... in determining and counseling patients regarding the risk of post-transplant HAT.

Primary biliary cirrhosis and liver transplantation

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: N Akamatsu - 2012 - 被引用次数: 1 - 相关文章

The highest incidence and prevalence rates were reported from the UK (19,20). ... Finally, recurrence of



[全部](#) [新闻](#) [图片](#) [视频](#) [更多 ▾](#) [搜索工具](#)

找到约 129,000 条结果 (用时 0.83 秒)

Infections in liver transplant recipients

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: FA Romero - 2011 - [被引用次数: 50](#) - [相关文章](#)

2011年4月27日 - [Survival after liver transplantation](#) has improved over the years, ... double the risk), [graft loss](#) (39% vs 7%), and [re-transplantation](#)[17]. ... and norfloxacin as prophylaxis for [spontaneous bacterial peritonitis](#), ... It is proposed that the indirect [effects](#) of CMV result from [its](#) immunomodulatory property[54-58].

Recurrence of hepatitis C after liver transplantation

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: C Vinaixa - 2013 - [被引用次数: 18](#) - [相关文章](#)

Recurrence of hepatitis C virus (HCV) infection [following](#) liver [transplantation](#) is a ... Reinfection of liver allografts is universal in [patients](#) with [pre-transplantation](#) ... [Graft failure](#) secondary to recurrent HCV represents the most frequent cause of ... Infection, particularly [spontaneous bacterial peritonitis](#), is a potentially severe ...

Pre-transplant portal vein thrombosis is an independent risk factor for ...

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: JG Stine - 2016 - [相关文章](#)

2015年12月10日 - [Pre-transplant](#) portal vein thrombosis is an independent risk factor for [graft loss](#) due to hepatic artery thrombosis in liver transplant recipients ... network; PVT: portal vein thrombosis; [SBP](#): [spontaneous bacterial peritonitis](#) ... in determining and counseling [patients](#) regarding the risk of [post-transplant](#) HAT.

Peritonitis after liver transplantation: Incidence, risk factors ...

onlinelibrary.wiley.com/doi/10.1002/lt.20801/pdf ▾ [翻译此页](#)

作者: S Pungpapong - 2006 - [被引用次数: 26](#) - [相关文章](#)



全部 新闻 图片 视频 更多 ▾ 搜索工具

找到约 104,000 条结果 (用时 0.94 秒)

Peritonitis after liver transplantation: Incidence, risk factors ...

onlinelibrary.wiley.com/doi/10.1002/lt.20801/pdf ▾ 翻译此页

作者: S Pungpapong - 2006 - 被引用次数: 26 - 相关文章

Peritonitis occurring after liver transplantation (PLT) has been poorly ... PLT after their first LT had a significantly greater risk of graft loss or mortality ... ed.2 Spontaneous bacterial peritonitis (SBP) is character- differences of the survival times between groups. Out- ... veloped in patients with a diagnosis of pre-LT refractory.

Infectious Complications After Liver Transplantation

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: MDP Hernandez - 2015 - 相关文章

Orthotopic liver transplantation (OLT) is the standard of care for patients with ... has improved graft and patient survival after orthotopic liver transplantation (OLT) but has also The effect of pretransplant infections on the incidence and severity of ... known that SBP is the most common infection among patients awaiting liver ...

Primary biliary cirrhosis and liver transplantation

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: N Akamatsu - 2012 - 被引用次数: 2 - 相关文章

The highest incidence and prevalence rates were reported from the UK (19,20), ... Finally, recurrence of PBC after liver transplantation supports the idea that bile duct ascites and spontaneous bacterial peritonitis, recurrent variceal bleeding, ... impact on patient or graft survival following liver transplantation in the short- ...

Medical Care of the Liver Transplant Patient

<https://books.google.com.hk/books?isbn=1444398423> - 翻译此页

Diogo Alain-Claudio, James E. Trotter, 2014, Medical