



28451-Review

Quotes Included
Bibliography Included3%
SIMILARName of Journal: *World Journal of Hepatology*

ESPS Manuscript NO: 28451

Manuscript Type: ORIGINAL ARTICLE

Observational Study


Prevalence of **significant liver disease in human immunodeficiency virus-infected patients exposed to Didanosine: A** cross sectional study

Sarah Logan, Alison Rodger, Laura Maynard-Smith, James O'Beirne, Thomas Fernandez, Filippo Ferro, Colette Smith, Sanjay Bhagani

Abstract**AIM**

We aimed to identify significant liver disease (including Nodular regenerative hyperplasia [NRH]) in asymptomatic Didanosine (DDI) exposed **human immunodeficiency virus** (HIV) positive patients.

Match Overview



- | | | |
|---|--|-----|
| 1 | Crossref 50 words
"Poster abstracts", HIV Medicine, 04/2011 | 2% |
| 2 | Internet 21 words
crawled on 21-Apr-2012
bhiya.org | 1% |
| 3 | Internet 11 words
crawled on 03-Nov-2015
www.science.gov | <1% |

[全部](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 39,700 条结果 (用时 0.34 秒)

Google 学术: Prevalence of significant liver disease in HIV-infected patients exposed to Didanosine: a cross sectional study

... fatty liver disease in HIV-infected patients referred to a ... - Guaraldi - 被引用次数: 118

Liver cirrhosis in HIV - infected patients: prevalence, ... - Castellares - 被引用次数: 99

... fatty liver disease (NAFLD) among HIV-infected ... - Crum-Cianflone - 被引用次数: 107

小提示: 仅限搜索简体中文结果。您可以在设置中指定搜索语言

Prevalence and risk factors for abnormal liver stiffness in HIV-infected ...

www.ncbi.nlm.nih.gov/pubmed/20710057 - [翻译此页](#)

作者: N Merchante - 2010 - 被引用次数: 33 - [相关文章](#)

Prevalence and risk factors for abnormal liver stiffness in HIV-infected patients without viral hepatitis coinfection: role of didanosine. Merchante N(1) ... METHODS: A cross-sectional prospective study was conducted. LS was assessed by TE in ... Long-term exposure to ddl is a major cause of liver damage in these patients.

Prevalence and risk factors for significant liver fibrosis among HIV ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ [翻译此页](#)

作者: M DallaPiazza - 2010 - 被引用次数: 55 - [相关文章](#)

2010年5月13日 - We conducted a cross-sectional study among HIV-infected patients ... risk factors associated with liver disease progression, hepatic fibrosis, In this cross-sectional study, we were not able to assess the duration of exposure to ART, ... older non-nucleoside (i.e., nevirapine) and nucleoside (i.e., didanosine, ...

[PDF] Original article Prevalence and risk factors for abnormal liver stiffnes...

<https://www.intmedpress.com/serveFile.cfm?slIID=dd6711c2-8299-445a> - [翻译此页](#)

[全部](#)[新闻](#)[图片](#)[地图](#)[更多 ▾](#)[搜索工具](#)

找到约 82,700 条结果 (用时 0.66 秒)

Human immunodeficiency virus infection and the liver - NCBI

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ [翻译此页](#)

作者: M Crane - 2012 - 被引用次数: 40 - 相关文章

2012年3月27日 - Liver disease in human immunodeficiency virus (HIV)-infected individuals ... of liver disease in HIV-infected patients, in the absence of co-infection with A strong association between prolonged exposure to didanosine (ddl) and the ... Two cross sectional studies looking at associations between multiple ...

Prevalence and risk factors for significant liver fibrosis among HIV ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ [翻译此页](#)

作者: M DallaPiazza - 2010 - 被引用次数: 58 - 相关文章

2010年5月13日 - We conducted a cross-sectional study among HIV-infected patients APRI accurately represents liver fibrosis only when liver disease has In this cross-sectional study, we were not able to assess the duration of exposure to ART. older non-nucleoside (i.e., nevirapine) and nucleoside (i.e., didanosine, ...

High Prevalence of Liver Fibrosis Associated with HIV Infection: A ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ [翻译此页](#)

作者: L Stabinski - 2011 - 被引用次数: 46 - 相关文章

Liver disease is a leading cause of mortality among HIV-infected persons in ... The prevalence of significant fibrosis was 17% among HIV-infected and 11% in ... This cross-sectional study was conducted in the Rakai District of rural ... 500 unselected HIV-infected patients of the Rakai Health Sciences Program (RHSP) HIV ...

[PDF] Epidemiology of Liver Disease in Human Immunodeficiency Virus ...

www.springer.com/cda/content/document/.../9781441917119-c1.pdf?... ▾ [翻译此页](#)

Liver disease in HIV-infected persons may be due to the virus itself ... moderate elevations do not seem

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多 ▾](#)[搜索工具](#)

找到约 82,100 条结果 (用时 1.04 秒)

Human immunodeficiency virus infection and the liver - NCBI

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: M Crane - 2012 - 被引用次数: 41 - 相关文章

跳到CLINICAL PRESENTATIONS OF **LIVER DISEASE IN HIV** MONO ... - ... exposure to didanosine (ddl) and the ... Two cross sectional studies ... Fatty liver disease in HIV ... the prevalence of NAFLD was ... Insulin resistance and exposure to ... in one study of 77 patients[62]. ... Significant fibrosis (\geq F2, liver ...

Prevalence and risk factors for significant liver fibrosis among HIV ...

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: M DallaPiazza - 2010 - 被引用次数: 61 - 相关文章

2010年5月13日 - We conducted a cross-sectional study among HIV-infected patients APRI accurately represents liver fibrosis only when liver disease has In this cross-sectional study, we were not able to assess the duration of exposure to ART. ... older non-nucleoside (i.e., nevirapine) and nucleoside (i.e., didanosine, ...

Assessment of liver fibrosis and associated risk factors in HIV-infected ...

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: J Vermehren - 2012 - 被引用次数: 17 - 相关文章

2012年3月27日 - Liver fibrosis in human immunodeficiency virus (HIV)-infected individuals is ... of other risk factors, including prolonged exposure to combined antiretroviral ... Overall, factors independently associated with significant fibrosis as assessed by ... for each individual patient at baseline in the cross-sectional study.

[PDF] Human immunodeficiency virus infection and the liver

www.wjnet.com/esps/DownloadFile.aspx?Type=Digital...1... ▾ [翻译此页](#)

作者: M Crane - 2012 - 被引用次数: 41 - 相关文章

HIV infection and the liver liver disease in HIV-infected patients with a specific focus ... EPIDEMIOLOGY OF LIVER DISEASE ... between prolonged exposure to didanosine (ddl) and ...