

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

 包括专利 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 [学术搜索设置](#) 指定搜索语言**Hepcidin Is Down - regulated in Alcoholic Liver Injury: Implications for the Pathogenesis of Alcoholic Liver Disease**KR Bridle, TK Cheung, TL Murphy... - *Alcoholism: Clinical ...*, 2006 - Wiley Online Library... Nicolas G, Bennoun M, Devaux I, Beaumont C, Grandchamp B, Kahn A, Vaulont S (2001) Lack of hepcidin gene expression and severe tissue iron overload in upstream stimulatory factor 2 (USF2) knockout mice. *Proc Natl Acad Sci USA* 98: 8780-8785. ...被引用次数: 113 [相关文章](#) [所有 7 个版本](#) [引用](#) [保存](#)**[HTML] Female hepatology: favorable role of estrogen in chronic liver disease with hepatitis B virus infection.** europepmc.org 中的 [\[HTML\]](#)I Shimizu, N Kohno, K Tamaki, M Shono... - *World journal of ...*, 2007 - europepmc.org

... These findings indicate that alcohol-induced oxidative stress downregulates hepcidin transcription, leading ... estrogen levels in menopausal women are associated with a loss in subcutaneous ... becomes evident spontaneously in aromatase-deficient mouse, which lack the ability ...

被引用次数: 93 [相关文章](#) [所有 13 个版本](#) [引用](#) [保存](#)**[HTML] Altered lipid metabolism in Hfe-knockout mice promotes severe NAFLD and early fibrosis** physiology.org 中的 [\[HTML\]](#)

TCH Tan, DHG Crawford... - ... and Liver ... , 2011 - Am Physiological Soc

... HFE interacts with cell surface transferrin receptors to modulate cellular iron uptake and signal the control of the expression of hepcidin, the master ... Here, we tested the hypothesis that the loss of HFE functioning plays a role in the development of NAFLD, by examining ...

被引用次数: 18 [相关文章](#) [所有 7 个版本](#) [引用](#) [保存](#)**Iron metabolism in nonalcoholic fatty liver disease**JE Nelson, H Klintworth, KV Kowdley - *Current gastroenterology reports*, 2012 - Springer

... and decreased platelets compared to the other iron deposition patterns or a lack of iron ... Diabetics

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 21933

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Lack of hepcidin expression attenuates steatosis and causes fibrosis in the liver

Sizhao Lu, Robert G Bennett, Kusum Kharbanda, Duygu Dee Harrison-Findik

Abstract

AIM: To investigate the role of key iron-regulatory protein, hepcidin in non-alcoholic fatty liver disease (NAFLD).

METHODS: Hepcidin (*Hamp1*) knockout and floxed control mice were administered a high fat and high sucrose (HES) or a regular control diet for 3 or 7 months. Steatosis

Match Overview

- 1** **CrossCheck** 27 words
Ann K Daly. "Genetic determinants of susceptibility and severity in nonalcoholic fatty liver disease". Expert Review of G <1%
e, and S. Dooley. "Animal Models of Chronic Liver Disease
- 2** **CrossCheck** 25 words
Liu, Y., C. Meyer, C. Xu, H. Weng, C. Hellerbrand, P. ten D ... <1%
e, and S. Dooley. "Animal Models of Chronic Liver Disease
- 3** **CrossCheck** 12 words
Cauchy, François. "Metabolic syndrome and non-alcoholi ... <1%
atty liver disease in liver surgery: The new scourges?", Wor

Lack of hepcidin expression attenuates steatosis and causes fibrosis in the liver



网页 图片 视频 新闻 购物 地图 图书

找到约 18,300 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Google 学术: Lack of hepcidin expression attenuates steatosis and causes fibrosis in the liver](#)

- [Hepcidin Is Down-regulated in Alcoholic Liver Injury: ...](#) - Bridle - 被引用次数: 118
- [... role of estrogen in chronic liver disease with hepatitis ...](#) - Shimizu - 被引用次数: 98
- [... mice promotes severe NAFLD and early fibrosis](#) - Tan - 被引用次数: 18

[New advances in molecular mechanisms and emerging therapeutic ...](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC4177473/ ▾

28 Sep 2014 ... In most heavy alcohol consumers, **steatosis** is **caused** by inhibiting fatty acid ... elevation in serum ALT levels, **steatosis** and some mild **liver fibrotic** changes[43], **attenuated** ethanol-induced mitochondrial damage in the **liver**. and a **lack of hepcidin** gene **expression causes** iron overload in tissues, ...

[Alcohol Activates TGF-Beta but Inhibits BMP Receptor-Mediated ...](#)

www.hindawi.com/journals/ijh/2012/459278/ ▾

7 Aug 2011 ... Alcohol induced **steatosis** and TGF-beta expression. **Liver** ..., Alcohol downregulates **hepcidin** expression in the **liver**, which leads to an The deletion of Smad4 in the **liver** also **attenuates** **hepcidin** expression and **causes** pronounced ... involved in the progression of **liver fibrosis** in alcoholic **liver** disease.

[High Fat Diet Induces Liver Steatosis and Early Dysregulation of Iron ...](#)

journals.plos.org/plosone/article?id=10.1371/journal.pone... ▾

21 Jun 2013 ... In **liver** of HFD fed animals an increased time-dependent activity of iron regulatory ... In the **liver** the **expression** of **hepcidin** is increased dramatically in ... of ferritin level observed at 5 weeks, that results **attenuated** after 8 weeks. weeks HFD, **steatosis** and mild inflammation without a clear **fibrosis**) and to ...

[\[PDF\] PDF\(403K\) - Wiley Online Library](#)

onlinelibrary.wiley.com/doi/10.1111/j.1522-0277.2009.01015.x/pdf



ack of hepcidin expression attenuates steatosis and causes fibrosis in the liver



[网页](#) [图片](#) [新闻](#) [购物](#) [视频](#) [更多](#) [搜索工具](#)

找到约 15,300 条结果 (用时 0.51 秒)

Google 学术 : Lack of hepcidin expression attenuates steatosis and causes fibrosis in the liver

[Hepcidin Is Down-regulated in Alcoholic Liver Injury: ...](#) - [Bridle](#) - 被引用次数 : 119

[... role of estrogen in chronic liver disease with hepatitis ...](#) - [Shimizu](#) - 被引用次数 : 98

[... mice promotes severe NAFLD and early fibrosis](#) - [Tan](#) - 被引用次数 : 18

New advances in molecular mechanisms and emerging ...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▼ [翻译此页](#)

作者 : [JA Williams](#) - 2014 - 被引用次数 : 12 - [相关文章](#)

2014年9月28日 - In most heavy alcohol consumers, **steatosis** is **caused** by inhibiting fatty acid ... elevation in serum ALT levels, **steatosis** and some mild **liver fibrotic** changes[43]. **attenuated** ethanol-induced mitochondrial damage in the **liver**. and a **lack of hepcidin** gene **expression causes** iron overload in tissues, ...

High Fat Diet Induces Liver Steatosis and Early ...

journals.plos.org/plosone/article?id=10.1371/journal.pone... ▼ [翻译此页](#)

作者 : [R Meli](#) - 2013 - 被引用次数 : 18 - [相关文章](#)

2013年6月21日 - In **liver** of HFD fed animals an increased time-dependent activity of iron regulatory ... In the **liver** the **expression** of **hepcidin** is increased dramatically in ... of ferritin level observed at 5 weeks, that results **attenuated** after 8 weeks. weeks HFD, **steatosis** and mild inflammation without a clear **fibrosis**) and to ...

Alcohol Activates TGF-Beta but Inhibits BMP Receptor ...

www.hindawi.com/journals/ijh/2012/459278/ ▼ [翻译此页](#)

作者 : [LN Gerjevic](#) - 2011 - 被引用次数 : 24 - [相关文章](#)

2011年8月7日 - Alcohol induced **steatosis** and TGF-beta expression. **Liver** ... Alcohol