

Name of Journal: *World Journal of Hepatology*

ESPS Manuscript NO: 29817

Manuscript Type: Case Report

**18-Fluoro-deoxyglucose uptake in inflammatory hepatic adenoma: A case report**

**Willy Liu, Jean Delwaide, Noella Bletard, Philippe Delvenne, Paul Meunier, Roland Hustinx, Olivier Detry**

#### Abstract

Positron emission tomography computed tomography (PET-CT) using 18-Fluoro-deoxyglucose (<sup>18</sup>FDG) is an imaging modality that reflects cellular glucose metabolism. Most cancers show an uptake of <sup>18</sup>FDG and benign tumors do not usually behave in such a way. The authors report herein the

#### Match Overview

Rank	Source	Words	Crawled On	Similarity (%)
1	Internet	72 words	crawled on 23-Feb-2017	5%
	www.wjgnet.com			
2	Crossref	35 words	Nakashima, Takaaki, Yukihisa Takayama, Akihiro Nishie, Yoshiki Asayama, Shingo Baba, Youichi Yamashita, Ken	2%
3	Internet	21 words	crawled on 08-Jan-2014	1%
	www.ncbi.nlm.nih.gov			
4	Crossref	18 words	Tateishi, Yoko, Mitsuiko Furuya, Fukuo Kondo, Ikue Torii, Kazunori Nojiri, Yukichi Tanaka, Shigeaki Umeda, Koji O	1%
5	Internet	16 words	crawled on 04-May-2016	1%
	www.authorstream.com			
6	Internet	14 words	crawled on 11-Jun-2013	1%
	radiographics.rsna.org			
7	Internet	12 words	crawled on 22-Jun-2015	1%
	www.ncbi.nlm.nih.gov			



全部 图片 新闻 视频 更多 ▾ 搜索工具

找到约 41,700 条结果 (用时 0.31 秒)

### Google 学术: 18-Fluorodeoxyglucose uptake in inflammatory hepatic adenoma: a case report

Increased 18 F-fluorodeoxyglucose uptake in benign, ... - Metser - 被引用次数: 132

Incidental Colonic Fluorodeoxyglucose Uptake: ... - Tatlidil - 被引用次数: 197

... -18 fluorodeoxyglucose positron emission tomography ... - Zimny - 被引用次数: 245

### Hepatic adenoma mimicking a metastatic lesion on computed ... - NCBI

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: D Lim - 2013 - 被引用次数: 7 - 相关文章

2013年7月21日 - Positron emission tomography (PET) using <sup>18</sup>F-fluorodeoxyglucose ... We report the case of a 44-year-old Chinese female patient with a history of .... In infections or inflammation, increased FDG uptake in lesions can be ...

### [PDF] Increased (18)F-fluorodeoxyglucose uptake in benign, nonphysiologi...

[medlib.yu.ac.kr/eur\\_j\\_oph/se\\_n\\_u/s\\_n\\_u/37\\_3\\_206.pdf](http://medlib.yu.ac.kr/eur_j_oph/se_n_u/s_n_u/37_3_206.pdf) ▾ 翻译此页

作者: U Metser - 被引用次数: 132 - 相关文章

the reports of PET/computed tomography (CT) studies performed in oncologic patients during a 6-month .... Increased uptake of 18F-FDG in inflammation may be explained by ..... as is the case in amebic hepatic abscess,<sup>33</sup> or from sepsis via the hepatic ..... adenoma. Spleen. Gaucher's disease. Adrenal. Adenoma. Benign.

### 'False-Positive' Uptake of FDG in a Hepatic Adenoma - ResearchGate

[https://www.researchgate.net/.../13991543\\_'False-Positive'\\_Uptake\\_of\\_FDG\\_...](https://www.researchgate.net/.../13991543_'False-Positive'_Uptake_of_FDG_...) - 翻译此页

Uptake of 18 FDG by a Hepatic Adenoma on Positron Emission Tomography ... the importance and implications of a false-positive result, as in this case report.

### A case of multiple hepatic angiomyolipomas with high 18 F ...

[bmcmedimaging.biomedcentral.com/articles/10.1186/1471-2342-14-17](http://bmcmedimaging.biomedcentral.com/articles/10.1186/1471-2342-14-17) ▾ 翻译此页

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 97,900 条结果 (用时 0.86 秒)

您是不是要找：[18-Fluorodeoxyglucose uptake in inflammatory hepatic adenoma: A case report](#)

Google 学术：[18-Fluoro-deoxyglucose uptake in inflammatory hepatic adenoma: A case report](#)

... of pancreatic carcinoma: a report of 106 cases - Zimny - 被引用次数 : 247

Metabolic characterization of breast tumors with ... - Avril - 被引用次数 : 288

... 'disease measured by F-18-fluoro-deoxyglucose ... - Boerner - 被引用次数 : 53

[Hepatic adenoma mimicking a metastatic lesion on computed ... - NCBI](#)  
<https://www.ncbi.nlm.nih.gov> , NCBI , Literature , PubMed Central (PMC) - 翻译此页

作者 : D Lim - 2013 - 被引用次数 : 7 - 相关文章

2013年7月21日 - Positron emission tomography (PET) using <sup>18</sup>F-fluorodeoxyglucose (<sup>18</sup>F-FDG) is an ... We report the case of a 44-year-old Chinese female patient with a history of ... lesion (standardized uptake value 7.9) in segment IVb of the liver. .... In infections or inflammation, increased FDG uptake in lesions can be ...

[PET-avid hepatocellular adenomas: incidental findings ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov> , NCBI , Literature , PubMed Central (PMC) - 翻译此页

作者 : SY Lee - 2016 - 被引用次数 : 1 - 相关文章

2015年12月11日 - Hepatocellular adenoma (HCA) is the second most common benign liver neoplasm and ... Positron emission tomography (PET) using [18F] fluorodeoxyglucose (FDG) is ... adenomas, Hepatocellular adenomas, False positive, inflammatory, ... reports of 10 patients with FDG-avid HCA, including a small case ...

[The clinical use of PET with <sup>11</sup>C-acetate](#)

<https://www.ncbi.nlm.nih.gov> , NCBI , Literature , PubMed Central (PMC) - 翻译此页

全部

图片

新闻

视频

购物

更多

设置

工具

找到约 104,000 条结果 (用时 0.81 秒)

### Google 学术: 18-Fluoro-deoxyglucose uptake in inflammatory hepatic adenoma: A case report

... of pancreatic carcinoma: a report of 106 cases - Zimny - 被引用次数: 247

Metabolic characterization of breast tumors with ... - Avril - 被引用次数: 288

... 'disease measured by F-18-fluoro-deoxyglucose ... - Boerner - 被引用次数: 53

### Hepatic adenoma mimicking a metastatic lesion on computed ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: D Lim - 2013 - 被引用次数: 7 - 相关文章

2013年7月21日 - Positron emission tomography (PET) using <sup>18</sup>F-fluorodeoxyglucose (<sup>18</sup>F-FDG) is an ...

We report the case of a 44-year-old Chinese female patient with a history of ... lesion (standardized uptake value 7.9) in segment IVb of the liver. .... In infections or inflammation, increased FDG uptake in lesions can be ...

### PET-avid hepatocellular adenomas: incidental findings associated ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: SY Lee - 2016 - 被引用次数: 1 - 相关文章

2015年12月11日 - Hepatocellular adenoma (HCA) is the second most common benign liver neoplasm and ... Positron emission tomography (PET) using [18F] fluorodeoxyglucose (FDG) is ... adenomas, Hepatocellular adenomas, False positive, inflammatory, ... reports of 10 patients with FDG-avid HCA, including a small case ...

### A case of multiple hepatic angiomyolipomas with high 18 F ...

[www.ncbi.nlm.nih.gov](https://www.ncbi.nlm.nih.gov) > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: S Kumasaka - 2014 - 被引用次数: 3 - 相关文章

2014年5月20日 - Although several case reports of hepatic angiomyolipoma have been ... In this case,