



Name of Journal: *World Journal of Hepatology*

ESPS Manuscript NO: 29879

Manuscript Type: MINIREVIEWS

Iatrogenic amyloid polyneuropathy after domino liver transplantation

Diana Mnatsakanova, Saša A Živković

Match Overview

1	Internet 16 words crawled on 11-Dec-2007 www.forbes.com	1%
2	Crossref 13 words Waddington Cruz, Márcia, and Merrill D. Benson. "A Review ... of Tafamidis for the Treatment of Transthyretin-Related Am	1%
3	Crossref 13 words Adams, David, Marie Théaudin, Cecile Cauquil, Vincent Algarrondo, and Michel Slama. "FAP Neuropathy and Emei ...	1%
4	Crossref 13 words Harit V. Desai. "Cardiac Amyloidosis : Approaches to Diagnosis and Management", <i>Cardiology in Review</i> , 01/2010	1%
5	Crossref 10 words Ando, Y., and M. Ueda. "Diagnosis and Therapeutic Approaches to Transthyretin Amyloidosis", <i>Current Medicinal Che</i>	1%
6	Internet 10 words crawled on 05-Nov-2015 www.science.gov	1%

学术搜索

获得 102 条结果 (用时0.13秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[HTML] iatrogenic amyloid neuropathy in a Japanese patient after sequential liver transplantation

T Goto, T Yamashita, M Ueda... - ... of transplantation, 2006 - Wiley Online Library

... Our findings confirmed iatrogenic sensory neuropathy induced by amyloid deposition 7 years ... Research Committee; the Pathogenesis, Therapy of Hereditary Neuropathy Research Committee ... through the blood-nerve barrier in recipients of familial amyloid polyneuropathy livers. ...

被引用次数: 54 相关文章 所有 9 个版本 引用 保存

Clinical symptomatic de novo systemic transthyretin amyloidosis 9 years after domino liver transplantation

AP Barreiros, C Geber, F Birklein, PR Galle... - ... Transplantation, 2010 - Wiley Online Library

... In order to alleviate the graft shortage, livers from patients with familial amyloidotic polyneuropathy (FAP) are used as domino grafts for other patients under the assumption that several decades ... iatrogenic amyloid neuropathy in a Japanese patient after sequential liver ...

被引用次数: 23 相关文章 所有 6 个版本 引用 保存

[HTML] Risk of transmission of systemic transthyretin amyloidosis after domino liver transplantation

L Lladó, C Baliellas, C Casasnovas, I Ferrer... - ... Transplantation, 2010 - Wiley Online Library

... minimal role for diabetes in the development of the severe polyneuropathy associated with ... biopsy samples will develop clinical symptoms and EMG lesions consistent with amyloid neuropathy. ... eg, rejection and infection) may promote the production of amyloid and accelerate ...

被引用次数: 47 相关文章 所有 5 个版本 引用 保存

Symptomatic and proven de novo amyloid polyneuropathy in familial amyloid polyneuropathy domino liver recipients

D Adams, C Lacomis, T Antonini, B Lazzaro, C Denier... Amyloid, 2011 Taylor & Francis

学术搜索

获得 109 条结果 (用时0.11秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 [学术搜索设置](#) 指定搜索语言

[HTML] [iatrogenic amyloid neuropathy in a Japanese patient after sequential liver transplantation](#)

T Goto, T Yamashita, M Ueda... - ... of [transplantation](#), 2006 - Wiley Online Library

... Our findings confirmed [iatrogenic](#) sensory [neuropathy](#) induced by [amyloid](#) deposition 7 years after sequential liver transplantation. Abbreviations: FAP familial [amyloidotic polyneuropathy](#), TTR transthyretin. ATTR ... [amyloidotic polyneuropathy](#) (FAP). [Amyloid](#) 2003; 10: S55-S66. ...

被引用次数: 54 [相关文章](#) 所有 9 个版本 [引用](#) [保存](#)

[HTML] [Risk of transmission of systemic transthyretin amyloidosis after domino liver transplantation](#)

L Lladó, C Baliellas, C Casasnovas, I Ferrer... - ... [Transplantation](#), 2010 - Wiley Online Library

... a fairly minimal role for diabetes in the development of the severe [polyneuropathy](#) associated with ... related to [transplantation](#) have also been implicated in the early development of [amyloidotic](#) changes in ... First, the presence of [amyloid](#) fibers in the graft may trigger the disease.27 ...

被引用次数: 47 [相关文章](#) 所有 5 个版本 [引用](#) [保存](#)

[Clinical symptomatic de novo systemic transthyretin amyloidosis 9 years after domino liver transplantation](#)

AP Barreiros, C Geber, F Birklein, PR Galle... - ... [Transplantation](#), 2010 - Wiley Online Library

... In order to alleviate the graft shortage, [livers](#) from patients with familial [amyloidotic polyneuropathy](#) (FAP) are used as [domino](#) grafts for other patients under the assumption that several decades ... [iatrogenic amyloid neuropathy](#) in a Japanese patient after sequential liver ...

被引用次数: 22 [相关文章](#) 所有 6 个版本 [引用](#) [保存](#)

[Symptomatic and proven de novo amyloid polyneuropathy in familial amyloid polyneuropathy domino liver recipients](#)

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 3,840 条结果 (用时 0.61 秒)

Google 学术: Iatrogenic amyloid polyneuropathy after domino liver transplantation

Iatrogenic amyloid neuropathy in a Japanese patient ... - Goto - 被引用次数: 54

... amyloidosis after domino liver transplantation - Lladó - 被引用次数: 49

... amyloidosis 9 years after domino liver transplantation - Barreiros - 被引用次数: 22

Iatrogenic Amyloid Neuropathy in a Japanese ... - Wiley Online Library

onlinelibrary.wiley.com/doi/10.1111/j.1600-6143.2006.01484.x/pdf ▾ 翻译此页

作者: T Goto - 2006 - 被引用次数: 54 - 相关文章

Our findings confirmed iatrogenic sensory neuropathy induced by amyloid deposition 7 years after sequential liver transplantation. Key words: Amyloid, domino ...

Iatrogenic Amyloid Neuropathy in a Japanese Patient After Sequential ...

onlinelibrary.wiley.com/doi/10.1111/j.1600-6143.2006.01484.x/full - 翻译此页

作者: T Goto - 2006 - 被引用次数: 54 - 相关文章

2006年8月1日 - Iatrogenic Amyloid Neuropathy in a Japanese Patient After Sequential Domino liver transplantation is normally performed for patients with ...

Domino transplantation using livers from patients with familial ...

onlinelibrary.wiley.com/doi/10.1002/lt.21073/pdf - 翻译此页

作者: BG Ericzon - 2007 - 被引用次数: 17 - 相关文章

Familial amyloidotic polyneuropathy (FAP) is a neuro- degenerative ... TTR, needed for the formation of amyloid fibrils was reduced to of FAP livers after domino liver transplantation. How- Iatrogenic