

Name of journal: *World Journal of Hepatology*

ESPS Manuscript NO: 31871

Manuscript Type: Original Article

Observational Study

Risk factors for hepatocellular carcinoma in cirrhosis due to nonalcoholic fatty liver disease: ¹A multicenter, case-control study

¹Kathleen E Corey, Samer Gawrieh, Andrew S deLemos, Hui Zheng, Andrew E Scanga, Jennifer W Haglund, Jorge Sanchez, Christopher J Danford, Megan Comerford, Krista Bossi, Samina Munir, Naga Chalasani, Julia Wattacheril

Abstract

AIM

Corey and Colleagues (1)

Match Overview

1	Internet 29 words crawled on 27-Aug-2016 onlinelibrary.wiley.com	1%
2	Internet 16 words crawled on 28-Sep-2016 hec2016.online-registry.net	1%
3	Crossref 15 words Satapathy, Sanjaya Kumar, Satheesh Nair, and Jason M. ... natta. "Nonalcoholic fatty liver disease following liver transp	1%

Risk factors for hepatocellular carcinoma in cirrhosis due to nonalcoholic fatty live

全部 图片 视频 新闻 地图 图书

找到约 124,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[PDF] [Nonalcoholic fatty liver disease and hepatocellular carcinoma: A ...](#)
[onlinelibrary.wiley.com/doi/10.1002/hep.23594/pdf](#) ▼

... or cryptogenic **cirrhosis**, which is associated with the typical **risk factors** for NASH. ... of **disease** ranging from iso- lated **hepatic steatosis** to **nonalcoholic** steatohepatitis and NAFLD.67-69 More recently, a **case-control study** in the United States **multicenter** comparative study with chronic hepatitis B. Gastroenterol-

[The Natural Course of Non-Alcoholic Fatty Liver Disease - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4881593/>

20 May 2016 ... **Risk factors** for fibrosis progression in **non-alcoholic fatty liver** ... The degree of **hepatic steatosis** does not appear to predict **disease** progression in NASH. Finally, Angulo et al. conducted an international **multicentre** cohort study to diabetes in patients with cryptogenic **cirrhosis**: A **case-control study**.

[Nonalcoholic fatty liver disease, metabolic risk factors, and ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4394070/>

14 Apr 2015 ... **Non-alcoholic liver disease** (NAFLD) defines **liver** abnormalities ranging from simple ... The current definition of NAFLD does not require secondary **hepatic fat** ... Major **risk factors** for **hepatocellular carcinoma** (HCC), the most ... A systematic review and **meta-analysis** of 26 **studies** (13 **case-control** and 13 ...

[Distinctive features of hepatocellular carcinoma in non-alcoholic ...](#)

[thelancet.com/journals/langas/article/PIIS2468-1253\(16\)30018-8/fulltext](#)

4 Oct 2016 ... **Non-alcoholic fatty liver disease** (NAFLD) and non-alcoholic incidence, prevalence, and **hepatocellular carcinoma risk factors** in C-related **cirrhosis**, suggesting that **hepatocellular carcinoma** risk in With this unknown contribution as the main rationale, a **multicentre case-control study** conducted in ...

[PDF] [Viral hepatitis, non-alcoholic fatty liver disease and alcohol as risk ...](#)

Risk factors for hepatocellular carcinoma in cirrhosis due to nonalcoholic fatty live

全部 图片 视频 新闻 地图 图书

找到约 124,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[PDF] [Nonalcoholic fatty liver disease and hepatocellular carcinoma: A ...](#)
[onlinelibrary.wiley.com/doi/10.1002/hep.23594/pdf](#) ▼

... or cryptogenic **cirrhosis**, which is associated with the typical **risk factors** for NASH. ... of **disease** ranging from iso- lated **hepatic steatosis** to **nonalcoholic** steatohepatitis and NAFLD.67-69 More recently, a **case-control study** in the United States **multicenter** comparative study with chronic hepatitis B. Gastroenterol-

[The Natural Course of Non-Alcoholic Fatty Liver Disease - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4881593/>

20 May 2016 ... **Risk factors** for fibrosis progression in **non-alcoholic fatty liver** ... The degree of **hepatic steatosis** does not appear to predict **disease** progression in NASH. Finally, Angulo et al. conducted an international **multicentre** cohort study to diabetes in patients with cryptogenic **cirrhosis**: A **case-control study**.

[Nonalcoholic fatty liver disease, metabolic risk factors, and ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4394070/>

14 Apr 2015 ... **Non-alcoholic liver disease** (NAFLD) defines **liver** abnormalities ranging from simple ... The current definition of NAFLD does not require secondary **hepatic fat** ... Major **risk factors** for **hepatocellular carcinoma** (HCC), the most ... A systematic review and **meta-analysis** of 26 **studies** (13 **case-control** and 13 ...

[Distinctive features of hepatocellular carcinoma in non-alcoholic ...](#)

[thelancet.com/journals/langas/article/PIIS2468-1253\(16\)30018-8/fulltext](#)

4 Oct 2016 ... **Non-alcoholic fatty liver disease** (NAFLD) and non-alcoholic incidence, prevalence, and **hepatocellular carcinoma risk factors** in C-related **cirrhosis**, suggesting that **hepatocellular carcinoma** risk in With this unknown contribution as the main rationale, a **multicentre case-control study** conducted in ...

[PDF] [Viral hepatitis, non-alcoholic fatty liver disease and alcohol as risk ...](#)

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 142,000 条结果 (用时 1.27 秒)

Cirrhosis and Chronic Viral Hepatitis as Risk Factors for ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: CV Constantin - 2010 - 被引用次数: 8 - 相关文章

Keywords: **Hepatocellular carcinoma**, liver **cirrhosis**, viral hepatitis, HbsAg ... Recent **studies** suggest it is more likely this is a result of gender-**related** inflammatory response (8-10). ... among high prevalence of **non-alcoholic fatty liver disease** (NAFLD) **cases** (18-21). ... Distribution of **risk factors** in the **study** and **control** groups.

Advances in hepatocellular carcinoma: Nonalcoholic steatohepatitis ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: FZ Khan - 2015 - 被引用次数: 18 - 相关文章

跳到**RISK FACTORS FOR NASH-RELATED HCC** - NASH-**related** HCC patients undergoing **liver** ... Yasui et al[16] performed a **multicenter** retrospective study in Japan to understand the pathological ... between non-**cirrhotic fatty liver disease** and HCC. ... a matched, nested **case control study** of 1303 diabetics ...

Nonalcoholic fatty liver disease, metabolic risk factors, and ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: LAM Streba - 2015 - 被引用次数: 26 - 相关文章

2015年4月14日 - **Non-alcoholic liver disease** (NAFLD) defines **liver** abnormalities ranging from simple ... The current definition of NAFLD does not require secondary **hepatic fat** ... Major **risk factors** for **hepatocellular carcinoma** (HCC), the most ... A systematic review and meta-analysis of 26 **studies** (13 **case-control** and 13 ...

Distinctive features of hepatocellular carcinoma in non-alcoholic fatty ...

[thelancet.com/journals/langas/article/PIIS2468-1253\(16\)30018-8/fulltext](http://thelancet.com/journals/langas/article/PIIS2468-1253(16)30018-8/fulltext) - 翻译此页

2016年10月4日 - **Non-alcoholic fatty liver disease** (NAFLD) and non-alcoholic ... including **hepatic**