

3

Name of journal: *World Journal of Gastrointestinal Pharmacology and Therapeutics*

ESPS Manuscript NO: 28184

Manuscript Type: Original Article

Randomized Clinical Trial

Efficacy of small-volume simethicone given at least 30 min before gastroscopy

1

Mingjun Song, Andrew Boon Eu Kwek, Ngai Moh Law, Jeannie Peng Lan Ong, Jessica Yi Tan, Prem Harichander Thurairajah, Danhne Shih Wen Ang

Match Overview



1

Crossref 15 words

"ASGE Program", Gastrointestinal Endoscopy, 2016.

1%

2

Internet 14 words

crawled on 22-Jun-2016

www.wjgnet.com

<1%

3

Internet 13 words

crawled on 02-Jun-2016

pancreas.imedpub.com

<1%



Efficacy of small-volume simethicone given at least 30 minutes before gastrc

全部

图片

视频

新闻

购物

地图

图书

找到约 1,420 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Small Volume Simethicone Before Gastroscopy: Any Benefit? - Full ...](https://clinicaltrials.gov/ct2/show/NCT02555228)

<https://clinicaltrials.gov/ct2/show/NCT02555228>

13 Sep 2015 ... Endpoint Classification: **Efficacy** Study Intervention Model: Parallel ... Liquid **simethicone** (1ml **volume**) in 5mls of water. Drug: **Simethicone** ... Other: Water. 5mls of water is **given at least 30 minutes before the gastroscopy**.

[Small Volume Simethicone Before Gastroscopy: Any Benefit?](http://www.bioportfolio.com/.../Small-Volume-Simethicone-Before-Gastroscopy-Any-Benefit.html)

www.bioportfolio.com/.../Small-Volume-Simethicone-Before-Gastroscopy-Any-Benefit.html

22 Sep 2015 the **efficacy** of liquid **simethicone** (100mg) in 5 mls water, versus placebo (5mls of water), as premedication (**given at least 30 minutes**) **before** ...

[The effectiveness of simethicone in improving visibility during ...](http://www.giejournal.org/article/S0016-5107(95)70041-2/abstract)

[www.giejournal.org/article/S0016-5107\(95\)70041-2/abstract](http://www.giejournal.org/article/S0016-5107(95)70041-2/abstract)

November 1995Volume 42, Issue 5, Pages 413–415 ... Benefits of **Small Volume Simethicone** Premedication If **Given at Least 30 Minutes Before Gastroscopy**; ...

[Efficacy of simethicone and N-acetylcysteine as premedication in ...](http://www.ncbi.nlm.nih.gov/pubmed/24325147)

www.ncbi.nlm.nih.gov/pubmed/24325147

Efficacy of **simethicone** and N-acetylcysteine as premedication in improving visibility during

Efficacy of small-volume simethicone given at least 30 min before gastroscopy

全部

图片

视频

新闻

购物

地图

图书

找到约 1,520 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Small Volume Simethicone Before Gastroscopy: Any Benefit? - Full ...](https://clinicaltrials.gov/ct2/show/NCT02555228)

<https://clinicaltrials.gov/ct2/show/NCT02555228>

13 Sep 2015 ... Endpoint Classification: **Efficacy** Study Intervention Model: ... Other: Water. 5mls of water is **given at least 30 minutes** before the **gastroscopy**.

[Small Volume Simethicone Before Gastroscopy: Any Benefit?](http://www.bioportfolio.com/.../Small-Volume-Simethicone-Before-Gastroscopy-Any-Benefit.html)

www.bioportfolio.com/.../Small-Volume-Simethicone-Before-Gastroscopy-Any-Benefit.html

22 Sep 2015 ... the **efficacy** of liquid **simethicone** (100mg) in 5 mls water, versus placebo (5mls of water), as premedication (**given at least 30 minutes**) **before** ...

[\[PDF\] Efficacy of simethicone and Nacetylcysteine as premedication in ...](https://onlinelibrary.wiley.com/doi/10.1111/jgh.12487/pdf)

onlinelibrary.wiley.com/doi/10.1111/jgh.12487/pdf

19 Nov 2013 ... **volume**, a 5-mL **simethicone** suspension administered more than **30 min prior** to upper **endoscopy** ... causing **small** bubbles to coalesce and collapse to release trapped ... visibility was between 10 and **30 min before EGD**. 12 Three liquid The amount of water to be **given** with **simethicone** is a matter of.

[Efficacy of premedication with activated Dimethicone or N ... - NCBI](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3208366/)

www.ncbi.nlm.nih.gov/pmc/articles/PMC3208366/

7 Oct 2011 ... AIM: To assess the **efficacy** of N-acetylcysteine (NAC) and activated ... with

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 1,610 条结果 (用时 0.53 秒)

Small Volume Simethicone Before Gastroscopy: Any Benefit? - Full ...

<https://clinicaltrials.gov/ct2/show/NCT02555228> ▾ [翻译此页](#)

2015年9月13日 - Endpoint Classification: **Efficacy** Study Intervention Model: ... Other: Water. 5mls of water is **given at least 30 minutes before the gastroscopy**.

Small Volume Simethicone Before Gastroscopy: Any Benefit?

www.bioportfolio.com ▾ [Topics](#) ▾ [Patient Monitoring](#) ▾ [Research](#) ▾ [翻译此页](#)

2015年9月22日 - ... which compares the **efficacy** of liquid **simethicone** (100mg) in 5 mls wa... as premedication (**given at least 30 minutes**) before gastroscopy ...

Sa1039 The Benefits of Small Volume Simethicone Premedication If ...

https://www.researchgate.net/.../301938493_Sa1039_The_Benefits_of_Small... - [翻译此页](#)

2016年5月8日 - Sa1039 The Benefits of **Small Volume Simethicone** Premedication If **Given at Least 30 Minutes Before Gastroscopy**; a Randomised, ...

Simethicone for the Preparation before Esophagogastroduodenoscopy

www.ncbi.nlm.nih.gov ▾ [NCBI](#) ▾ [Literature](#) ▾ [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: M Ahsan - 2011 - 被引用次数: 6 - 相关文章

2011年8月4日 - Administration of **simethicone** prior to EGD reduces the amount of ... Fifteen to **30 minutes before EGD**, in the presence of the researcher, ... to the **small** amount of duodenal air bubbles during the procedure, ... We suggest comparison of **effectiveness** of different doses, **timing** of administration, and **volume** of ...

Efficacy of simethicone and Nacetylcysteine as premedication in ...

onlinelibrary.wiley.com/doi/10.1111/jgh.12487/pdf ▾ [翻译此页](#)

2013年11月19日 - **volume**, a 5-mL **simethicone** suspension administered more than **30 min prior to upper endoscopy** ... causing **small** bubbles to coalescence and collapse to release trapped ... visibility was between 10 and **30 min before EGD**. 12 Three liquid The amount of water to be **given** with