

74748_Auto_Edited.docx

Name of Journal: *World Journal of Radiology*

Manuscript NO: 74748

Manuscript Type: EVIDENCE REVIEW

Expanding utility of cardiac computed tomography in infective endocarditis: A contemporary review

Cardiac CT in Infective Endocarditis

Diarmaid Hughes, Richard Linchangco, Reza Reyaldeem, Bo Xu

Abstract

There is increasing evidence on the utility of cardiac CT (CCT) in infective endocarditis (IE) to investigate the valvular pathology, the extra-cardiac manifestations of IE and pre-operative planning. CCT can assist in the diagnosis of perivalvular complications, such as pseudoaneurysms and abscesses, and can help identify embolic events to the lungs or systemic vasculature. CCT has also been shown to be beneficial in the pre-operative planning of patients by delineating the coronary artery anatomy and the major cardiovascular structures in relation to the sternum. Finally, hybrid nuclear/computed tomography techniques have been shown to increase the diagnostic accuracy in prosthetic valve endocarditis. This manuscript aims to provide a contemporary update of the existing evidence base for the use of CCT in IE.

Key Words: Infective Endocarditis; Cardiac CT; Multimodality Cardiac Imaging

Hughes D, Linchangco R, Reyaldean R, Xu B. Expanding utility of cardiac computed tomography in infective endocarditis: A contemporary review. *World J Radiol* 2022; In press

Core Tip: Cardiac CT (CCT) has an expanding role in the management of infective endocarditis (IE). It has been shown to be superior to echocardiography for diagnosing perivalvular complications such as pseudoaneurysms and abscesses. CCT can also diagnose extra-cardiac manifestations of IE such as septic emboli to the lungs. It can assist in pre-operative planning by delineating the coronary anatomy and assessing vascular structures. Herein, we review the role of CCT in IE including the evidence base comparing CCT to echocardiography in diagnosing the valvular complications of IE and the use of CT in IE beyond valvular assessment.

1 INTRODUCTION

Infective endocarditis (IE) is an infection of the endocardium, heart valves or intra-cardiac devices. It remains a challenging disease to diagnose and manage with high rates of morbidity and mortality^[1,2]. Echocardiography remains the main imaging modality used in IE; more recently, however, there is an increasing evidence base for a multimodality imaging approach for IE. Complementary imaging modalities including cardiac computed tomography (CCT) now play increasingly important roles in diagnosis, risk stratification and management of IE. CCT has certain advantages compared to echocardiography in being able to investigate for perivalvular extension, extra-cardiac complications of IE, including metastatic spread and planning for surgery including assessing for coronary artery disease. Advancements in CT technologies, including the use of dedicated cardiac gated four-dimensional CCT, have expanded the applications of CT in IE, demonstrating good sensitivity and specificity for diagnosing the complications of IE. This article aims to review the available evidence for the use of CCT in IE.

CLINICAL CONSIDERATIONS

The incidence of IE in the United States is estimated to be approximately 15 per 100000 persons annually^[3,4], with *Staphylococcus aureus* (SA) being the most common pathogen followed by Viridans group Streptococci^[5]. A number of risk factors have been identified for acquiring IE, including the presence of a prosthetic valve, a previous episode of IE, patients with untreated cyanotic congenital heart disease, injection drug use, poor dentition and pre-existing valvular heart disease^[6]. The clinical presentation of IE can vary significantly from an acute life threatening illness to a more indolent chronic disease^[7]. The most common presenting symptoms are: fever, cardiac murmur, heart failure or complications from septic emboli^[8].

The Duke criteria were developed in 1994 to assist in the risk stratification of patients with suspected IE into definite, possible and rejected cases of IE^[9]. These criteria have been since validated by a number of retrospective analyses^[10,11], and underwent further

modification in 2000 to reflect changing clinical practice and the emergence of SA as the most common pathogen encountered^[12]. Despite the updated clinical criteria for diagnosis of IE, there often remains a delay in diagnosis for many patients, commonly due to a lack of microbiological criteria from inappropriate antibiotic use, with worse outcomes seen in these patients^[13,14]. Advances in CT technologies including improvements in both temporal and spatial resolution have enabled greater use of CCT for the diagnosis of IE. The European Society of Cardiology (ESC) guidelines for the management of infective endocarditis reflect these advances in imaging techniques and include paravalvular lesions detected by CCT to be a major imaging criterion^[15]. The 2020 American College of Cardiology (ACC) / American Heart Association (AHA) Guideline for the management of patients with valvular heart disease also recommend the use of CCT as an adjunctive imaging modality for IE^[16].

Echocardiography

Echocardiography, including transthoracic echocardiography (TTE) and transesophageal echocardiography (TEE) where appropriate, remain the first line imaging modality for the diagnosis and monitoring of IE^[17,18]. There are three main echocardiographic findings that are considered major criteria for the diagnosis of IE: vegetations, abscesses/pseudoaneurysms and new dehiscence of a prosthetic valve^[15]. In native valve endocarditis (NVE), the sensitivities for the diagnosis of a vegetation are approximately 70% for TTE and 96% for TEE, respectively^[17]. For prosthetic valve endocarditis (PVE), the sensitivities for diagnosing a vegetation are approximately 50% for TTE and 92% for TEE, respectively^[17]. There are many challenges in the diagnosis of IE with echocardiography, including small vegetations or embolization of the vegetation prior to imaging, difficulty visualizing the lesion in the setting of pre-existing valvular disease or with prosthetic valves. Small abscesses may also be challenging to diagnose, especially by TTE^[19]. There are also many mimics of IE that could result in a false positive diagnosis, such as Lambl's excrescence, fibroelastomas, thrombus, degenerative lesions, prosthetic material/sutures or marantic lesions^[20]. CCT

can therefore be helpful to assist in the diagnosis of IE, when there are equivocal findings by echocardiography or in challenging cases involving prosthetic valves^[21].

Dedicated Cardiac CT Protocol for Endocarditis Evaluation

For evaluation of cardiac valves, multiphase imaging using a retrospectively ECG-gated acquisition is required to obtain an isotropic data set. Images are acquired in spiral mode utilizing a low pitch of 0.16 to 0.5 during 5-10 R-R intervals with a section thickness of 0.60 mm. Thin collimation is used for optimal visualization of the valve leaflets, typically 64 x 0.6 mm. The tube voltage used is adapted to the patient's weight, and can vary between 100 to 120 kV. Gantry rotation time of 0.28 to 0.35 s is used^[22-27]. The scan is typically performed in a limited field of view from the level of the carina to the cardiac apex.

Timing of the iodinated contrast bolus is important to optimize visualization of the involved cardiac valves. A monophasic contrast injection is most commonly used, with timing of the contrast bolus chosen for optimal visualization of the expected involved valve and cardiac chambers. Alternatively, biphasic contrast injection may be performed, which allows evaluation of all cardiac chambers and valves^[28]. Premedication with beta blockers may be used to regulate heart rate to less than 65-70 bpm if not contraindicated. This improves image quality by reducing artefacts related to cardiac motion and valvular motion.

The isotropic data set acquired from the retrospectively gated acquisition allows for reconstruction in any desired plane. In addition to static images, imaging at multiple points during the cardiac cycle also allows for creation of 4D cine images, allowing for evaluation of valve leaflet motion and planimetry.

Because images are acquired throughout the entire cardiac cycle, this results in a significantly higher radiation dose penalty compared to prospectively gated CT as used

typically in CT angiography of the coronary arteries. Radiation dose may be lowered utilizing methods such as iterative reconstruction and ECG-triggered radiation dose modulation. However, ECG-triggered dose modulation may result in suboptimal evaluation during the phase of reduced tube current, typically the systolic phase^[29]. There are specific protocols used for visualizing the various cardiac and extra manifestations of IE and for pre-operative planning. Herein, we group all of these into an umbrella term of CCT, referring to ECG-gated CT of the chest with contrast. There are some situations, such as during investigation for septic emboli to the visceral organs, when abdominal imaging may also be needed.

UTILITY OF CARDIAC CT IN INFECTIVE ENDOCARDITIS

CCT has the ability to assess for valvular lesions, perivalvular extension, metastatic spread/embolization, as well as aortic anatomy. CCT may also be used in appropriate cases for the assessment of coronary artery disease^[30]. This is particularly relevant in patients with aortic valve IE and vegetations, whereby invasive coronary angiography may be relatively contraindicated, due to the potential risk of causing the vegetations to embolize during the procedure. CCT can detect valvular lesions, such as vegetations, prosthetic valve dehiscence in addition to perivalvular lesions such as abscesses, fistulae and pseudoaneurysms^[26,31]. Table 1 compares the various definitions for IE detected on CCT *vs* TEE.

Perivalvular complications

Pseudoaneurysms and abscesses

Perivalvular extension of IE, which includes pseudoaneurysms, abscesses and fistulae are associated with a higher rate of operative management and mortality^[32-34]. A pseudoaneurysm is a perivalvular cavity that is in communication with the cardiovascular lumen which results from an abscess rupturing into a cavity^[17]. On echocardiography, this appears as a pulsatile echo-free space with detectable Doppler

color flow, while on CCT, it appears as an abnormal cavity close to the valve with direct communication with the heart chambers or major blood vessel^[30]. An abscess is a closed cavity with necrosis and purulent material not in communication with a cardiovascular cavity^[17]. On echocardiography, this appears as a thickened perivalvular area with a homogenous echo-dense or echo-lucent appearance. On CCT, abscesses appear as perivalvular collections of fluid encased in a thick layer of inflammatory tissue enhanced by the injection of contrast medium. See Figures 1-4 for examples of comparisons of TEE and CCT images in patients with perivalvular complications of IE.

A 2009 study by Feuchter *et al*^[24] to investigate the value of CCT for the assessment of valvular abnormalities included 37 consecutive patients (29 of whom went on to have surgery) with clinically suspected IE who underwent both CCT and TEE. CCT identified all pseudoaneurysms and abscesses in this study with sensitivity and specificity of 100%, which was superior to TEE (sensitivity of 89% and a specificity of 100%)^[24]. CCT was also shown to be superior to TEE for perivalvular extension of the IE, identifying myocardial and pericardial extension more often than TEE^[24]. In a 2009 prospective study, 19 patients with aortic valve endocarditis requiring surgical intervention underwent CCT pre-operatively^[25]. The majority of patients (approx. 90%) had native valve IE (NVE). This study showed that CCT had sensitivity and specificity for diagnosing pseudoaneurysms of 100% and 92%, respectively, and CCT correctly identified all cases where there was extension of IE into the intervalvular fibrosa^[25]. This paper did not report the TEE findings for their participants ^[25].

A paper by Fagman *et al*^[23] reported in 2012 on 27 consecutive patients who had TEE findings of aortic valve PVE and investigated the strength of agreement between the TEE and CT results. They found a strength of agreement compared to TEE was 0.68 for abscesses and 0.75 for dehiscence^[23]. However, using surgery as the reference standard (16 patients went on to have surgery), CCT had sensitivity of 91% to detect pseudoaneurysms /abscesses compared to 82% for TEE^[23]. A 2013 study investigated

the additional value of CCT beyond the usual evaluation with TEE in PVE in 28 patients, with a final diagnosis being either determined clinically or at the time of surgery as reported by Habets *et al*^[35]. They reported that usual evaluation had sensitivity of 68% for detecting periannular complications (mycotic aneurysms and abscesses), which was increased to 100% with the use of CCT^[35]. Koo *et al*^[27] also compared CCT *vs* TEE using intra-operative findings as the reference standard in 2018. They enrolled 49 patients, 12 of whom had PVE^[27]. The overall detection of IE by CCT was 94%, compared to 96% by TEE^[27]. CCT performed better than TEE at detecting abscess/pseudoaneurysms, with sensitivities of 60% for CCT and 40% for TEE, respectively^[27]. A retrospective study from 2018 by Sims *et al*^[36] investigated the performance of CCT in the pre-operative evaluation of IE. In total, they had 251 patients undergoing TEE with 34 of these patients also having a CCT^[36]. The sensitivity of CCT for detecting abscesses/pseudoaneurysms was 91%, which was superior to TEE at 78%^[36]. CCT was reported to have a lower sensitivity for detecting fistulae at 50% *vs* 79% for TEE, and dehiscence at 57% *vs* 70% for TEE^[36].

Two studies from 2018 (Ouchi *et al* and Koneru *et al*^[31,37]) retrospectively investigated the utility of CCT in IE with intra-operative findings as the reference standard. CCT performed better than TEE in detecting abscess/pseudoaneurysm in prosthetic valves with sensitivity of 81% (versus 64% for TEE) in the study by Koneru *et al*^[31]. CCT had sensitivity of 100% sensitivity for detecting perivalvular complications, such as pseudoaneurysms in the paper by Ouchi *et al*^[37]. A 2019 study by Hryniewiecki *et al*^[26] investigated 53 consecutive patients who had perivalvular complications from IE, who also underwent CCT and TEE pre-operatively. They showed the sensitivity and specificity for detecting abscesses/pseudoaneurysms for CCT were 81% and 90%, respectively, compared to 63% and 90%, respectively for TEE^[26]. A 2020 study of 68 patients reported by Sifaoui *et al* with definite left-sided IE who underwent CCT and TEE reported the comparison of CCT and TEE to detect perivalvular complications^[38].

They showed again that CT had a higher sensitivity for detecting pseudoaneurysms at 100%, compared to TEE at 67%^[38].

Overall, the current evidence base suggests that the diagnostic performance of CCT is likely superior to that of TEE for the detection of pseudoaneurysms and abscesses in appropriately selected cases. A recent meta-analysis^[39] reported pooled sensitivity and specificity for CCT for the detection of peri-annular complications of 88% and 93%, respectively, compared to TEE at 70% and 96%, respectively.

The identification of perivalvular complications is important for prognostic and management considerations. These sequelae of invasive IE, which are more common with aortic valve endocarditis and PVE, have been associated with increased rates of surgical management, and may confer an increased risk of mortality^[32,33]. Therefore, a multimodality imaging strategy for IE that includes CCT would have the ability to identify more of these complications, compared to using TEE alone, and therefore impact on decision making for patients.

Vegetations

A vegetation is a mass-like lesion of infected material composed of fibrin, platelets and microorganisms attached to an endocardial structure or on an implanted cardiac device (CIED)^[40,41]. On echocardiography, this appears as an oscillating or non-oscillating intracardiac echodensity, which can be attached to a valve, other endocardial surface or cardiac device^[15]. A vegetation tends to move with the cardiac cycle, and is more frequently found on the atrial side of the atrioventricular valves and the ventricular side of the semi-lunar valves^[15]. On CT, vegetations appear as hypodense homogeneous irregular masses, which can be attached to a valve or other cardiac structures^[30].

In the 2009 paper by Feuchter *et al*^[24] using surgical/pathological diagnosis as the reference standard, CCT had sensitivity of 96% and specificity of 97% for the diagnosis of vegetations. 5 vegetations were missed by CCT (11%) either due to artefact or small size ($\leq 4\text{mm}$)^[24]. The performance of TEE was similar to CCT with sensitivity of 96% and specificity of 100%^[24]. CCT was found to be inferior to TEE for detecting leaflet perforations^[24]. The study by Gahide *et al*^[25] on aortic valve IE showed CCT had a sensitivity of 71% and a specificity of 100% for detecting vegetations, though the sensitivity was increased to 100% for large vegetations ($>10\text{mm}$). The 2012 paper by Fagman *et al*^[23] found that CCT detected vegetations in 7 out of 13 cases (54%), with a lower detection rate being potentially explained by artefact from the prosthetic valves obscuring the CCT images. The 2013 study by Habets *et al*.^[35] found additional benefit with CCT in addition to usual work-up with TEE in PVE, with a final diagnosis being either determined clinically or at the time of surgery. They reported that usual work-up had sensitivity of 63% for detecting vegetations, which was increased to 100% with the use of CCT^[35]. The 2018 study by Koo *et al*^[27] reported sensitivity for CCT to detect vegetations of 91%, compared to 100% by TEE. Missed vegetations were smaller, and the authors also listed motion artefact and beam hardening from mechanical valves as reasons for the failure of CCT to detect the vegetations^[27]. Sims *et al*^[36] reported the sensitivity for detecting vegetations to be 70% for CCT (34 patients) and 96% for TEE (251 patients). The study by Oucho *et al*^[37] reported sensitivity of 92% for detecting vegetations for CCT, correctly identifying 12 of 13 cases who had vegetations confirmed at the time of surgery. The retrospective review on 122 patients by Koneru *et al*^[31] showed TEE to have a statistically significantly higher sensitivity for detecting vegetations compared to CCT at 85% *vs* 16%, though CCT did have a higher specificity at 96% compared to 69% for TEE. The lower sensitivity in detecting vegetations by CCT in this study may be related to only reviewing single-phase images, and the fact that the slice thickness used was 3 mm which was thicker than the other studies referenced above^[31]. In the 2019 paper by Hryniewiecki *et al*^[26] the sensitivity and specificity for detecting vegetations by CT were 89% and 71%, respectively, compared to TEE at 97%

and 42%, respectively. The 2020 study by Sifaoui *et al*^[38] showed that TEE had a higher area under the curve (AUC) than CCT for detecting vegetations, with AUC for TEE of 0.86 *vs* AUC for CT of 0.69.

Overall, the current evidence base suggests that TEE is overall superior to CCT for the detection of vegetations, particular small vegetations, with pooled sensitivity for TEE from a recent meta-analysis of 94%^[42]. CCT demonstrated a lower pooled sensitivity, at 64% for the detection of vegetations^[42]. There was a wide range of results reported likely related to small sample sizes, differing patient populations and different protocols used for imaging. While CCT should not replace echocardiography as the first line imaging tool in the majority of patients primarily to detect vegetations, in a small subset of patients who could not undergo clinical indicated TEE (e.g. esophageal pathology), CCT may add diagnostic value. CCT has also been shown to improve the diagnostic accuracy overall, when used in combination with TEE^[35]. For example figure 5 shows a hypoattenuating lesion on a mechanical aortic valve that was more clearly defined on CCT. Figures 6 and 7 shows an examples of an aortic graft and aortic stent infections that can be difficult to image with TEE.

INCREMENTAL VALUE OF CCT IN INFECTIVE ENDOCARDITIS

Pre-operative assessment

In addition to the advantages related to the management of IE as outlined above, CCT can also assist in the pre-operative planning of IE surgery. In patients with prior cardiothoracic surgery, CCT can delineate the relationship of cardiovascular structures to the sternum and the location of the coronary artery bypass grafts (CABG). See Figure 8 for CCT images of a patient with a prior CABG. For all patients, CCT can identify calcification of the ascending aorta ('porcelain aorta'), which may preclude surgery as well as give precise anatomic location and extent of the degree of calcification of the subclavian, axillary and femoral arteries. The advantage of having a pre-operative CCT was described by Merlo *et al*^[43] with reported lower rates of stroke and mortality in

patients undergoing pre-operative CCT followed by primary cardiac surgery, *vs* those without pre-operative CCT imaging. Figure 9 shows a CCT in a patient with vascular calcification.

Pre-operative assessment

In addition to the advantages related to the management of IE as outlined above, CCT can also assist in the pre-operative planning of IE surgery. In patients with prior cardiothoracic surgery, CCT can delineate the relationship of cardiovascular structures to the sternum and the location of the coronary artery bypass grafts (CABG). See Figure 8 for CCT images of a patient with a prior CABG. For all patients, CCT can identify calcification of the ascending aorta ('porcelain aorta'), which may preclude surgery as well as give precise anatomic location and extent of the degree of calcification of the subclavian, axillary and femoral arteries. The advantage of having a pre-operative CCT was described by Merlo *et al*^[43] with reported lower rates of stroke and mortality in patients undergoing pre-operative CCT followed by primary cardiac surgery, *vs* those without pre-operative CCT imaging. Figure 9 shows a CCT in a patient with vascular calcification.

CONCLUSION

With improvements in the temporal and spatial resolution of CCT technology, including the use of dedicated 4D CCT, there has been an expanding role of CCT imaging in IE. CCT has been shown to be superior to TEE for the identification of pseudoaneurysms and abscesses in appropriately selected cases, while the combination of both modalities results in the greatest sensitivity for detection. TEE is superior to CCT for small vegetations; however this advantage is less marked for larger vegetations. In addition, CCT has a number of adjunctive uses in IE beyond evaluation of the valvular pathology. CCT can aid in the diagnosis of embolic events, both pulmonary complications in RSIE, and systemic embolism. It can also be used to diagnose significant CAD in low to intermediate risk patients preoperatively, or when

there is a contraindication to ICA, such as when there is a large aortic valve vegetation. CCT can also be helpful for pre-operative planning to assess the relationship of the cardiovascular structures in relation to the sternum, which is particularly helpful in redo sternotomy cases. The addition of hybrid techniques such as PET or SPECT/CT, has been shown to improve the diagnostic accuracy in challenging cases of PVE. Greater awareness of the strengths, weaknesses and appropriate applications of CCT in IE will assist in its optimal use for improved diagnosis and management of this challenging condition.

3%

SIMILARITY INDEX

PRIMARY SOURCES

1	link.springer.com Internet	54 words — 1%
2	eagleseye.acc.org Internet	22 words — 1%
3	G. Habib. "Guidelines on the prevention, diagnosis, and treatment of infective endocarditis (new version 2009): The Task Force on the Prevention, Diagnosis, and Treatment of Infective Endocarditis of the European Society of Cardiology (ESC)", European Heart Journal, 08/27/2009 Crossref	16 words — < 1%
4	www.science.gov Internet	13 words — < 1%