

7 Name of journal: World Journal of Radiology

ESPS Manuscript NO: 17275

Columns Case Report

Isolated renal hydatid presenting as a complex renal lesion followed by spontaneous hydatiduria

Anil Bhaya, Archana P Shinde

Abstract

Echinococcosis is a zoonotic disease. **4** Liver is the most common site of involvement. Renal involvement is seen in 2 to 3 % of patients. CT findings in renal hydatid typically include: a cyst with thick or calcified wall, unilocular cyst with detached membrane, a multiloculated cyst with mixed internal density and daughter cysts with lower density than maternal matrix. Rarely type IV hydatid cysts may mimic hypovascular renal cell carcinoma. We report a case of previously asymptomatic middle aged female who presented with mild intermittent pain and a complex renal lesion on imaging which was considered to be a hypovascular renal carcinoma or urothelial neoplasm. However, by serendipity, the patient had spontaneous hydatiduria and later was definitively diagnosed and stented. Hydatid disease **11** should always be considered amongst the top differential diagnosis of an isolated "complex" renal lesion which remains indeterminate on imaging.

Key Words: - Hydatid, Renal, Hydatidosis, Echinococcosis, Hydatiduria, Kidney

Match Overview

1	Internet 71 words crawled on 26-Sep-2014 www.libreriauniverso.it	4%
2	Internet 64 words crawled on 16-Jan-2010 www.ajronline.org	4%
3	Internet 20 words crawled on 07-Jan-2004 indianpediatrics.net	1%
4	Internet 17 words crawled on 29-Apr-2014 fr.slideshare.net	1%
5	CrossCheck 12 words Turgut, A.T.. "Unusual imaging characteristics of complicated hydatid disease", European Journal of Radiology, 2007	1%
6	Internet 10 words crawled on 21-Oct-2010 bhj.org	1%
7	Internet 10 words crawled on 09-Apr-2015	1%



ISOLATED RENAL HYDATID PRESENTING AS COMPLEX RENAL LES



网页

图片

视频

新闻

更多 ▾

搜索工具

找到约 2,890 条结果 (用时 0.71 秒)

您是不是要找: ISOLATED RENAL HYDATID PRESENTING
AS COMPLEX RENAL LESION FOLLOWED BY
SPONTANEOUS *HYDATID UREA*

[PDF] The Rupture Of A Renal Hydatid Cyst Into The Renal Pelv...
tip.harran.edu.tr/tipdergisi/arsiv/200851_8.pdf ▾ 翻译此页

作者: MD Murat Savas - 相关文章

presented a woman with renal hydatid cysts with symptoms of ureteral colic and ... Key words: Hydatid disease, Renal hydatid cyst, renal colic, ureteral colic, hydatiduria ... and complex cystic lesion in kidney, diagnosis of ... spontaneous or traumatic rupture of renal hydatid ... simplest technique and can be followed safely by.

Isolated unilocular renal hydatid cyst: a rare diagnostic ...
www.researchgate.net/.../6662636_Isolated_unilocular_renal_h... - 翻译此页

Publication » Isolated unilocular renal hydatid cyst: a rare diagnostic difficulty with simple ... finding that is pathognomonic for hydatid disease except for hydatiduria. ... or computed tomography cannot always show a hydatidosis as a specific lesion. ... Article: Imaging features of renal hydatid cyst presenting with hydatiduria.

Translumbar puncture of a renal hydatid cyst. - ResearchGate

网页 图片 视频 新闻 更多 ▾ 搜索工具

找到约 3,210 条结果 (用时 0.97 秒)

您是不是要找: **Isolated renal hydatid presenting as a complex renal lesion followed by spontaneous *hydatid urea***

Primary Hydatid Cyst of the Kidney and Ureter with ...

www.hindawi.com/journals/crin/2012/596923/ ▾ 翻译此页

作者: V Seetharam - 2012 - 被引用次数: 2 - 相关文章

2012年10月17日 - Most common localization of hydatid cyst occurs in liver followed by lungs. Renal hydatid cyst constitutes about 2–4% of all locations. ... It is caused in humans by larval stage of Echinococcus granulosus complex, E. multilocularis, working in laboratories and presenting with mass lesions in relevant ...

[PDF] The Rupture Of A Renal Hydatid Cyst Into The Renal Pelv...

tip.harran.edu.tr/tipdergisi/arsiv/200851_8.pdf ▾ 翻译此页

作者: MD Murat Savas - 相关文章

presented a woman with renal hydatid cysts with symptoms of ureteral colic and ... Key words: Hydatid disease, Renal hydatid cyst, renal colic, ureteral colic, hydatiduria ... and complex cystic lesion in kidney, diagnosis of ... spontaneous or traumatic rupture of renal hydatid ... simplest technique and can be followed safely by.

Isolated renal hydatid cyst managed by laparoscopic ...

www.ncbi.nlm.nih.gov ▸ ... ▸ PubMed Central (PMC) ▾ 翻译此页

[网页](#)[图片](#)[视频](#)[新闻](#)[更多 ▾](#)[搜索工具](#)

找到约 3,880 条结果 (用时 0.72 秒)

Isolated renal hydatid cyst managed by laparoscopic ...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: KJ Shah - 2009 - 被引用次数: 11 - 相关文章

Isolated right renal hydatid cyst was removed in toto. ... We present a case of a 45-year-old female, with isolated renal echinococcosis, a very rare presentation, and its management by ... grape-sized, balloon-like structures in the urine (hydatiduria) for last 10 years ... There were no similar lesions in other abdominal viscera.

Imaging Features of Renal Hydatid Cyst Presenting with ...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: MU Amin - 2009 - 被引用次数: 5 - 相关文章

2009年3月1日 - Hydatiduria is an extremely rare manifestation of renal hydatid cyst. We report such ... Solid avascular component was also seen within the lesion. Upper pole of Ultrasonography shows a complex cyst in all of the patients. ... Isolated Unilocular Renal Hydatid Cyst: A Rare Diagnostic Difficulty with simple.

Primary Hydatid Cyst of the Kidney and Ureter with ...

www.hindawi.com/journals/crin/2012/596923/ ▾ [翻译此页](#)

作者: V Seetharam - 2012 - 被引用次数: 2 - 相关文章

2012年10月17日 - Most common localization of hydatid cyst occurs in liver followed by lungs. Renal hydatid cyst constitutes about 2-4% of all locations. ... It is caused in humans by larval stage of Echinococcus granulosus complex. E. multilocularis ... working in