

Name of Journal: *World Journal of Radiology*

ESPS Manuscript NO: 22504

Manuscript Type: FRONTIER

Brainstem Tegmental Lesions in neonates with hypoxic-ischemic encephalopathy: MR diagnosis and clinical outcome

Quattrocchi Carlo Cosimo, Fariello Giuseppe, Longo Daniela

Abstract

Lesions of the brainstem have been reported in the clinical scenarios of hypoxic-ischemic

Match Overview

| | | |
|---|--|----|
| 1 | CrossCheck 57 words SLATTERY, JUSTINE, ANGELA MORGAN, and JACINTA DOUGLAS. "Early sucking and swallowing problems as | 1% |
| 2 | CrossCheck 49 words Sugama, S.. "Brainstem lesions in children with perinata l brain injury", <i>Pediatric Neurology</i> , 200303 | 1% |
| 3 | CrossCheck 45 words Sugiura, Hiroshi, Masanori Kouwaki, Tohru Kato, Tsutomu Ogata, Rie Sakamoto, Atsushi Ieshima, and Kenji ... | 1% |
| 4 | CrossCheck 35 words Shioda, M.. "Lesions in the central tegmental tract in ai ... opsy cases of developmental brain disorders", <i>Brain an</i> | 1% |
| 5 | CrossCheck 27 words Martinez-Biarge, Miriam, Jesus Diez-Sebastian, Courtney J. Wusthoff, Stacey Lawrence, Annie Aloysius, Mary A. | 1% |
| 6 | CrossCheck 24 words Martinez-Biarge, M.. "Outcomes after central grey matte ... njury in term perinatal hypoxic-ischaemic encephalopath | 1% |



Brainstem Tegmental Lesions in neonates with hypoxic-ischemic encephalopa ▾



学术搜索

找到约 322 条结果 (用时0.06秒)

我的著作引用情况



文章

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

我的图书馆

[HTML] In Re: Barkovich AJ. MR and CT evaluation of profound neonatal and infantile asphyxia. ajnr.org 中的 [HTML]
AJNR Am J Neuroradiol 1992; 13: 959-972

ER Heinz - American Journal of Neuroradiology, 2000 - Am Soc Neuroradiology
... circulatory arrest and asphyxia (3). The basal ganglia, diencephalon, tegmentum, brain stem,
and spinal ... Lesions of the telencephalon and cerebral cortex were less prominent. ... Anoxic
encephalopathy with predominant involvement of basal ganglia, brain stem and spinal cord in ...

被引用次数: 242 相关文章 所有 7 个版本 引用 保存

Neonatal encephalopathy: an inadequate term for hypoxic-ischemic encephalopathy ucsfmedicalcenter.org 中的 [PDF]

JJ Volpe - Annals of neurology, 2012 - Wiley Online Library
... hippocampus), basal ganglia, thalamus, and brainstem (inferior colliculus and tegmental structures,
including ... pattern of injury, that is, basal ganglia/thalamus and brainstem tegmentum, with relative ...
the time of delivery have delineated a topography of lesions highly correlated ...

被引用次数: 41 相关文章 所有 7 个版本 引用 保存

Watershed infarcts in the fetal and neonatal brainstem. An aetiology of central hypoventilation, ejpn-journal.com 中的 [HTML]
dysphagia, Möbius syndrome and micrognathia

HB Sarnat - European Journal of Paediatric Neurology, 2004 - Elsevier
... 17 Much of it lies within the tegmentum. ... In severe cases of fetal and neonatal hypoxic/ischemic
encephalopathy, tegmental infarction may be present but, rather than restricted to an ... corresponding
mostly to watershed zones of the basilar circulation at the level of the brainstem. ...

被引用次数: 71 相关文章 所有 5 个版本 引用 保存

The EEG in coma

GB Young - Journal of Clinical Neurophysiology, 2000 - journals.lww.com
... Absolutely invariant EEGs usually indicate a severe encephalopathy and often a poor prognosis

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

创建快讯

学术搜索

找到约 330 条结果 (用时0.06秒)

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☒ 包含引用☐ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[HTML] In Re: Barkovich AJ. MR and CT evaluation of profound neonatal and infantile asphyxia. AJNR Am J Neuroradiol 1992; 13: 959-972

ER Heinz - American Journal of Neuroradiology, 2000 - Am Soc Neuroradiology

... circulatory arrest and asphyxia (3). The basal ganglia, diencephalon, tegmentum, brain stem, and spinal ... Lesions of the telencephalon and cerebral cortex were less prominent. ... Anoxic encephalopathy with predominant involvement of basal ganglia, brain stem and spinal cord in ...

被引用次数: 243 相关文章 所有 7 个版本 引用 保存

Neonatal encephalopathy: an inadequate term for hypoxic-ischemic encephalopathy

JJ Volpe - Annals of neurology, 2012 - Wiley Online Library

... hippocampus), basal ganglia, thalamus, and brainstem (inferior colliculus and tegmental structures, including ... pattern of injury, that is, basal ganglia/thalamus and brainstem tegmentum, with relative ... the time of delivery have delineated a topography of lesions highly correlated ...

被引用次数: 48 相关文章 所有 7 个版本 引用 保存

Watershed infarcts in the fetal and neonatal brainstem. An aetiology of central hypoventilation, dysphagia, Möbius syndrome and micrognathia

HB Sarnat - European Journal of Paediatric Neurology, 2004 - Elsevier

... 17 Much of it lies within the tegmentum. ... In severe cases of fetal and neonatal hypoxic/ischemic encephalopathy, tegmental infarction may be present but, rather than restricted to an ... corresponding mostly to watershed zones of the basilar circulation at the level of the brainstem. ...

被引用次数: 72 相关文章 所有 5 个版本 引用 保存

The EEG in coma

GB Young - Journal of Clinical Neurophysiology, 2000 - journals.lww.com

... Absolutely invariant EEGs usually indicate a severe encephalopathy and often a poor prognosis (Cant and ... into consideration the clinical picture, age, etiology, acuity, and the integrity of brainstem

ajnr.org 中的 [HTML]

ucsfmedicalcenter.org 中的 [PDF]

ejpn-journal.com 中的 [HTML]

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 6,410 条结果 (用时 0.68 秒)

Google 学术: Brainstem tegmental lesions in neonates with hypoxic-ischemic encephalopathy: Magnetic resonance diagnosis and clinical outcome

... inadequate term for hypoxic-ischemic encephalopathy - Volpe - 被引用次数: 49

... diagnosis of white matter injury in premature newborns ... - Miller - 被引用次数: 187

Neurologic aspects of neonatal hypoglycemia - Alkalay - 被引用次数: 26

Dorsal Brain Stem Syndrome: MR Imaging Location of Brain ...

www.ajnr.org/content/31/8/1438.full ▾ 翻译此页

作者: CC Quattrocchi - 2010 - 被引用次数: 6 - 相关文章

2010年4月15日 - MR imaging showed hypoxic-ischemic lesions in supra- and ... Oral motor dysfunction was associated with brain stem tegmental lesions in posthypoxic-ischemic infants. This examination may be crucial in defining the clinical diagnosis of ... Magnetic resonance imaging of brain injury in the high-risk term ...

Dorsal Brain Stem Syndrome: MR Imaging Location of Brain ...

www.researchgate.net/.../43159922_Dorsal_Brain_Stem_Syindr... - 翻译此页

MR imaging showed hypoxic-ischemic lesions in supra- and infratentorial areas. ... was associated with brain stem tegmental lesions in posthypoxic-ischemic infants. ... of brainstem lesions in fullterm infants with hypoxic ischemic encephalopathy (HIE) ... Article: Magnetic resonance imaging in neonates with total asphyxia.

Anatomic Changes and Imaging in Assessing Brain Injury in ...

[www.ncbi.nlm.nih.gov/...>Literature>PubMedCentral\(PMC\)](http://www.ncbi.nlm.nih.gov/...>Literature>PubMedCentral(PMC)) - 翻译此页

作者: BK Lawrence - 2008 - 被引用次数: 16 - 相关文章