

[国内版](#)[国际版](#)[All](#)[Images](#)[Videos](#)[翻译成中文](#)[关闭取词](#)

28,500 Results

Any time ▾

Mediastinal traction injury to the recurrent laryngeal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3616173>

There have also been case reports of injury following carotid artery stenting and subclavian line insertion. Anatomically, the left vagus nerve arrives in the mediastinum via the thoracic inlet before giving off its recurrent laryngeal branch at the arch of the aorta.

Cited by: 1

Author: Michael K. Ng, Andrew Barling, Sorway C...

Publish Year: 2012

The Recurrent Laryngeal Nerves and the Thoracic Surgeon ...

<https://www.ctsnet.org/article/recurrent-laryngeal-nerves-and-thoracic-surgeon> ▾

Sep 19, 2017 · Introduction Many thoracic surgeons are terrified to come anywhere near the recurrent laryngeal nerve (RLN), especially on the left side. The reason for this fear is the dreaded complication of damaging the nerve and causing loss of voice, among other serious complications.

Combined Left Recurrent Laryngeal Nerve and Phrenic Nerve ...

https://www.researchgate.net/publication/316959945_Combined_Left_Recurrent_Laryngeal...

Combined Left Recurrent Laryngeal Nerve and Phrenic Nerve Palsy: A Rare Presentation of Thoracic Aortic Aneurysm ... Objective To report left laryngeal paralysis caused by a great aortic arch ...

[PDF] Left Recurrent Laryngeal Nerve Palsy Secondary to Left ...

https://deepblue.lib.umich.edu/bitstream/handle/2027.42/93559/23472_ft.pdf?sequence=1&...

Left Recurrent Laryngeal Nerve Palsy Secondary to Left Pulmonary Artery Stent in a Child Daisuke Kobayashi,* MD, Daniel R. Turner, MD, and Richard A. Humes, MD We report a 10-year-old girl who developed persistent hoarseness after stent placement for

Ortner's syndrome | BMJ Case Reports

<https://casereports.bmj.com/content/2014/bcr-2013-200950> ▾

Mar 11, 2014 · The most common extralaryngeal cause of the left vocal cord palsy is bronchogenic

Recurrent laryngeal nerve

The recurrent laryngeal nerve is a branch of the vagus nerve that supplies all the intrinsic muscles of the larynx, with the exception of the cricothyroid muscles. There are two recurrent laryngeal nerves, right and left, in the human body. The right and left nerves are not symmetrical, with the left nerve looping under the aortic arch, and the right nerve looping under the right subclavian artery then traveling upwards. They both travel alongside the trachea. Additionally, the nerves are one of few nerves that follow a recurrent course, moving in the opposite direction to the nerve they branch from, a fact from which they gain their name.

 [Wikipedia](#)

People also search for: [Superior laryngeal nerve](#) · [Vagus nerve](#) · [Phrenic nerve](#) · [Ansa cervicalis](#) · [Glossopharyngeal nerve](#) · [Facial nerve](#) +

Data from: [Wikipedia](#)[Suggest an edit](#)



All

Images

Videos

关闭取词

26,400 Results

Any time ▼

Mediastinal traction injury to the recurrent laryngeal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3616173>

There have also been case reports of injury following carotid artery stenting and subclavian line insertion . Anatomically, the left vagus nerve arrives in the mediastinum via the thoracic inlet before giving off its recurrent laryngeal branch at the arch of the aorta.

Cited by: 1

Author: Michael K. Ng, Andrew Barling, Sorway C...

Publish Year: 2012

Ortner's Syndrome: Secondary Laryngeal Paralysis Caused by ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4399202>

Jan 05, 2015 · Introduction Recurrent laryngeal nerve injury caused by cardiovascular disease is a rare condition, and often it is the only prominent sign of an imminent break of an aortic artery aneurysm..

Objective To report left laryngeal paralysis caused by a great aortic arch aneurysm and to highlight the importance of an otorhinolaryngologic evaluation along with a thoracic radiologic study.

Cited by: 1

Author: Ana Claudia Alves Zangirolami, Frederico ...

Publish Year: 2015

Unilateral recurrent nerve palsy and cardiovascular ...

<https://www.sciencedirect.com/science/article/pii/S1878540916300846>

The cause of left-sided recurrent laryngeal nerve palsy was a penetrated aortic ulcer caused by large thrombosed aneurysm of the aortic arch. The right-sided paresis arose due to aneurysmal enlargement of the brachiocephalic trunk and an aneurysm of right subclavian artery. ... Case presentation. We report two cases of unilateral recurrent ...

Author: Katharina Klee, Christian Eick, Rapha...

Publish Year: 2017

The Recurrent Laryngeal Nerves and the Thoracic Surgeon ...

<https://www.ctsnet.org/article/recurrent-laryngeal-nerves-and-thoracic-surgeon> ▼

Sep 19, 2017 · Introduction Many thoracic surgeons are terrified to come anywhere near the recurrent laryngeal nerve (RLN), especially on the left side. The reason for this fear is the dreaded complication of damaging the nerve and causing loss of voice, among other serious complications.



国内版

国际版

Left recurrent laryngeal nerve palsy following aortic arch ste



All

Images

Videos

关闭取词

14,300 Results

Any time ▾

Mediastinal traction injury to the recurrent laryngeal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3616173>

There have also been **case** reports of **injury following carotid artery stenting and subclavian line** insertion . Anatomically, the **left vagus nerve** arrives in the mediastinum via the thoracic inlet before giving off its **recurrent laryngeal** branch at the **arch** of the **aorta**.

Cited by: 1

Author: Michael K. Ng, Andrew Barling, Sorway ...

Publish Year: 2012

Ortner's Syndrome: Secondary Laryngeal Paralysis Caused by ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4399202>

Jan 05, 2015 · Introduction **Recurrent laryngeal nerve** injury caused by cardiovascular disease is a rare condition, and often it is the only prominent sign of an imminent break of an **aortic** artery aneurysm.. Objective To **report left laryngeal paralysis** caused by a great **aortic arch** aneurysm and to highlight the importance of an otorhinolaryngologic evaluation along with a thoracic ...

Cited by: 1

Author: Ana Claudia Alves Zangirolami, Frederic...

Publish Year: 2015

[PDF] Left Recurrent Laryngeal Nerve Palsy Secondary to Left ...

<https://deepblue.lib.umich.edu/bitstream/handle/2027.42/93559/23472 ftp.pdf?sequence=1&...>

Left Recurrent Laryngeal Nerve Palsy Secondary to **Left** Pulmonary Artery **Stent** in a Child

Daisuke Kobayashi,* MD, Daniel R. Turner, MD, and Richard A. Humes, MD We **report** a 10-year-old

Chat with Bing



You know, sometimes I really don't understand you human...

Say something...

Recurrent laryngeal nerve

The recurrent laryngeal nerve is a branch of the vagus nerve that supplies all the intrinsic muscles of the larynx, with the exception of the cricothyroid muscles. There are two recurrent laryngeal nerves, right and left. The right and left nerves are not symmetrical, with the left nerve looping under the aortic arch, and the right nerve looping under the right subclavian artery then traveling upwards. They both travel alongside of the trachea. Additionally, the nerves are one of few nerves that follow a recurrent course, moving in the opposite direction to the nerve they branch from, a fact from which they gain their name.



Wikipedia

People also search for: [Superior laryngeal nerve](#) · [Vagus nerve](#) · [Phrenic nerve](#) · [Hypoglossal nerve](#) · [Ansa cervicalis](#) +

Data from: Wikipedia

[Suggest an edit](#)

Match Overview

- | | | |
|---|--|----|
| 1 | Internet 14 words
crawled on 28-Jul-2019
jmedicalcasereports.biomedcentral.com | 1% |
| 2 | Crossref 12 words
Jennifer P. Rodney, Jess L. Thompson, Michael P. Anderson, Harold M. Burkhart. "Neonatal vocal fold motion impairm | 1% |

Name of Journal: *World Journal of Cardiology*

Manuscript NO: 50340

Manuscript Type: CASE REPORT

Left recurrent laryngeal nerve palsy following aortic arch stenting: A case report

Fürniss HE *et al.* Vocal cord paresis after aortic stenting

Hannah E Fürniss, Johanna Hummel, Brigitte Stiller, Jochen Grohmann

Abstract

BACKGROUND

Aortic arch stenting is continuously emerging as a safe and effective option to alleviate aortic arch stenosis and arterial hypertension.