

¹¹
Name of journal: *World Journal of Cardiology*
Manuscript NO: 55079
Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Impact of cardiologist intervention on guideline-directed use of statin therapy

Cassagnol M *et al.* Guideline-directed use of statin therapy

Manouchkathé Cassagnol, Ofek Hai, Shaqeel A Sherali, Kyla D'Angelo, David Bass,
 Roman Zeltser, Amgad N Makaryus

Abstract

BACKGROUND

Match Overview

6 matches

1	Internet 88 words crawled on 24-Jan-2020 jamanetwork.com	3%
2	Internet 32 words crawled on 25-Mar-2019 journals.plos.org	1%
3	Internet 20 words crawled on 31-Jan-2020 www.wjgnet.com	1%
4	Internet 19 words crawled on 19-Oct-2009 cimed.ucr.ac.cr	1%
5	Internet 19 words crawled on 28-Jun-2020 www.acc.org	1%
6	Internet 19 words crawled on 10-Apr-2019 link.springer.com	1%
7	Crossref 14 words Mataaka, M.S.. "Is alternate daily dose of atorvastatin effective in treating patients with hyperlipidemia? The Alter ...	<1%
8	Crossref 13 words April Hinds, Debra Lopez, Karen Rascati, Jason Jokerst, Maaya Srinivasa. "Adherence to the 2013 Blood Chole: ...	<1%
9	Internet 10 words koara.lib.keio.ac.jp	<1%
10	Internet 10 words crawled on 27-Jun-2020 article.sciencepublishinggroup.com	<1%
11	Internet 10 words crawled on 20-Dec-2019 f6publishing.blob.core.windows.net	<1%



ALL

IMAGES

VIDEOS

3,130,000 Results

Any time ▾

Guideline-based statin use, adherence higher with ...

<https://www.healio.com/news/cardiology/20190805/...> ▾

Analysis of 1,249,061 Veterans Affairs patients with clinical ASCVD showed while 19.9% of patients overall were not prescribed statins for ASCVD, patients who visited a cardiologist had a 25%...

2019 ACC/AHA Guideline on the Primary Prevention of ...

<https://www.acc.org/latest-in-cardiology/ten...> ▾

- Scope of Guideline. The guideline is a compilation of the most important studies and guidelines for ...
- Assessment of ASCVD Risk. Assessment of ASCVD risk is the foundation of primary prevention. ...
- Estimating Risk of ASCVD. Electronic and paper chart risk estimators are available that utilize ...
- Nutrition. Dietary patterns associated with CVD mortality include—sugar, low-calorie sweeteners, ...

See full list on acc.org

Impact of Statin Therapy on Clinical Outcome in Patients ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4889205>

Most patients of the present study received low-intensity statin therapy probably because Japanese are known to have higher responsiveness to statin treatment compared with Caucasian.²² The 2013 American College of Cardiology/AHA guidelines recommend the use of higher dose and aggressive statin therapy in high-risk patients with clinically ...

Cited by: 13

Author: Masanobu Ishii, Koichi Kaikita, Koji Sato,...

Publish Year: 2016

Statin Use for the Primary Prevention of Cardiovascular ...

<https://www.aafp.org/afp/2017/0115/od1.html> ▾



Risk Factors For CVD

10-Year Risk of CVD Events

Screening and Stati



For the purposes of this recommendation, dyslipidemia is defined as an LDL-C level greater than 130 mg/dL or a high-density lipoprotein cholesterol (HDL-C) level less than 40 mg/dL (to convert HDL-C values to mmol/L, multiply by 0.0259). Most participants enrolled in trials of statin use for the prevention of CVD had an LDL-C level of 130 to 190 mg/dL or a diabetes diagnosis; hypertension and smoking were also common among trial participants.⁶ Persons with an LDL-C level greater than 190 mg/d...

See more on aafp.org



3,130,000 Results Any time ▾

[Guideline-based statin use, adherence higher with ...](#)

<https://www.healio.com/news/cardiology/20190805/...> ▾

Analysis of 1,249,061 Veterans Affairs patients with clinical ASCVD showed while 19.9% of patients overall were not prescribed statins for ASCVD, patients who visited a cardiologist had a 25%...

[Statin Use for the Primary Prevention of Cardiovascular ...](#)

<https://www.aafp.org/afp/2017/0115/od1.html> ▾

< **Risk Factors For CVD** 10-Year Risk of CVD Events Screening and Statir >

For the purposes of this recommendation, dyslipidemia is defined as an LDL-C level greater than 130 mg/dL or a high-density lipoprotein cholesterol (HDL-C) level less than 40 mg/dL (to convert HDL-C values to mmol/L, multiply by 0.0259). Most participants enrolled in trials of statin use for the prevention of CVD had an LDL-C level of 130 to 190 mg/dL or a diabetes diagnosis; hypertension and smoking were also common among trial participants.⁶ Persons with an LDL-C level greater than 190 mg/d...

[See more on aafp.org](#)

[Impact of Statin Therapy on Clinical Outcome in Patients ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4889205>

Most patients of the present study received low-intensity **statin therapy** probably because Japanese are known to have higher responsiveness to **statin** treatment compared with Caucasian.²² The 2013 American College of **Cardiology**/AHA guidelines recommend the **use** of higher dose and aggressive **statin therapy** in high-risk patients with clinically ...

Cited by: 13 **Author:** Masanobu Ishii, Koichi Kaikita, Koji Sato, ...

Publish Year: 2016

[2019 ACC/AHA Guideline on the Primary Prevention of ...](#)

<https://www.acc.org/latest-in-cardiology/ten...> ▾

- Scope of Guideline. The guideline is a compilation of the most important studies and guidelines for ...
- Assessment of ASCVD Risk. Assessment of ASCVD risk is the foundation of primary prevention. ...
- Estimating Risk of ASCVD. Electronic and paper chart risk estimators are available that utilize ...
- Nutrition. Dietary patterns associated with CVD mortality include: sugar, low-calorie sweeteners