

90854.docx

Name of Journal: World Journal of Cardiology

Manuscript Type: EDITORIAL

Mechanistic Insights into Fasting-Induced Autophagy in the Aging Heart

Hannaneh Parvaresh, Katarzyna Paczek, Abdul Alim Al-Bari, Nabil Eid

Abstract

Autophagy is a prosurvival mechanism for the clearance of accumulated abnormal proteins, damaged organelles, and excessive lipids within mammalian cells. A growing body of data indicates that autophagy is reduced in aging cells. This reduction leads to various diseases, such as myocardial hypertrophy, infarction, and atherosclerosis. Recent studies in animal models of an aging heart showed that fasting-induced autophagy improved cardiac function and longevity. This improvement is related to autophagic clearance of damaged cellular components via either bulk or selective autophagy (such as mitophagy). In this editorial, we summarize the mechanisms of autophagy in normal and aging hearts. In addition, the protective effect of fasting-induced autophagy in cardiac aging has been highlighted.

Key words: Aging; Autophagy; Heart; Fasting; Mitophagy.

Core tip: Autophagy is an essential mechanism for the clearance of harmful cellular components, which accumulate with age. However, autophagic machinery decreases with age, resulting in various diseases, such as cardiac hypertrophy. Recently, fasting-induced autophagy has been reported to improve cardiac function in animal models of aging via normalization of defective

autophagic machinery. Therefore, autophagy is an important target for the prevention of cardiac pathologies in the geriatric population.

INTRODUCTION

Cardiac Aging

Improvements in treatment procedures have contributed to increased life expectancy and growth in the aged population, especially in industrialized countries ^[1]. Aging is associated with a structural and functional decline in multiple organs, such as the heart. A sedentary lifestyle can also accelerate the incidence of aging-related diseases, including cardiovascular disease (CVD) ^[2-5].

Cardiovascular aging affects both the heart and the blood circulation system through slow and progressive alterations that can result in the development of left ventricular hypertrophy, diastolic dysfunction, coronary artery disease, stroke, hypertension, atherosclerosis, atrial fibrillation, and heart failure ^[6-9]. Aortic valve sclerosis is a valvulopathy associated with aging and is characterized by myxomatous degeneration, collagen deposition, and progression to aortic stenosis (AS) ^[10]. AS is an indicator of increased CVD risk and is mainly defined as increased leaflet calcification and decreased leaflet mobility ^[11]. Moreover, approximately 13–16% of elderly

people suffer from aortic regurgitation ^[12], which results in left ventricular dilation and dysfunction over time. Another valvular change related to aging is mitral annular calcification (MAC), which usually accompanies aortic valve sclerosis ^[13].

The free radical theory of aging and the mitochondrial theory have been suggested to explain the cellular deterioration observed in aging and suggest that the age-related decline in mitochondrial function and structure is a major driver of cardiomyocyte senescence, which causes endothelial dysfunction, alteration in the vasculature, and/or vascular injury ^[14].

Cellular senescence ² is activated following multiple stressors, including the elevation of reactive oxygen species; proinflammatory cytokines; and metabolic, mechanical, and chemical toxicity. Cellular senescence impairs the repair and regeneration of damaged cells in cardiovascular tissues ^[15-17].

Cellular senescence is characterized by genome instability, telomere attrition, and mitochondrial dysfunction ^[18].

Dysfunctional mitochondria produce less ATP while also generating increased amounts of reactive oxygen species (ROS) ^[19], exposing aged cardiomyocytes to high levels of oxidative stress. Autophagic and proteasomal degradation are the main mechanisms for the removal of damaged mitochondria and abnormal proteins in aged postmitotic cardiomyocytes. However, these mechanisms decline with age ^[20]. Eventually, when these mechanisms are unable to compensate for the accumulated cellular damage, stem-cell exhaustion and altered intercellular communication occur, further contributing to aging ^[18].

Autophagy in Cardiac Aging: Reduced Autophagy Accelerates Cardiac Aging

Autophagy activity is usually reduced with age ^[21]. A decrease in autophagy in the hearts of aged flies ^[22] and aged C57BL/6 mice (20–26 months old) has been reported ^[23, 24].

Autophagy is a protective housekeeping mechanism critical for cellular homeostasis and survival. Long-lived, damaged, and dysfunctional organelles; misfolded proteins; and invading pathogens are eliminated through this degradation process, providing building components for cellular renovation to effectively adapt cells to stressful conditions, such as nutrient deprivation, hypoxia, or oxidative stress ^[25, 26].

Autophagy can be selective or nonselective. Under starvation conditions, the protein functions non-selectively, and any cytoplasmic content can be targeted for catabolic recycling to maintain cellular energy production. However, there are also selective forms of autophagy that specifically target damaged organelles. For instance, mitophagy is a type of autophagy that selectively removes damaged mitochondria ^[27]. Mitochondria play a substantial role in cellular functions as well as cellular death. Thus, mitochondrial dysfunction is a crucial determinant of lifespan across species ^[28, 29].

Three types of autophagy have been recognized: macroautophagy, microautophagy, and chaperone-mediated autophagy, all of which lead to the turnover of intracellular components via various mechanisms. “Autophagy” is a term that generally refers to macroautophagy, which is the most prevalent form of autophagy^[30,31].

Molecular Machinery of Autophagy

Autophagy is initiated when several autophagy-related gene products (Atg1–Atg12) and other proteins are organized to form a phagophore. These proteins consist of at least five molecular components that mediate fusion between autophagosomes and lysosomes: (1) the Atg1/unc-51-like kinase (ULK) complex; (2) the Beclin 1/class III phosphatidylinositol 3-kinase (PI3K) complex; (3) Atg9 and vacuole membrane protein 1 (VMP1); (4) two ubiquitin-like proteins (Atg12 and Atg8/LC3) conjugation systems; and (5) proteins that mediate fusion between autophagosomes and lysosomes^[25, 32].

The initial step of autophagosome formation starts with Beclin1 (Atg6) and class III PI3K, which play crucial roles in vesicle isolation. Other Atg proteins are involved in Beclin-1-mediated formation of the Class III PI3K complex. In the next step, the autophagosome undergoes elongation via two conjugation systems. First, Atg12 is conjugated to Atg5 with the help of Atg7 and Atg10^[33, 34], followed by the conjugation of phosphatidylethanolamine to microtubule-associated protein 1 light chain 3 (LC3) via Atg4, Atg7 and Atg3. Consequently, the cytoplasmic LC3 (LC3-I) is converted to membranous (LC3-II) form, which is responsible for formation and maturation of the autophagosome^[35]. In the end, fusion of autophagosomes and lysosomes occurs with the formation of autolysosomes for degradation and recycling^[36].

The protein kinases mTOR and AMPK are implicated in the regulatory mechanisms of autophagy^[37]. Autophagy is inhibited by the mammalian target of rapamycin (mTOR). Phosphorylation of Ulk1 by AMPK is involved in autophagy promotion, although mTOR represses this process^[38]. Figure 1 demonstrates the various mechanisms of autophagy in mammalian cells.

Autophagy in the Heart

Accumulating evidence reveals that autophagy plays essential homeostatic roles in the heart under normal physiological conditions and during the aging process^[39]; additionally, it has an essential role in improving the immune response and reducing inflammation^[40]. Consequently, any perturbations to this process in the cardiovascular system can elicit harmful effects on health.

Autophagy attenuates with age and has serious implications for heart structure and function. A decrease in autophagy causes the development of heart failure, hypertension, atherosclerosis, and ischemic heart disease^[41].

Mitophagy is the selective autophagic clearance of damaged mitochondria and is crucial for the bioenergetics of the cardiovascular system; thus, mitophagy dysfunction is generally accompanied by cardiac disorders [27, 42, 43]. In addition, studies have suggested that autophagy decelerates cardiovascular senescence and has a positive effect on the healthy lifespan of animals [44-46].

Age-induced impairment of autophagy

Cardiomyocytes undergo age-related changes in proteostasis pathways, resulting in calcium homeostasis impairment, reactive oxygen species induction, hypertrophy and fibrosis, and eventual structural damage and diminished cardiac function. Moreover, with age, the MTOR-1 complex is significantly upregulated, and the AMPK pathway is downregulated. In addition, transcription factors involved in autophagy and lysosomal proteins such as TFEB and FOXO 3 are deactivated with advanced aging, resulting in reduced expression of autophagy genes [28-31].

Any defect in the autophagy process accelerates aging; likewise, aging is suppressed when autophagy is stimulated. Deletion of atg5, a cardiac-specific autophagy-related gene, in adult mice leads to an accelerated aging phenotype, including the development of cardiac hypertrophy, left ventricular dilatation, and contractile dysfunction [20, 47].

Mutations in the atg4c gene increase the risk of heart disease in elderly patients and eventually death [48]. Cardiomyocyte-specific deletion of glycogen synthase kinase-3 (GSK-3 α) in mice reduced basal autophagy levels and accelerated cardiac aging [49]. Dysfunction of autophagy with age slows the turnover of damaged proteasomes and contributes to age-associated CVD and cardiomyocyte senescence [50]. Mitophagy is impaired in aged mice, and mitophagy induction improves mitochondrial function and reduces arterial wall stiffness [51].

Acyl-coenzyme A binding protein (ACBP), which is encoded by a diazepine binding inhibitor (DBI), acts as an extracellular feedback inhibitor of autophagy [52]. It appears that high ACBP/DBI values correlate with future cardiovascular events (such as heart surgery, myocardial infarction, and stroke), suggesting that ACBP/DBI is indeed a biomarker of biological aging [39].

Mechanisms Underlying Age-Related Cardiac Remodeling: Involvement of Autophagy

Although there are many potential causes underlying the decline in cardiovascular function with age, a major determinant of the aging process is likely the progressive loss of quality control due to reduced autophagy.

Hyperactivation of mTOR and reduced AMPK activity^[53] in old age can directly inhibit autophagy by inactivating the pro-autophagic ULK1 (Unc-51 like autophagy activating kinase-1) complex^[54], contributing to the downregulation of autophagy activity.

It is conceivable that exposure to excessive ROS during aging promotes the accumulation of oxidized proteins, mitochondrial DNA mutations, and protein misfolding^[55]. Additionally, several cytosolic and mitochondrion-localized proteins involved in autophagy regulation become dysfunctional, thus contributing to abnormal mitochondrial turnover and the removal of damaged mitochondria^[56]. This chain of events results in impaired autophagy due to exhaustion of the aged autophagic machinery.

In addition, it has been proposed that a hallmark of aging in postmitotic cells, such as cardiomyocytes, is the aggregation of nondegradable structures inside lysosomes, termed lipofuscin, which impedes lysosomal function and therefore can likely inhibit autophagy^[57].

It has been shown that intracellular calcium has a key regulatory effect on cardiomyocyte autophagy. Inositol 1,4,5-trisphosphate (IP3) receptors mediate calcium release and transfer to mitochondria. This process inhibits autophagy by suppressing AMPK activation^[58]. Since evidence has shown that IP3 receptors are upregulated in the aged, hypertrophied, and failing myocardium of rodents^[59] and humans^[60], increased IP3 receptor-mediated calcium signaling likely exacerbates autophagy in the aging heart^[61].

Forkhead transcription factor (FOXO) and sirtuin (SIRT) proteins are also major metabolic regulators that mediate age-related vascular changes, particularly endothelial dysfunction^[9].

Dietary Activation of Autophagy in the Heart via Caloric Restriction or Fasting

Dietary interventions involving caloric restriction (CR) and fasting are among several stress stimuli that can induce autophagy in response to food deprivation^[62-64]. CR was defined as a reduction in caloric intake using a diet containing adequate amounts of protein, vitamins, and minerals^[65]. CR is a potent inducer of autophagy in the heart^[66], and its positive impacts on health and lifespan in various model organisms, primates and humans have been studied^[67-69]. CR is the most potent physiological stimulus of autophagy and ameliorates cardiac dysfunction (systolic and diastolic) and attenuates myocardial hypertrophy and fibrosis at the cardiomyocyte level. CR reduces mitochondrial damage, lipid accumulation, oxidative stress, apoptosis, telomere shortening, senescence marker levels, and circulating proinflammatory cytokine levels^[70].

Autophagy plays an important role in CR-mediated longevity^[71] via clearance of damaged mitochondria, reduction of oxidative stress, improvement of insulin sensitivity and suppression of inflammatory responses^[63, 64].

Short-term CR for 10 weeks in mice rejuvenated symptoms of the aging heart, such as significant improvement in diastolic function and regression of age-dependent cardiac hypertrophy^[72]. Moreover, CR reversed age-dependent cardiac proteome remodeling and mitigated oxidative damage and ubiquitination in these mice.

In aged animals, hypertrophy, and fibrosis, as well as systolic and diastolic dysfunctions, improved after CR^[70, 73]. The beneficial effects of CR observed in cardiomyocytes include enhanced mitochondrial fitness and reduced oxidative stress, apoptotic cell death, inflammation, and importantly, senescence^[70]. In vasculature, CR helps improve endothelial cell function and attenuates collagen deposition, elastin remodeling, and oxidative stress; as a result, CR reduces arterial stiffness^[74]. Another study¹² revealed improvements in numerous markers of cardiovascular health in humans after short-term periodic fasting, which is also a pro-autophagic dietary regimen^[75].

Intermittent fasting (IF) has attracted the attention of researchers as a dietary intervention associated with better compliance¹ and long-term adherence than CR in recent years^[76]. IF consists of regular cycles of times with no or minimal caloric intake interrupted by periods of normal food consumption¹. Alternate day fasting delays cardiac aging in rats, as determined by reduced hypertrophy and fibrosis^[77, 78] and extended lifespan^[79]. The advantageous effects of life-long alternate-day fasting were attributed to reduced phosphoinositide 3-kinase signaling, which was associated with reduced myocardial collagen deposition, oxidative stress, inflammatory markers, and B-type natriuretic peptide levels^[77, 80].

A fasting-mimicking diet (FMD) is considered another form of dietary intervention in which individuals consume low amounts of calories, sugars, and proteins but high amounts of unsaturated fats. Studies of FMD effects in mice have shown improved cognitive function and a rejuvenated immune system, in addition to promoting lifespan and health factors by reducing cancer incidence, obesity, and inflammation¹^[81]. FMD was investigated in humans, and the findings showed reduced age-related CVD risk factors, including reduced blood pressure, body mass index, fasting glucose, and inflammation, as well as an improved lipid profile^[82].

The efficacy of fasting on autophagy in the heart was assessed in male FBN rats by randomly dividing them into different groups of equal amounts of protein, vitamin, and mineral intake, while the CR groups received 20% less food from a 125% fortified diet for six weeks. Additionally, in addition to one simple CR group, two other CR groups were given 5 or 50 mg/kg/day

resveratrol. Compared with AL group, a marked reduction of expression of p62 (autophagy substrate) in the left ventricle was observed in the CR and Resv-50 rats, indicating enhanced cardiac autophagy in the CR group. Similarly, a significant overexpression of beclin-1 was found in the Resv-50 and CR animals. The CR + Resv-50 group of rats showed dramatically attenuated doxorubicin-induced damage, which can be due to enhanced autophagy [83]. Another study investigated the autophagic response of CR on diabetic rat hearts. Diabetic and nondiabetic rats were exposed to a CR diet (30% energy reduction) for 32 weeks. Compared with those of diabetic AL rats, diabetic CR rats exhibited an increase in the hepatic and cardiac LC3-II/LC3-I ratio (indicating enhanced autophagy) [84].

A high-fat diet (fat 60% kcal/100 kcal fat) was given to the FVB/N male mice for 4–20 weeks, after which they were subjected to overnight fasting to study the mechanisms of fasting-induced autophagy in the fatty mice heart [85]. After 24 hours of fasting, there was a significant conversion of LC3-I to LC3-II in lean mice heart but was not associated with a change in diet-induced obesity (DIO) mice. Furthermore, fasting suppressed mTOR in both lean and DIO mice, as indicated by increased AMPK phosphorylation and enhanced dephosphorylation of S6. Interestingly, mTOR inhibition was greater in obese mice. Taken together, these findings indicate that fasting activates autophagy in the hearts of lean mice [85].

Godar et al. investigated the impacts of IF on the autophagy–lysosome machinery in the myocardium. The authors studied the effects of fasting after 24 hours, followed by 24 hours of refeeding or 24 and 48 hours of fasting for six weeks [86]. The autophagosome abundance increased dramatically after 48 hours of fasting. Treatment with chloroquine (an autophagy inhibitor) was associated with a significant increase in LC3-II and SQSTM1/p62 after 24 hours of fasting but not in fed mice. Thus, fasting induces autophagy in cardiomyocytes; however, autophagy returns to basal levels on gestational days.

The effects of IF on right ventricular (RV) function in a rat model of pulmonary arterial hypertension (characterized by RV mitochondrial dysfunction and resultant lipotoxicity and microbiome dysbiosis) were explored. IF improved RV systolic and diastolic function and decreased RV cardiomyocyte hypertrophy and fibrosis, which was likely mediated by autophagy activation [87]. These protective effects could be related to autophagy activation.

Recent findings from studies also show that cardiometabolic parameters (e.g., adiposity, insulin sensitivity, and cardiac function) can be influenced by the time of day at which food is consumed [88]. To test the hypothesis that fasting during the sleep period elicits beneficial adaptation effects on cardiac function, wild-type mice were fasted for 24 hours or for either the 12-hour light/sleep phase or the 12-hour dark/awake phase. Repression of myocardial p-mTOR and protein synthesis occurred during the dark phase; both parameters remained elevated in the hearts of fasted mice during the light phase. In contrast, markers of autophagy (e.g., LC3-II) exhibited peak responses

to fasting during the light phase. Collectively, these data show that the responsiveness of the heart to fasting is temporally partitioned ^[88].

IF alleviated ² high-fat diet (HFD)-induced obesity cardiomyopathy in male C57BL/6J mice by improving cardiac functional and structural impairment and serum lipid metabolic disorders induced by HFD through decreasing lipid deposition, apoptosis and m6A methylation in the heart ^[89].

Researchers compared the effects of alternate-day fasting on elderly (aged 24 months) and young (aged 6 months) male rats. The results of this study indicated that alternate day fasting protected against inflammation and fibrosis in the heart during aging by inhibiting oxidative damage and NF- κ B activation ^[78]. Other studies have shown that fasting preconditioning activates AMPK, induces autophagy, decreases ROS levels, and inhibits NF- κ B signaling in the cardiac tissues of rats ^[90]. In addition, compared with fasting controls, IF in human subjects resulted in autophagy upregulation and reduced levels of proinflammatory cytokines, indicating the protective effects of fasting on the vascular system. This effect is most likely mediated by the anti-inflammatory effects of autophagy ^[91]. We investigated fasting-induced autophagy among large groups of population in the UAE during Ramadan (the holy Islamic fasting month). The results of this study will be revealed shortly. An accepted abstract related to this research will be presented at the Sharjah First International Conference on Fasting, February 28-29, 2024, held at Sharjah University, United Arab Emirates ^[92].

CONCLUSIONS

In conclusion, fasting-induced autophagy is beneficial for ensuring cardiac function, preventing disease, and improving longevity. However, additional studies in vivo in animal models of cardiac aging are needed to determine the specific molecular mechanisms involved in normalizing autophagy by fasting. In addition, large-scale studies on humans are needed. Ramadan fasting, a type of IF (a common religious practice) in Islamic countries, could be investigated in large groups of geriatric people with or without cardiac diseases. Importantly, further in vitro research should be directed toward human cardiac tissues to better understand the molecular mechanisms of fasting-induced autophagy and its beneficial effects on longevity pathways and prevention of cardiovascular diseases.

16%

SIMILARITY INDEX

PRIMARY SOURCES

- 1

Mahmoud Abdellatif, Simon Sedej, Didac Carmona-Gutierrez, Frank Madeo, Guido Kroemer. "Autophagy in Cardiovascular Aging", Circulation Research, 2018

Crossref

113 words — 4%
- 2

www.ncbi.nlm.nih.gov

Internet

108 words — 3%
- 3

www.science.gov

Internet

78 words — 2%
- 4

link.springer.com

Internet

49 words — 2%
- 5

www.researchgate.net

Internet

40 words — 1%
- 6

Leon, Leonardo J., and Åsa B. Gustafsson. "Staying young at heart: autophagy and adaptation to cardiac aging", Journal of Molecular and Cellular Cardiology, 2015.

Crossref

34 words — 1%
- 7

Vanessa Obas, Ramachandran S. Vasan. "The aging heart", Clinical Science, 2018

Crossref

18 words — 1%
- 8

www.fightaging.org

Internet

16 words — < 1%

9	La Rovere, Rita M.L., Gemma Roest, Geert Bultynck, and Jan B. Parys. "Intracellular Ca ²⁺ signaling and Ca ²⁺ microdomains in the control of cell survival, apoptosis and autophagy", Cell Calcium, 2016. <small>Crossref</small>	14 words — < 1%
10	assets.cureus.com <small>Internet</small>	13 words — < 1%
11	ia601901.us.archive.org <small>Internet</small>	13 words — < 1%
12	Mahmoud Abdellatif, Senka Ljubojevic-Holzer, Frank Madeo, Simon Sedej. "Autophagy in cardiovascular health and disease", Elsevier BV, 2020 <small>Crossref</small>	12 words — < 1%
13	www.mdpi.com <small>Internet</small>	12 words — < 1%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 12 WORDS
EXCLUDE MATCHES < 12 WORDS