

Match Overview

| | | |
|---|---|-----|
| 1 | CrossCheck 259 words Cabanas-Grandio, P., E. Abu-Assi, S. Raposeiras-Roubin, B. Alvarez-Alvarez, C. Gonzalez-Cambeiro, S. G. Romani, | 3% |
| 2 | CrossCheck 93 words Abumuaileq, R. R.-Y., E. Abu-Assi, S. Raposeiras-Roubin, A. Lopez-Lopez, A. Redondo-Dieguez, D. Alvarez-Iglesias, | 1% |
| 3 | Internet 89 words crawled on 08-Jan-2013 www.glucose.sextreffnorge.com | 1% |
| 4 | Internet 67 words crawled on 23-Apr-2015 cjasn.asnjournals.org | 1% |
| 5 | Internet 42 words crawled on 13-Feb-2015 circ.ahajournals.org | 1% |
| 6 | Internet 27 words crawled on 08-Dec-2013 beckerinfo.net | <1% |
| 7 | Internet 24 words crawled on 25-Feb-2015 circheartfailure.ahajournals.org | <1% |
| 8 | Internet 24 words crawled on 17-Feb-2014 www.revistanefrologia.com | <1% |
| 9 | Internet 23 words crawled on 18-May-2015 www.researchgate.net | <1% |

40

Name of Journal: World Journal of Cardiology

ESPS Manuscript NO: 20651

Manuscript Type: Original Article

Retrospective Cohort Study

47

Renal function assessment in atrial fibrillation: Usefulness of chronic kidney disease epidemiology collaboration vs re-expressed 4 variable modification of diet in renal disease

2

Rami Riziq-Yousef Abumuaileq, Emad Abu-Assi, Andrea López-López, Sergio Raposeiras-Roubin, Moisés Rodríguez-Mañero, Luis Martínez-Sande, Francisco Javier García-Seara, Xesus Alberte Fernandez-López, Jose Ramón González-Juanatey

Abstract

AIM: To compare the performance of the re-expressed Modification of Diet In Renal Disease equation vs the new Chronic Kidney Disease Epidemiology Collaboration equation in patients with non-valvular atrial fibrillation.



Renal function assessment in atrial fibrillation: usefulness of CKD-EPI ve



网页

新闻

图片

视频

更多 ▾

搜索工具

找到约 2,900,000 条结果 (用时 0.85 秒)

Performance of the Cockcroft-Gault, MDRD and CKD-EPI ...

www.jafib.com/published/published.php?id...is=previous... ▾ 翻译此页

作者: S Barra - 被引用次数: 1 - 相关文章

Atrial fibrillation (AF) is the most common sustained arrhythmia and associates ... In the ROCKET-AF and ATRIA study cohorts, impaired renal function was ... of the re-expressed 4-variable MDRD, Cockcroft-Gault and CKD-EPI formulae in a ... assessing the severity of suspected or previously known mitral valve disease (19 ...

Cross-Sectional Evaluation of Kidney Function in ... - Karger

www.karger.com/Article/FullText/355813 ▾ 翻译此页

作者: D Santoro - 2014 - 相关文章

2014年12月19日 - The most common methods of assessment of kidney function are the estimation ... The CKD-EPI (Chronic Kidney Disease Epidemiology ... Glomerular filtration rate was estimated by using the re-expressed 4-variable MDRD Study with warfarin in relation to renal function in patients with atrial fibrillation: ...

Patient safety and estimation of renal function in patients ...

bmjopen.bmj.com/content/3/9/e003343.full ▾ 翻译此页

作者: PK MacCallum - 2013 - 被引用次数: 29 - 相关文章

2013年9月27日 - Given the potentially widespread use of these agents, particularly in ... For each patient in the study, renal function was calculated using both ... and better-performing MDRD formula and a newer CKD-EPI formula ... Dabigatran versus warfarin in patients with atrial fibrillation. Re:Reply to Rapuch et al.

Renal function estimations and dose recommendations for ...

bmjopen.bmj.com/content/3/4/e002686.full ▾ 翻译此页



Renal function assessment in atrial fibrillation: Usefulness of chronic ki

网页

图片

新闻

视频

更多 ▾

搜索工具

找到约 71,500 条结果 (用时 0.78 秒)

Performance of the Cockcroft-Gault, MDRD and CKD-EPI ...www.jafib.com/published/published.php?id...is=previous... ▾ 翻译此页

作者: S Barra - 被引用次数: 1 - 相关文章

Renal dysfunction is a strong predictor of adverse events in patients with ... The Cockcroft-Gault, Modification of Diet in Renal Disease (MDRD) and Chronic Kidney Disease-Epidemiology Collaboration (CKD-EPI) equations are available for ... of the re-expressed 4-variable MDRD, Cockcroft-Gault and CKD-EPI formulae in a ...

[PDF] Performance of the Cockcroft-Gault, MDRD and CKD-EPI ...www.jafib.com/published/webFormat/Sergio/sergio.pdf ▾ 翻译此页

作者: S Barra - 被引用次数: 1 - 相关文章

Modification of Diet in Renal Disease (MDRD) and Chronic Kidney Disease-Epidemiology ... AUC=0.586±0.054, p=0.041) and CKD-EPI (AUC=0.673±0.049 vs. formulae for the estimation of GFR (Cockcroft-Gault, re-expressed Chronic Kidney Disease Epidemiology Collaboration equation; AF → atrial fibrillation; TIA →

Expressing the Modification of Diet in Renal Disease Study ...www.clinchem.org/content/53/4/766.full ▾ 翻译此页

作者: AS Levey - 2007 - 被引用次数: 853 - 相关文章

2007年3月1日 - The reexpressed 4-variable MDRD Study equation for Scr (mg/dL) is GFR = 175 ... To facilitate detection, evaluation, and management of chronic kidney disease (CKD),1 ... the 4-variable (modified) MDRD Study equation (6) for use with ... Chronic Kidney Disease Epidemiology Collaboration is a research ...

Cross-Sectional Evaluation of Kidney Function in ... - Karger<https://www.karger.com/Article/FullText/355813> - 翻译此页

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 71,200 条结果 (用时 0.84 秒)

Performance of the Cockcroft-Gault, MDRD and CKD-EPI ...

www.jafib.com/published/published.php?id...is=previous... ▾ 翻译此页

作者: S Barra - 被引用次数: 1 - 相关文章

Renal dysfunction is a strong predictor of adverse events in patients with ... The Cockcroft-Gault, Modification of Diet in Renal Disease (MDRD) and Chronic Kidney Disease-Epidemiology Collaboration (CKD-EPI) equations are available for ... of the re-expressed 4-variable MDRD, Cockcroft-Gault and CKD-EPI formulae in a ...

[PDF] Performance of the Cockcroft-Gault, MDRD and CKD-EPI ...

www.jafib.com/published/webFormat/Sergio/sergio.pdf ▾ 翻译此页

作者: S Barra - 被引用次数: 1 - 相关文章

Modification of Diet in Renal Disease (MDRD) and Chronic Kidney Disease-Epidemiology ... recommended, the re-expressed version of the MDRD formula was ... 4-variable MDRD, Cockcroft-Gault and CKD-EPI formulae in a ... Chronic Kidney Disease Epidemiology Collaboration equation; AF – atrial fibrillation; TIA –

Expressing the Modification of Diet in Renal Disease Study ...

www.clinchem.org/content/53/4/766.full ▾ 翻译此页

作者: AS Levey - 2007 - 被引用次数: 853 - 相关文章

2007年3月1日 - The reexpressed 4-variable MDRD Study equation for Scr (mg/dL) is ... Chronic Kidney Disease Epidemiology Collaboration is a research ... assay vs NIST-assigned creatinine values using LC-IDMS on CAP LN24 survey samples. ... to Baseline Renal Function in Patients With Atrial Fibrillation: A RE-LY ...

Cross-Sectional Evaluation of Kidney Function in ... - Karger

<https://www.karger.com/Article/FullText/355813> - 翻译此页

作者: D Santoro - 2014 - 相关文章

2014年12月19日 - The incidence of chronic kidney disease (CKD) continues to increase