

Name of Journal: *World Journal of Biological Chemistry*

ESPS Manuscript NO: 31747

Manuscript Type: Editorial

A common therapeutic target for both cancer and obesity

Yie-Hwa Chang

Abstract

Obesity and cancer are two interrelated conditions of high epidemiological need, with studies showing that obesity is responsible for nearly 25% of the relative contribution to cancer incidence. Given the connection between these conditions, a drug that can

Match Overview

1	Crossref 97 words Arico-Muendel, C. C., B. Belanger, D. Benjamin, H. S. Blanchette, T. M. Caiazzo, P. A. Centrella, J. DeLorey, E.	4%
2	Internet 96 words crawled on 18-Dec-2016 www.dovepress.com	4%
3	Internet 73 words crawled on 06-Sep-2016 www.univerciencia.org	3%
4	Crossref 61 words Kim, Dennis D., Janakan Krishnarajah, Stephen Lillioja, Ferdinandus de Looze, Jo Marjason, Joseph Proietto, S	2%
5	Internet 51 words crawled on 23-Mar-2010 www.e-medicum.com	2%
6	Crossref 47 words Sang Joon Shin. "A Phase Ib pharmacokinetic study of the anti-angiogenic agent CKD-732 used in combination	2%

找到约 35,000 条结果 (用时 0.75 秒)

Google 学术: Roles of Methionine aminopeptidase in Cancer and Obesity

Inhibition of the methionine aminopeptidase 2 enzyme ... - Joharapurkar - 被引用次数: 20

Methionine aminopeptidase-2 as a selective target of ... - Kass - 被引用次数: 13

... and differentiation, in tumor suppression, and in obesity - Datta - 被引用次数: 7

METAP2 - Wikipedia

<https://en.wikipedia.org/wiki/METAP2> ▼ 翻译此页

Methionine aminopeptidase 2 is an enzyme that in humans is encoded by the METAP2 gene. ... is necessary for the progression of diseases including solid tumor cancers and rheumatoid arthritis. ... While previous studies have indicated MetAP2 catalyzes the removal of N-terminal methionine residues in vitro, the function ...

Inhibition of the methionine aminopeptidase 2 enzyme for the ... - NCBI

[https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral\(PMC\)](https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC)) - 翻译此页

作者: AA Joharapurkar - 2014 - 被引用次数: 20 - 相关文章

2014年2月28日 - The main function of WAT is to store energy in the form of lipids, whereas the ...

Methionine aminopeptidases (MetAP2) in angiogenesis therapy for cancer, MetAP2 inhibition for obesity treatment might not lead to drug ...

Inhibition of the methionine aminopeptidase 2 enzyme for the ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/24611021> - 翻译此页

作者: AA Joharapurkar - 2014 - 被引用次数: 20 - 相关文章

2014年2月28日 - Worldwide prevalence of obesity has nearly doubled since 1980. ... anti-cancer agents,

methionine aminopeptidase (MetAP2) inhibitors induce ...



Common therapeutic target for both cancer and obesity



全部

新闻

图片

购物

视频

更多

设置

工具

找到约 3,670,000 条结果 (用时 0.90 秒)

Google 学术: Common therapeutic target for both cancer and obesity

... hypoxia-inducible factor 1 α in common human cancers ... - Zhong - 被引用次数: 2643

Diabetes and cancer: a consensus report - Giovannucci - 被引用次数: 1313

World cancer report 2014 - Stewart - 被引用次数: 946

Adipose tissue angiogenesis as a therapeutic target for obesity and ...

www.nature.com > [Journal home](#) > [Archive](#) > [Perspectives](#) > [Opinion](#) ▾ [翻译此页](#)

作者: Y Cao - 2010 - 被引用次数: 233 - [相关文章](#)

Molecular mechanisms of switching on angiogenic phenotypes in both healthy ... shows that pathological angiogenesis is regulated by common molecular mechanisms. The original hypothesis that angiogenesis could be a target for obesity ... Clinical experiences with anti-angiogenic drugs for cancer therapy such as ...

Cellular Bioenergetics as a Target for Obesity Therapy - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: YH Tseng - 2010 - 被引用次数: 294 - [相关文章](#)

Cellular Bioenergetics as a Target for Obesity Therapy This cycling is involved in the thermogenesis associated with burn injuries, cancer cachexia, and ... DIT and NST share common features in that both occur in brown fat and both are ...

Diabetes and cancer: two diseases with obesity as a common risk factor

[全部](#)[新闻](#)[图片](#)[购物](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 3,540,000 条结果 (用时 0.69 秒)

Google 学术: Common therapeutic target for both cancer and obesity

... hypoxia-inducible factor 1 α in common human cancers ... - Zhong - 被引用次数: 2652

Diabetes and cancer: a consensus report - Giovannucci - 被引用次数: 1326

World cancer report 2014 - Stewart - 被引用次数: 983

Diabetes and cancer: two diseases with obesity as a common risk factor

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: SK Garg - 2014 - 被引用次数: 53 - 相关文章

2013年6月12日 - In fact, both obesity and T2DM have been shown to be ... by the multidrug therapy required for the treatment of diabetes, obesity and cancer. levels of IGF-1 reduce apoptosis and increase cell proliferation in target cells, ...

Adipose tissue angiogenesis as a therapeutic target for obesity and ...

www.nature.com > Journal home > Archive > Perspectives > Opinion ▼ 翻译此页

作者: Y Cao - 2010 - 被引用次数: 238 - 相关文章

Molecular mechanisms of switching on angiogenic phenotypes in both healthy ... shows that pathological angiogenesis is regulated by common molecular mechanisms. The original hypothesis that angiogenesis could be a target for obesity Clinical experiences with anti-angiogenic drugs for cancer therapy such as ...

British Journal of Cancer - From obesity to diabetes and cancer ...

www.nature.com > Journal home > Archive > Minireview - 翻译此页

作者: C García-Jiménez - 2016 - 被引用次数: 11 - 相关文章

2016年2月23日 - Given the global epidemic of diabetes, preventing both cancer occurrence in ... and both common and independent risk factors may contribute to cancer Two major therapies targeting cancer growth or survival have been ...