



**Name of Journal:** *World Journal of Clinical Oncology*

**ESPS Manuscript NO:** 31729

**Manuscript Type:** Case Report

**Target migration from re-inflation of adjacent atelectasis during lung stereotactic body radiotherapy**

**Bijing Mao, Vivek Verma, Dandan Zheng, Xiaofeng Zhu, Nathan R Bennion, Abhijeet R Bhirud, Maria A Poole, Weining Zhen**

#### Abstract

Stereotactic body radiotherapy (SBRT) is a widely accepted option for the treatment of medically inoperable early-stage non-small cell lung cancer

#### Match Overview

1	Internet 39 words crawled on 10-Nov-2014 <a href="http://worldwidescience.org">worldwidescience.org</a>	2%
2	Internet 17 words crawled on 20-Sep-2014 <a href="http://effiloop.com">effiloop.com</a>	1%
3	Internet 16 words crawled on 22-Mar-2016 <a href="http://www.eunethta.eu">www.eunethta.eu</a>	1%
4	Internet 13 words crawled on 07-Sep-2016 <a href="http://www.osti.gov">www.osti.gov</a>	1%
5	Internet 13 words crawled on 24-Nov-2006 <a href="http://www.cshs.org">www.cshs.org</a>	1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 9,340 条结果 (用时 0.37 秒)

[S. Siva - IASLC | International Association for the Study of Lung Cancer](#)  
[library.iaslc.org/search-speaker?search\\_speaker=18248](#) ▾ [翻译此页](#)

Reinflation of atelectatic lung in response to tumour size reduction may require .... to stereotactic ablative body radiotherapy (SABR) for lung cancer patients in Asia. .... Two markers were placed in adjacent airways in one patient, and the ... No marker migration was observed over the treatment time for both marker types.

[Radiotherapy - IASLC | International Association for the Study of Lung ...](#)  
[library.iaslc.org/virtual-library-search?product\\_id=6...](#) [Radiotherapy](#) ▾ [翻译此页](#)

Recent Advances in Radiation Treatment Technique for Lung Cancer Lung cancer ... early cancer and targeted correctly by stereotactic body radiation therapy (SBRT). .... Advances in Stereotactic Body Radiotherapy Matthias Guckenberger, ..... to assess the toxicity and outcome of SBRT re-irradiation (reRT) in patients with ...

[\[PDF\] acosog z4099 / rtog 1021 - Advanced Technology Consortium](#)  
[atc.wustl.edu/protocols/rtog/1021/Z4099-1021\\_A0\\_05-02-2011.pdf](#) ▾ [翻译此页](#)

2011年5月2日 - Brachytherapy) versus Stereotactic Body Radiation Therapy in High Risk Patients with Stage I Non-Small Cell Lung ..... the edge of the target resulting in a disappointing 32% Kaplan-Meier local ..... After securing brachytherapy implant and with lung reinflation, positioning of the sources should be verified.

[Lung deformations and radiation-induced regional lung collapse in ...](#)  
[https://www.ncbi.nlm.nih.gov/pubmed/26520737](#) - [翻译此页](#)

作者: Q Diot - 2015 - [相关文章](#)

Lung deformations and radiation-induced regional lung collapse in patients treated with stereotactic body radiation therapy. Diot Q(1), Kavanagh B(1), ...

缺少字词: target migration re inflation adjacent

[\[PDF\] The Management of Respiratory Motion in Radiation Oncology - AA...](#)

[All](#)[Images](#)[News](#)[Videos](#)[Shopping](#)[More](#)[Settings](#)[Tools](#)

About 52,000 results (0.67 seconds)

[S. Siva - IASLC | International Association for the Study of Lung Cancer](#)[library.iaslc.org/search-speaker?search\\_speaker=18248](https://library.iaslc.org/search-speaker?search_speaker=18248) ▼

Reinflation of atelectatic lung in response to tumour size reduction may require .... to stereotactic ablative body radiotherapy (SABR) for lung cancer patients in Asia. .... Two markers were placed in adjacent airways in one patient, and the ... No marker migration was observed over the treatment time for both marker types.

[D. Ball - IASLC | International Association for the Study of Lung Cancer](#)[library.iaslc.org/search-speaker?search\\_speaker=15505](https://library.iaslc.org/search-speaker?search_speaker=15505) ▼

Reinflation of atelectatic lung in response to tumour size reduction may require .... to detect tumor and normal tissue response to radiotherapy during treatment. ... for targeted preventative strategies to reduce LC-associated TE, and in .... Quality Assurance for Lung Cancer Stereotactic Ablative Body Radiotherapy (ID 2559).

[Radiotherapy - IASLC | International Association for the Study of Lung ...](#)[library.iaslc.org/virtual-library-search?product\\_id=6&author...Radiotherapy](https://library.iaslc.org/virtual-library-search?product_id=6&author...Radiotherapy) ▼

Advances in Stereotactic Body Radiotherapy Matthias Guckenberger, ... Stereotactic Body Radiation Therapy for Early-Stage Non-Small Cell Lung Cancer. .... SC03.05 - Radiotherapy Combined with Targeted Therapies or ..... This study aimed to assess the toxicity and outcome of SBRT re-irradiation (reRT) in patients with ...

[The management of respiratory motion in radiation oncology report of ...](#)[onlinelibrary.wiley.com/doi/10.1118/1.2349696/full](https://onlinelibrary.wiley.com/doi/10.1118/1.2349696/full)

by PJ Keall - 2006 - Cited by 1235 - Related articles

Lung cancer accounts for 28% of all cancer deaths in the U.S. (American .... the leaves of a multileaf collimator (MLC) and the component of target motion ... The intercostal muscles connect adjacent ribs and also participate in normal inhalation. .... single or few-fraction treatments common for stereotactic body radiotherapy.



[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 66,100 条结果 (用时 0.61 秒)

## S. Siva - IASLC | International Association for the Study of Lung Cancer

[library.iaslc.org/search-speaker?search\\_speaker=18248](http://library.iaslc.org/search-speaker?search_speaker=18248) ▾ [翻译此页](#)

Reinflation of atelectatic lung in response to tumour size reduction may require .... to stereotactic ablative body radiotherapy (SABR) for lung cancer patients in Asia. .... Two markers were placed in adjacent airways in one patient, and the ... No marker migration was observed over the treatment time for both marker types.

## [PDF] acosog z4099 / rtog 1021 - Advanced Technology Consortium

[atc.wustl.edu/protocols/rtog/1021/Z4099-1021\\_A0\\_05-02-2011.pdf](http://atc.wustl.edu/protocols/rtog/1021/Z4099-1021_A0_05-02-2011.pdf) ▾ [翻译此页](#)

2011年5月2日 - Brachytherapy) versus Stereotactic Body Radiation Therapy in High Risk Patients with Stage I Non-Small Cell Lung ..... the edge of the target resulting in a disappointing 32% Kaplan-Meier local ..... After securing brachytherapy implant and with lung reinflation, positioning of the sources should be verified.

## The management of respiratory motion in radiation oncology report of ...

[onlinelibrary.wiley.com/doi/10.1118/1.2349696/full](http://onlinelibrary.wiley.com/doi/10.1118/1.2349696/full) - [翻译此页](#)作者: PJ Keall - 2006 - 被引用次数: 1246 - [相关文章](#)

Lung cancer accounts for 28% of all cancer deaths in the U.S. (American .... the leaves of a multileaf collimator (MLC) and the component of target motion ... The intercostal muscles connect adjacent ribs and also participate in normal inhalation. .... single or few-fraction treatments common for stereotactic body radiotherapy.

## Pet/Ct Radiotherapy Planning - European Association of Nuclear ...

[www.eanm.org/content.../gl\\_PET-CT\\_Radiotherapy\\_Planning\\_Part\\_3.pdf](http://www.eanm.org/content.../gl_PET-CT_Radiotherapy_Planning_Part_3.pdf) - [翻译此页](#)

Section 4 – Future Possibilities for PET/CT in Radiotherapy Planning. 86 .... control by escalating radiation dosages to re- ... invade adjacent tissues and thence travel ..... Figure 3: Dose-volume histograms for the planning target volume (PTV) red ... tion of rotational IMRT for treatment of lung