

Name of Journal: *World Journal of Clinical Oncology*

Manuscript NO: 31714

Manuscript Type: Case Report

Charcot-Marie-Tooth hereditary neuropathy revealed after administration of docetaxel in advanced breast cancer

Hampig Raphael Kourie, Nicolas Mavroudakis, Philippe Aftimos, Martine Piccart

Abstract

Charcot-Marie-Tooth (CMT) neuropathy is the most common hereditary cause of neuropathy. Diagnosis is usually not made during the childhood but in adolescence or late adulthood. It is reported in the literature that some neurotoxic chemotherapeutical agents can reveal an asymptomatic CMT IA hereditary neuropathy. To our knowledge, we report here the first case of CMT IA revealed in a 55 year-old woman after the administration of docetaxel/trastuzumab/perituzumab for metastatic breast cancer. This case

Match Overview

- | | | |
|---|--|----|
| 1 | Crossref 25 words
Carla C P Verstappen. "Neurotoxic Complications of Chemotherapy in Patients with Cancer", Drugs, 2003 | 2% |
| 2 | Internet 25 words
crawled on 28-Oct-2015
www.ncbi.nlm.nih.gov | 2% |
| 3 | Crossref 14 words
William D. Graf. "Severe vincristine neuropathy in Charcot-Marie-Tooth disease type 1A", Cancer, 04/01/199 | 1% |

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 33,400 条结果 (用时 0.81 秒)

[\[PDF\] Full-Text PDF - MDPI.com](#)

www.mdpi.com/2305-6304/3/2/152/pdf ▾ [翻译此页](#)

作者: R Velasco - 2015 - [被引用次数: 3](#) - [相关文章](#)

2015年4月28日 - neoadjuvant **breast cancer** patients, dose-modification of taxane planned women with established **docetaxel**-induced peripheral neuropathy ... recordings before and **after** taxane **administration**, with paclitaxel patients with the hereditary neuropathy **Charcot-Marie-Tooth** (CMT) being poorly tolerant of.

[Chemotherapy-induced polyneuropathy. Part I. Pathophysiology - NCBI](#)

<https://www.ncbi.nlm.nih.gov> ▸ [NCBI](#) ▸ [Literature](#) ▸ [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: K Brzeziński - 2012 - [被引用次数: 10](#) - [相关文章](#)

2012年2月29日 - In most cases, they are **administered** in multi-drug schemes, while **After** treatment discontinuation, the symptoms disappear slowly (up to Severe vincristine neuropathy in **Charcot-Marie-Tooth** disease type 1A. in patients with **advanced breast cancer** pretreated with paclitaxel – a phase II study.

[Presentation and management of docetaxel-related adverse effects in ...](#)

<https://www.ncbi.nlm.nih.gov> ▸ [NCBI](#) ▸ [Literature](#) ▸ [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: MY Ho - 2014 - [被引用次数: 8](#) - [相关文章](#)

2014年5月27日 - **Maria** Y Ho and John R Mackey ... A large retrospective study published by the British Columbia Agency **Breast Tumor** Group **revealed** that patients who were For those patients with HER-2-positive **metastatic breast cancer**, **docetaxel** Extended benefit from sequential **administration of docetaxel after** ...

缺少字词: ~~ehareet tooth~~

[Chemotherapy-induced peripheral neurotoxicity: A critical analysis ...](#)

onlinelibrary.wiley.com/doi/10.3322/caac.21204/full ▾ [翻译此页](#)

作者: SB Park - 2013 - [被引用次数: 87](#) - [相关文章](#)

2013年11月11日 - Symptoms may continue to worsen even **after** treatment has ceased, [132] In

找到约 53,000 条结果 (用时 0.65 秒)

Charcot-Marie-Tooth gene, SBF2, associated with taxane-induced ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5347688/> - 翻译此页

作者: BP Schneider - 2016 - 被引用次数: 1 - 相关文章

2016年10月9日 - **Advanced** - Journal list - Help ... Taxane-induced peripheral **neuropathy** (TIPN) is one of the most ... paclitaxel in the adjuvant, randomized phase III **breast cancer** trial, ... **After** removing variants with missing rate > 20% and those which ... a subtype of **Charcot-Marie-Tooth** (CMT), a known **hereditary** form of ...

Chemotherapy-induced polyneuropathy. Part I. Pathophysiology

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: K Brzeziński - 2012 - 被引用次数: 11 - 相关文章

2012年2月29日 - Chemotherapy-induced peripheral **neuropathy** (CIPN) is a toxic from the body within 96 hours **after** cessation of drug **administration** Severe vincristine **neuropathy** in **Charcot-Marie-Tooth** disease type 1A. **Cancer**. 1996;77:1356-62 patients with **advanced breast cancer** pretreated with paclitaxel - a ...

[PDF] Taxane-Induced Peripheral Neurotoxicity - MDPI

www.mdpi.com/2305-6304/3/2/152/pdf ▼ 翻译此页

作者: R Velasco - 2015 - 被引用次数: 4 - 相关文章

2015年4月28日 - neoadjuvant **breast cancer** patients, dose-modification of taxane planned women with established **docetaxel**-induced peripheral neuropathy ... recordings before and **after** taxane **administration**, with paclitaxel patients with the **hereditary neuropathy Charcot-Marie-Tooth** (CMT) being poorly tolerant of.

The administration of chemotherapy in a patient with Charcot-Marie ...

https://www.researchgate.net/.../7875374_The_administration_of_chemother... - 翻译此页

She developed a distal sensory and motor neuropathy **after** her first ... Upon transfer of care to our



全部

新闻

图片

视频

购物

更多

设置

工具

找到约 53,300 条结果（用时 0.76 秒）

Charcot-Marie-Tooth gene, SBF2, associated with taxane-induced ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5347688/> - 翻译此页

作者：BP Schneider - 2016 - 被引用次数：1 - 相关文章

2016年10月9日 - **Advanced** - Journal list - Help ... Taxane-induced peripheral **neuropathy** (TIPN) is one of the most ... paclitaxel in the adjuvant, randomized phase III **breast cancer** trial, ... **After** removing variants with missing rate > 20% and those which ... a subtype of **Charcot-Marie-Tooth** (CMT), a known **hereditary** form of ...

Chemotherapy-induced polyneuropathy. Part I. Pathophysiology

[https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral\(PMC\)](https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC)) - 翻译此页

作者：K Brzeziński - 2012 - 被引用次数：11 - 相关文章

2012年2月29日 - Chemotherapy-induced peripheral **neuropathy** (CIPN) is a toxic from the body within 96 hours **after** cessation of drug **administration** Severe vincristine **neuropathy** in **Charcot-Marie-Tooth** disease type 1A. *Cancer*. 1996;77:1356-62 patients with **advanced breast cancer** pretreated with paclitaxel - a ...

[PDF] Taxane-Induced Peripheral Neurotoxicity - MDPI

www.mdpi.com/2305-6304/3/2/152/pdf - 翻译此页

作者：R Velasco - 2015 - 被引用次数：4 - 相关文章

2015年4月28日 - neoadjuvant **breast cancer** patients, dose-modification of taxane planned women with established **docetaxel**-induced peripheral neuropathy ... recordings before and **after** taxane **administration**, with paclitaxel patients with the **hereditary neuropathy Charcot-Marie-Tooth** (CMT) being poorly tolerant of.

The administration of chemotherapy in a patient with Charcot-Marie ...

https://www.researchgate.net/.../7875374_The_administration_of_chemother... - 翻译此页

She developed a distal sensory and motor neuropathy **after** her first ... Upon transfer of care to our