

3,920 Results Any time

[\(PDF\) Hypercoagulability detected by thromboelastography ...](#)

<https://www.researchgate.net/publication/335813583...>

PDF | On Aug 1, 2019, Lui G , Bellomo R, McNicol L, Weinberg L, published Hypercoagulability detected by thromboelastography in patients undergoing elective total hip arthroplasty...

[Hypercoagulability detected by thromboelastography in ...](#)

<https://www.researchgate.net/publication/340964969...>

Hypercoagulability detected by thromboelastography in patients undergoing elective total hip arthroplasty April 2019 Conference: Australian and New Zealand College of ...

[Changes of thrombelastography in patients undergoing ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4100560>

In conclusion, our results indicate that thrombelastography was an effective way to identify hypo- and hypercoagulability in patients undergoing elective primary total knee and total hip replacement...

See more

229,000 Results Any time ▾

Changes of thromboelastography in patients undergoing ...
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4100560>

Jul 08, 2014 - Thromboelastography was an effective way to identify hypercoagulability in patients undergoing elective primary total knee and total hip replacement. Platelet may play an important role in the progress of blood hypercoagulability. Keywords: Thromboprophylaxis, Low molecular weight heparin, Arthroplasty, Thromboelastography, Hypercoagulability
Cited by: 11 Author: Yi Yang, Zhenjun Yao, Wenda Dai, Peng ...
Publish Year: 2014

Changes of thromboelastography in patients undergoing ...
<https://jor-online.biomedcentral.com/articles/10.1186/s13018-014-0052-0>

Jul 08, 2014 - Thromboelastography was an effective way to identify hypercoagulability in patients undergoing elective primary total knee and total hip replacement. Platelet may play an important role in the progress of blood hypercoagulability.
Cited by: 11 Author: Yi Yang, Zhenjun Yao, Wenda Dai, Peng ...
Publish Year: 2014 Estimated Reading Time: 8 mins

The application of thromboelastography in risk ...
<https://pubmed.ncbi.nlm.nih.gov/32787371>

The application of thromboelastography in risk stratification for selective thromboembolism prophylaxis after total joint arthroplasty in Chinese: a randomized controlled trial Ann Palliat Med. 2020 Sep;9(5):2498-2507. doi: 10.21037/apm-19-385
Author: Ziming Chen, Yuanchen Ma, Qingtian ... Publish Year: 2020

Thromboelastography

Thromboelastography (TEG) is a method of testing the efficiency of blood coagulation. It is a test mainly used in surgery and anesthesiology, although increasingly used in resuscitations in Emergency Departments, intensive care units, and labor and delivery suites. More common tests of blood coagulation include prothrombin time (PT) and partial thromboplastin time (aPTT) which measure coagulation factor function, but TEG also can assess platelet function, clot strength, and fibrinolysis which these other tests cannot.

 [Wikipedia](#)

Data from [Wikipedia](#)

[Suggest an edit](#)

Search Tools

[Turn off Hover Translation \(关闭翻译\)](#)



Name of Journal: *World Journal of Orthopedics*

Manuscript NO: 64320

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Thromboelastography in elective total hip arthroplasty

THA causes hypercoagulability demonstrated on TEG

Patryck Lloyd-Donald, Wen-Shen Lee, Guoming Liu, Rinaldo Bellomo, Larry McNicol, Laurence Weinberg

Match Overview

1	Crossref 19 words L. Weinberg, M. Scurnah, F. C. Parker, R. Daur, J. Marshall, P. Mccall, D. Story, C. Smith, L. McNicol. 'Markers of Coagulati	1%
2	Internet 16 words crawled on 08-Feb-2018 www.cmc.ca/infoc.org	1%
3	Internet 14 words crawled on 05-Sep-2016 www.science.gov	<1%
4	Crossref 13 words Gonzalez, E., "Differentiation of Enzymatic from Platelet Hy... coagulability Using the Novel Thromboelastography Parameter	<1%

国内版

国际版

Thromboelastography in elective total hip arthroplasty



ALL

IMAGES

VIDEOS

237,000 Results

Any time ▾

Changes of thrombelastography in patients undergoing ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4100560>

Jul 08, 2014 · Thrombelastography was an effective way to identify hypercoagulability in patients undergoing elective primary total knee and total hip replacement. Platelet may play an important role in the progress of blood hypercoagulability. Keywords: Thromboprophylaxis, Low molecular weight heparin, Arthroplasty, Thrombelastography, Hypercoagulability

Cited by: 11

Author: Yi Yang, Zhenjun Yao, Wenda Dai, Peng ...

Publish Year: 2014

The application of thromboelastography in risk ...

<https://pubmed.ncbi.nlm.nih.gov/32787371>

The application of thromboelastography in risk stratification for selective thromboembolism prophylaxis after total joint arthroplasty in Chinese: a randomized controlled trial *Ann Palliat Med* . 2020 Sep;9(5):2498-2507. doi: 10.21037/apm-19-385.

Author: Ziming Chen, Yuanchen Ma, Qingtian ... Publish Year: 2020

(PDF) Hypercoagulability detected by thromboelastography ...

<https://www.researchgate.net/publication/335813583...>

PDF | On Aug 1, 2019, Patryck Lloyd-Donald and others published Hypercoagulability detected by thromboelastography in patients undergoing elective total hip arthroplasty. | Find, read and cite all ...

Changes of thrombelastography in patients undergoing ...

<https://jor-online.biomedcentral.com/articles/10.1186/s13018-014-0052-0> ▾

Jul 08, 2014 · Thrombelastography was an effective way to identify hypercoagulability in patients undergoing elective primary total knee and total hip replacement. Platelet may play an important role in the progress of blood hypercoagulability.

Cited by: 11

Author: Yi Yang, Zhenjun Yao, Wenda Dai, Peng ...

Publish Year: 2014 Estimated Reading Time: 8 mins

Application of thrombelastography (TEG) for safety ...

<https://jor-online.biomedcentral.com/articles/10.1186/s13018-019-1250-6> ▾

Jul 15, 2019. Microclastic hemostatic assays (MHA) technology such as thrombelastography (TEG)