

87639_Auto_Edited.docx

Observational Study**Association between serum estradiol level and appendicular lean mass index in middle-aged postmenopausal women**

Fang Jin, Yanfei Wang, Zhongxin Zhu

Abstract**BACKGROUND**

Previous studies investigating the association between loss of estrogen at menopause and skeletal muscle mass came to contradictory conclusions.

AIM

To evaluate the association between serum estradiol level and appendicular lean mass index in middle-aged postmenopausal women using population-based data.

METHODS

This study included 673 postmenopausal women, aged 40–59 years, from the National Health and Nutrition Examination Survey between 2013 and 2016. Weighted multivariable linear regression models were used to evaluate the association between serum E2 Level and ALMI. When non-linear associations were found by using weighted generalized additive model and smooth curve fitting, two-piecewise linear regression models were further applied to examine the threshold effects.

RESULTS

There was a positive association between serum E2 Level and ALMI. Compared to individuals in quartile 1 group, those in other quartiles had higher ALMI levels. An inverted U-shaped curve relationship between serum E2 Level and ALMI was found on performing weighted generalized additive model and smooth curve fitting, and the inflection point was identified as a serum E2 Level of 85 pg/mL.

CONCLUSION

Our results demonstrated an inverted U-shaped curve relationship between serum E2 Levels and ALMI in middle-aged postmenopausal women, suggesting that low serum E2 Levels play an important in the loss of muscle mass in middle-aged postmenopausal women.

INTRODUCTION

Most women experience menopausal transition in middle age, when aging-related hormonal changes accelerate^[1]. The onset of sarcopenia, a multifactorial condition related to the loss of muscle mass and quality, has been intimately linked to menopause^[2, 3].

Compared with the anabolic effects of androgens on the skeletal muscle mass in men^[4, 5], the effects of estrogens on the skeletal muscle mass in women are less clearly understood^[6]. Moreover, previous studies on the association between the loss of estrogen at menopause and skeletal muscle mass or function came to contradictory conclusions^[7]. As the most potent estrogen hormone, estradiol (E2) is responsible for the maintenance of sexual characteristics and muscle health^[8]. Thus, we aimed to evaluate the association between serum E2 Level and appendicular lean mass index (ALMI) in middle-aged postmenopausal women using population-based data.

MATERIALS AND METHODS

Data source and study population

The National Health and Nutrition Examination Survey (NHANES) is a large, ongoing cross-sectional survey conducted annually in a nationally representative sample of the non-institutionalized United States population. Data for this study were pooled from the NHANES between 2013 and 2016. The study population was restricted to postmenopausal women aged 40–59 years. Individuals with a regular period in the past 12 months ($n = 840$), or with an unrecorded menopausal status ($n = 287$), as well as those with missing serum E2 Levels ($n = 69$) or ALMI data ($n = 171$) were excluded. Finally, 673 women were included in the analysis.

Written informed consent was obtained from all participants and the Institutional Review Board of the National Center for Health Statistics (NCHS) approved the survey protocols (Protocol #2011-17).

Study variables

The exposure variable was the serum E2 Level, which was measured based on the reference method of the National Institute for Standards and Technology, using isotope dilution liquid chromatography tandem mass spectrometry. The outcome variable was ALMI, which was measured by dual-energy X-ray absorptiometry whole-body scans and calculated as the appendicular lean mass (kg) divided by height squared (m^2). The covariates included in this study were age, race, educational level, body mass index (BMI), ratio of family income to poverty, moderate activities, total protein, blood urea nitrogen, and serum uric acid and calcium levels. Detailed information on these variables can be found on the NHANES website (<https://www.cdc.gov/nchs/nhanes/>).

Statistical analyses

All estimates were applied with weights, in accordance with the guidelines edited by the NCHS^[9], to account for the NHANES sampling method. All analyses were performed using EmpowerStats software (<http://www.empowerstats.com>) and R software (version 3.4.3). The statistical significance was set at $P < 0.05$. Weighted multivariable linear regression models were used to evaluate the association between serum E2 Level and ALMI. Following the Strengthening the Reporting of Observational

Studies in Epidemiology statement^[10], we constructed three models: Model 1, no covariates were adjusted; Model 2, age and race were adjusted; and Model 3, all covariates presented in Table 1 were adjusted. When non-linear associations were found by using weighted generalized additive model and smooth curve fitting, two-piecewise linear regression models were further applied to examine the threshold effects.

RESULTS

Demographic characteristics of the participants subclassified based on the serum E2 Level quartiles (Q1: ≤ 3.80 pg/mL; Q2: 3.88–7.42 pg/mL; Q3: 7.45–17.50 pg/mL; and Q4: ≥ 17.60 pg/mL) are shown in Table 1. Compared with the Q1 group, individuals in other groups were younger, and had lower levels of blood urea nitrogen, and higher levels of income to poverty ratio, BMI, total protein, serum uric acid, and ALMI.

The association between serum E2 Level and ALMI was positive in each model, with a significant P for trend among the different serum E2 Level quartile groups (Table 2). In the subgroup analysis stratified by BMI and race, this positive association was significant in the group with BMI < 25 kg/m² (Table 3).

An inverted U-shaped curve relationship between serum E2 Level and ALMI was found, as shown in Figure 1, and the inflection point was identified at a serum E2 Level of 85 pg/mL (Table 4).

DISCUSSION

This study evaluated the association between serum E2 Level and ALMI in middle-aged postmenopausal women, and found an inverted U-shaped curve relationship between them, with the point of inflection at a serum E2 Level of 85 pg/mL.

Estrogens, especially E2, are known to play an important role in the preservation of muscle health. Several studies have investigated the effects of hormone replacement therapy (HRT) and found that it has a positive and measurable impact on muscle function^[11, 12]. Conversely, other studies found that HRT does not protect against

muscle loss^[13, 14]. Moreover, it was reported that menopausal HRT was associated with an increased risk of adverse events, such as dementia^[15], stroke^[16], and breast cancer^[17]. Therefore, it is important to balance the potential benefits against risks. Our results revealed an inverted U-shaped curve relationship between serum E2 Level and ALMI, suggesting that adequate E2 supplementation may be a useful adjunct therapy for individuals with a low serum E2 Level.

The exact mechanism underlying the effects of E2 on skeletal muscle remains unclear. A possible explanation for the potentially beneficial effect is that E2 can stimulate the proliferative activity of the muscle satellite cells (stem cells) that are responsible for muscle tissue maintenance^[18, 19]. Another possible explanation is that estrogen deficiency results in the loss of muscle mass through apoptotic mechanisms^[20, 21]. Despite these possibilities, the molecular mechanism of the impact of E2 on muscle function needs to be further explored.

Data from the NHANES surveys were acquired following standard protocols, which ensured that the data were accurate and consistent. However, the limitations of this study should also be noted. First, a causal relationship between serum E2 Level and ALMI in middle-aged postmenopausal women could not be determined due to the cross-sectional design of the NHANES surveys. Second, biases caused by unmeasured confounding factors cannot be excluded. Third, the conclusion cannot be generalized to older women because the population of this study was restricted to middle-aged postmenopausal women.

CONCLUSION

Overall, this study showed an inverted U-shaped curve relationship between serum E2 Levels and ALMI in middle-aged postmenopausal women, suggesting that low serum E2 Levels play a crucial role in the loss of muscle mass in middle-aged postmenopausal women.

ARTICLE HIGHLIGHTS

Research background

The onset of sarcopenia, a multifactorial condition related to the loss of muscle mass and quality, has been intimately linked to menopause.

Research motivation

Compared with the anabolic effects of androgens on the skeletal muscle mass in men, the effects of estrogens on the skeletal muscle mass in women are less clearly understood. Moreover, previous studies on the association between the loss of estrogen at menopause and skeletal muscle mass or function came to contradictory conclusions.

Research objectives

We aimed to evaluate the association between serum E2 Level and appendicular lean mass index (ALMI) in middle-aged postmenopausal women using population-based data.

Research methods

This study included 673 postmenopausal women, aged 40–59 years, from the National Health and Nutrition Examination Survey between 2013 and 2016. Weighted multivariable linear regression models were used and when non-linear associations were found by using weighted generalized additive model and smooth curve fitting, two-piecewise linear regression models were further applied to examine the threshold effects.

Research results

There was a positive association between serum E2 Level and ALMI. Compared to individuals in quartile 1 group, those in other quartiles had higher ALMI levels. An inverted U-shaped curve relationship between serum E2 Level and ALMI was found on performing weighted generalized additive model and smooth curve fitting, and the inflection point was identified as a serum E2 Level of 85 pg/mL.

Research conclusions

Our results demonstrated an inverted U-shaped curve relationship between serum E2 Levels and ALMI in middle-aged postmenopausal women, suggesting that low serum E2 Levels play an important in the loss of muscle mass in middle-aged postmenopausal women.

Research perspectives

The molecular mechanism of the impact of E2 on muscle function needs to be further explored

6%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|----------|--|----------------------|
| 1 | www.ncbi.nlm.nih.gov <small>Internet</small> | 25 words — 2% |
| <hr/> | | |
| 2 | f6publishing.blob.core.windows.net <small>Internet</small> | 18 words — 1% |
| <hr/> | | |
| 3 | Zhongxin Zhu, Hongliang Zhou, Yanfei Wang, Xiacong Yao. "Associations between bone turnover markers and bone mineral density in older adults", Journal of Orthopaedic Surgery, 2021 <small>Crossref</small> | 14 words — 1% |
| <hr/> | | |
| 4 | ec.bioscientifica.com <small>Internet</small> | 14 words — 1% |
| <hr/> | | |
| 5 | Jiazhong Ji, Yue Hou, Zhaoyang Li, Ying Zhou, Huaming Xue, Tao Wen, Tao Yang, Long Xue, Yihui Tu, Tong Ma. "Association between physical activity and bone mineral density in postmenopausal women: a cross-sectional study from the NHANES 2007–2018", Journal of Orthopaedic Surgery and Research, 2023 <small>Crossref</small> | 13 words — 1% |
| <hr/> | | |
| 6 | www.hindawi.com <small>Internet</small> | 12 words — 1% |

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES OFF
EXCLUDE MATCHES < 12 WORDS