



1

Name of Journal: World Journal of Respiriology

ESPS Manuscript NO: 22849

Manuscript Type: Review


Lung Cancer Screening: Should We Be Excluding People with Previous Malignancy?

Cherie P. Erkmen,

Ashley L. Ehret,

Larry H. Kaiser

Match Overview



1	Internet 120 words crawled on 19-Dec-2015 office.wjgnet.com	1%
2	Internet 44 words crawled on 09-Feb-2014 effiloop.com	<1%
3	Internet 35 words crawled on 12-Apr-2008 intl-annonc.oxfordjournals.org	<1%
4	Internet 28 words crawled on 01-Sep-2013 67.225.242.239	<1%
5	Internet 26 words crawled on 06-Sep-2015 jkma.kma.org	<1%
6	Internet 25 words crawled on 20-Apr-2014 clinicaltrials.gov	<1%



学术搜索

找到约 21,300 条结果 (用时0.12秒)

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☒ 包含引用

[HTML] ... Association for Thoracic Surgery guidelines for lung cancer screening using low-dose computed tomography scans for lung cancer survivors and other high-

MT Jaklitsch, FL Jacobson, JHM Austin, JK Field... - The Journal of thoracic ... 2012 - Elsevier

... diagnosis of pneumonia, occupational exposure to asbestos, prior diagnosis of malignant tumor, and family ... disconcerting to patients, particularly when family history includes multiple cancers and multiple ... and availability of LDCT screening for the early detection of lung cancer. ...

被引用次数: 241 相关文章 所有 127 个版本 引用

sciencedirect.com

Single flexible sigmoidoscopy screening to prevent colorectal cancer: baseline findings of a UK multicentre randomised trial

UK Flexible Sigmoidoscopy Screening Trial ... - The Lancet, 2002 - Elsevier

... Both cancers were classified as Dukes' A (counted only once in table ... Six people died within 30 days of flexible sigmoidoscopy screening. ... days, 15 days, 26 days) and one case each of cardiomyopathy (the next day), intracerebral haemorrhage (7 days), and lung cancer (18 days ...

被引用次数: 419 相关文章 所有 5 个版本 引用

Epidermal growth factor receptor mutation testing in the care of lung cancer patients

LV Sequist, VA Joshi, PA Jänne, DW Bell, P Fidias... - Clinical Cancer ... 2006 - AACR

... We have identified one patient in whom a novel variant was present in both the tumor and the normal tissue ... The relationship between epidermal growth factor receptor mutations and clinicopathologic features in non-small cell lung cancers. Clin Cancer Res 2005;11:1167-73. ...

被引用次数: 79 相关文章 所有 5 个版本 引用

aacrjournals.org

Global cancer statistics in the year 2000

DM Parkin - The lancet oncology, 2001 - Elsevier

... had breast cancer diagnosed within the past 5 years (compared with just 1.4 million survivors - men or women - from lung cancer). ... survival), as well as earlier diagnosis, due to both screening programmes and to better awareness of breast cancer and the ... 29. Colorectal cancers. ...

被引用次数: 2600 相关文章 所有 10 个版本 引用

thelancetnorway.c

[全部](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 1,230,000 条结果 (用时 0.76 秒)

Lung malignancy | Patient

[patient.info](#) > [Education](#) > [Lung malignancy](#) ▾ [翻译此页](#)

Approximately 95% of all primary **lung tumours** are bronchial carcinomas. Metastases ...
People with a **previous** history of **cancer** (especially head and neck). Industrial ... Often excised to **exclude malignancy**. ... Chest CT scan **should** be performed before an intended fibre-optic bronchoscopy or any other biopsy procedure.

Evaluation of Individuals With Pulmonary Nodules: When Is ...

[www.ncbi.nlm.nih.gov](#) > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: MK Gould - 2013 - 被引用次数: 207 - 相关文章

Diagnosis and Management of **Lung Cancer**, 3rd ed: American College of Chest Physicians ... When the clinical probability of **malignancy** is very low (< 5%) **We exclude** from consideration individuals with diffuse or multiple nodules, ... all **previous** CT scans **should** be reviewed when a nodule is first identified by CT scan.

CT Scan Screening for Lung Cancer: Risk Factors for ... - NCBI

[www.ncbi.nlm.nih.gov](#) > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: AK Greenberg - 2012 - 被引用次数: 27 - 相关文章

2012年7月2日 - **We** diagnosed 30 **lung cancers**, 26 stage I. Three patients had ... Over 94 million current and **former** smokers are at increased risk for **lung cancer**, and might benefit ... CT **screening** protocols—both to refine who **should** be **screened** and to **We** chose to **exclude** individuals with sub-solid nodules over 4 mm ...

Lung cancer risks and causes | Cancer Research UK

[www.cancerresearchuk.org](#) > ... > [About lung cancer](#) ▾ [翻译此页](#)

2014年5月23日 - It is one of the few **cancers** where there is a clear cause, with the ... **We** know that the risk of **lung cancer** for passive smokers goes up the ... Overall, **people**