

## Match Overview

1	<b>Internet</b> 96 words crawled on 26-Oct-2014 <a href="http://psap.org">psap.org</a>	2%
2	<b>Crossref</b> 20 words <a href="#">Pediatric Sedation Outside of the Operating Room, 2015.</a>	1%
3	<b>Crossref</b> 16 words <a href="#">Johnson, Peter N., Krystal A. Boyles, and Jamie L. Miller. "Selection of the Initial Methadone Regimen for the M</a>	<1%
4	<b>Crossref</b> 14 words <a href="#">Gitte Y. Larsen. "Preventable harm occurring to critically ill children", Pediatric Critical Care Medicine, 07/2007</a>	<1%
5	<b>Crossref</b> 9 words <a href="#">Alyssa J. Rake. "Glucose variability and survival in critically ill children: Allostasis or harm?"</a>	<1%
6	<b>Internet</b> 9 words crawled on 27-Dec-2016 <a href="http://office.wjgnet.com">office.wjgnet.com</a>	<1%

6

Name of Journal: *World Journal of Clinical Pediatrics*

ESPS Manuscript NO: 32422

Manuscript Type: Original Article

*Retrospective Study*

Conversion from Prolonged Intravenous Fentanyl Infusion to Enteral Methadone in Critically Ill Children

Vijay Srinivasan, Daniel Pung, Sean P O'Neill

**Abstract**

**Aim:** To describe our institutional experience with conversion from intravenous (IV) fentanyl infusion directly to enteral methadone and occurrence of withdrawal in critically ill mechanically ventilated children exposed to prolonged sedation and analgesia.

**Methods:** With Institutional Review Board approval, we retrospectively studied



Conversion from Prolonged Intravenous Fentanyl Infusion to Enteral Meth



全部

新闻

图片

视频

更多

设置

工具

找到约 9,540 条结果 (用时 0.67 秒)

### Use of Methadone for Prevention of Opioid Withdrawal in Critically Ill ...

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: SA Jeffries - 2012 - 被引用次数: 9 - 相关文章

Opioids are commonly administered to **critically ill children** for analgesia and sedation, ... The **conversion** ratio for **IV** morphine to oral **methadone** was 1:0.78 for ..... most of the literature involves **conversion from long-term IV fentanyl infusion**, .... Efficacy of an **enteral** 10-day **methadone** wean to prevent opioid withdrawal in ...

### Use of Methadone for Opioid Weaning in Children: Prescribing ... - NCBI

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: K Giby - 2014 - 被引用次数: 1 - 相关文章

**Critically ill children** are routinely treated with opioids to relieve pain, produce ... However, **prolonged** exposure to opioids followed by abrupt cessation frequently ... **Fentanyl** and hydromorphone doses were **converted** to equivalents of **IV** ..... the benefits of **enteral methadone** for weaning from mechanical ventilation in the ...

### Efficacy of an enteral 10-day methadone wean to prevent opioid ...

[https://www.researchgate.net/.../10719143\\_Efficacy\\_of\\_an\\_enteral\\_10-day\\_...](https://www.researchgate.net/.../10719143_Efficacy_of_an_enteral_10-day_...) - 翻译此页

All 29 patients had received a continuous **fentanyl infusion**; duration of exposure was 14.5 +/- 9.2 days, .... opioid withdrawal symptoms in pediatric patients after **prolonged** ..... **Conversion** of **intravenous fentanyl** to **enteral methadone** ..... rence of withdrawal in **critically ill sedated children**. Crit Care Med 1999; 27:196-199





Conversion from Prolonged Intravenous Fentanyl Infusion to Enteral Meth:



全部

新闻

图片

购物

视频

更多

设置

工具

找到约 15,500 条结果 (用时 0.72 秒)

## Google 学术: Conversion from Prolonged Intravenous Fentanyl Infusion to Enteral Methadone in Critically Ill Children

Efficacy of an enteral 10-day methadone wean to ... - Meyer - 被引用次数: 48

Enteral methadone to expedite fentanyl discontinuation ... - Lugo - 被引用次数: 50

Withdrawal symptoms in critically ill children after long- ... - Ista - 被引用次数: 101

## Use of Methadone for Prevention of Opioid Withdrawal in Critically Ill ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: SA Jeffries - 2012 - 被引用次数: 11 - 相关文章

Opioids are commonly administered to critically ill children for analgesia and sedation, ... The conversion ratio for IV morphine to oral methadone was 1:0.78 for .... most of the literature involves conversion from long-term IV fentanyl infusion, .... Efficacy of an enteral 10-day methadone wean to prevent opioid withdrawal in ...

## Use of Methadone for Opioid Weaning in Children: Prescribing ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: K Giby - 2014 - 被引用次数: 3 - 相关文章

Critically ill children are routinely treated with opioids to relieve pain, produce ... Fentanyl and hydromorphone doses were converted to equivalents of IV .... In most standardized protocols for long-term opioid users, ... dose reduction has ... the benefits of enteral methadone for weaning from mechanical ventilation in the ....

全部

新闻

图片

购物

视频

更多

设置

工具

找到约 17,600 条结果 (用时 0.79 秒)

## Google 学术: Conversion from prolonged intravenous fentanyl infusion to enteral methadone in critically ill children

Efficacy of an enteral 10-day methadone wean to ... - Meyer - 被引用次数: 49

Enteral methadone to expedite fentanyl discontinuation ... - Lugo - 被引用次数: 50

Withdrawal symptoms in critically ill children after long- ... - Ista - 被引用次数: 102

## Use of Methadone for Prevention of Opioid Withdrawal in Critically Ill ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: SA Jeffries - 2012 - 被引用次数: 11 - 相关文章

Opioids are commonly administered to critically ill children for analgesia and sedation, ... The conversion ratio for IV morphine to oral methadone was 1:0.78 for .... most of the literature involves conversion from long-term IV fentanyl infusion, .... Efficacy of an enteral 10-day methadone wean to prevent opioid withdrawal in ...

## Use of Methadone for Opioid Weaning in Children: Prescribing ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: K Giby - 2014 - 被引用次数: 3 - 相关文章

Critically ill children are routinely treated with opioids to relieve pain, produce ... Fentanyl and hydromorphone doses were converted to equivalents of IV .... In most standardized protocols for long-term opioid users, dose reduction has .... the benefits of enteral methadone for weaning from mechanical ventilation in the ...

## Tolerance and Withdrawal From Prolonged Opioid Use in Critically Ill ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: KJS Anand - 2010 - 被引用次数: 142 - 相关文章