

Name of Journal: *World Journal of Clinical Urology*

ESPS Manuscript NO: 19762

Manuscript Type: REVIEW

Is continuous bladder irrigation after prostate surgery still needed?

Continuous bladder irrigation in contemporary practice

Chukwudi Ogonnaya Okorie

Abstract

Continuous bladder irrigation (CBI) is commonly prescribed after certain prostate surgeries to help prevent the clot formation and retention that are frequently associated

Match Overview

1	CrossCheck 126 words Okorie, C.O.. "Modified Suprapubic Prostatectomy Without Irrigation Is Safe", <i>Urology</i> , 201003	4%
2	CrossCheck 47 words Lezrek, M.. "Modified Denis technique: a simple solution for maximal hemostasis in suprapubic prostatectomy", <i>Urolog</i>	1%
3	CrossCheck 27 words L. F. Tinckler. "POST-PROSTATECTOMY MANAGEMENT-... NEW CATHETER SYSTEM", <i>British Journal of Urology</i> , 04/1	1%
4	CrossCheck 16 words Melcher, Barbara, John Schroeder, and Steve Shumaker. ... "Cured-in-Place Piping: The Question of External Point Rep	<1%
5	Internet 13 words crawled on 10-Jan-2011 lib.bioinfo.pl	<1%



小提示: 只搜索中文(简体)结果, 可在 [学术搜索设置](#) 指定搜索语言

Gyrus bipolar versus standard monopolar transurethral resection of the prostate: a randomized prospective trial

M de Sio, R Autorino, G Quarto, R Damiano, S Perdonà... - Urology, 2006 - Elsevier

... 4 and 5 In brief, it consists of a generator and a cutting loop that does not differ in ... The resection time and blood loss were similar between the two groups, but the need for continuous bladder irrigation after surgery, time to ... Is transurethral resection of the prostate still justified? ...

被引用次数: 158 [相关文章](#) [所有 6 个版本](#) [引用](#) [保存](#)

Laser ablation of the prostate in patients with benign prostatic hypertrophy

AJ Costello, WG Bowsher, DM Bolton... - British journal of ..., 1992 - Wiley Online Library

... Neodymium: YAG laser ablation of papillary transitional cell carcinoma of the bladder, but there does not appear to ... Regional or general anaesthesia is required and continuous bladder irrigation is often necessary post-operatively to evacuate clot ... Further evaluation is required. ...

被引用次数: 351 [相关文章](#) [所有 5 个版本](#) [引用](#) [保存](#)

Holmium laser enucleation of the prostate (HoLEP): the endourologic alternative to open prostatectomy

EA Elzayat, MM Elhilali - European urology, 2006 - Elsevier

... of a large prostate in much the same way as the surgeon's finger does during open ... Continuous bladder irrigation was necessary in 8 patients (3.5%) for transient postoperative hematuria. ... One of the advantages of HoLEP is that it has no size limitation, with immediate and ...

被引用次数: 155 [相关文章](#) [所有 7 个版本](#) [引用](#) [保存](#)

Complications of transurethral resection of the prostate (TURP)—incidence, management, and prevention

J Rassweiler, D Teber, R Kuntz, R Hofmann - European urology, 2006 - Elsevier

... and (cloudy and opaque) whereas urinary bleeding results in a dark red continuous irrigation fluid

[researchgate.net 中的 \[PDF\]](#)

[muacme.org 中的 \[PDF\]](#)



Is continuous bladder irrigation after prostate surgery still needed?



网页 视频 新闻 图片 更多 ▾ 搜索工具

找到约 104,000 条结果 (用时 0.54 秒)

BPH Surgery, Complications of Surgical Treatment for BPH ...

www.healthcommunities.com/bph...prostate/bph-surgery.shtml ▾ 翻译此页

1998年6月9日 - Surgical treatment for benign prostatic hyperplasia, BPH, involves ... water to flush the bladder;; eat a balanced diet;; use a laxative if necessary to ... Blood in the urine (hematuria) is common after TURP surgery and ... Urine output and color and continuous bladder irrigation (CBI), if present, are monitored.

What a Nurse Should Know About Continuous Bladder ...

educationcareerarticles.com/.../what-a-nurse-should-know-abou... ▾ 翻译此页

2013年9月9日 - Continuous bladder irrigation (CBI) is typically used following surgery, ... the catheter may need to be drained and irrigated every 20 minutes to ... of clot retention that may pop up after transurethral prostatectomy surgery, There's still time to apply to Grand Canyon University and pursue your education.

Management of clot retention following urological surgery ...

www.nursingtimes.net/management...surgery/206307.article ▾ 翻译此页

Improving hand hygiene after neurological injury to prevent clot retention by continuously irrigating the bladder via a three-way catheter (Gilbert and Gobbi, 1989). Bladder irrigation is required due to the vascular nature of the prostate, and to treatment for benign prostatic hyperplasia and may still suffer severe adverse ...

Instructions for Care Following Transurethral Resection of ...

www.med.umich.edu/1libr/urology/postcare/turp.htm ▾ 翻译此页

Instructions for Care Following Transurethral Resection of the Prostate ... Ask your nurse for pain medication, as you need them. ... You may have a continuous bladder irrigation (CBI) through the catheter to help keep the urine clear ... Ambulation: Purpose: Walking soon after surgery helps with early return of normal bowel