
Microsoft Bing	Trends of Central line-associated Bloodstream Infections in the Inte		Q	C	2	
	ALL IMAGES VIDEOS					
	12,000 Results Any time •					
	Trends in central-line-associated bloodstream infections			Search Tools		
	https://www.cambridge.org/core/journals/infection   Trends in central-line associated bloodstream infections and catheter-associated urinary tract infections in a large acute-care hospital system in New York City, 2016-2019 Published online by Cambridge				Turn off Hover Translation (美代取词)	
	Length of Stay in Patients With Central Line-Associated https://www.cureus.com/articles/42347-length-of + 0ct 06, 2020 - The rising use of central lines in routine clinical practice has made central line-associated bloodstream infections (CLASS) a leading cause of preventable healthcare infections. A 2017 report.					
	Author: Naif H Alotaihi Abdulrahman Barri Mu Publish Year 2020					



Impact of a Multidimensional Infection Control Approach on ... https://pubmed.ncbi.nlm.nih.gov/23641950
Background: Central line-associated bloodstream infections (CLABs) have long been associated with excess lengths of stay, increased hospital costs and mortality attributable to them. Different studies fro... Cited by: 68 Author: Hakan Leblebicioglu, Recep Öztürk, Victor ...
Publish Year: 2013

Six-year multicenter study on short-term peripheral venous ... https://pubmed.ncbi.nlm.nih.gov/32295756
Six-year multicenter study on short-term peripheral venous catheters-related bloodstream infection rates in 246 intensive units of 83 hospitals in 52 cities of 14 countries of Middle East: Bahrain, Egypt, Iran,...
Citted by: 3
Author: Victor D. Rosenthal, Souad Belkebir, Farid Z...
Publish Year: 2020

(PDF) Impact of a multidimensional infection control ... https://www.researchgate.net/publication/236636060...



## Trends in central-line-associated bloodstream infections ...

https://www.cambridge.org/core/journals/infection... •

Similarly, average age of the patients with CAUTIs remained constant over the 4 years: 62.8 ± 17.3 years in 2016,  $66.2 \pm 15.3$  years in 2017,  $65.4 \pm 15.6$  in 2018, and  $66.1 \pm 15.9$  in 2019. The number of infections per 1,000 device days progressively declined across the 11-hospital system (Table 1 ).

#### Reducing central line-associated bloodstream infections in ...

https://www.myamericannurse.com/strategies... •

First Intervention When we started the project in 2006, we had three aims—compliance with appropriate hand hygiene, a central-line insertion bundle, and a central-line maintenance bundle. Today, these are still our goals.

The Setback—And A Key Realizati...


See more on myamericannurse.com

Estimated Reading Time: 6 mins



Feedback

The Recovery



### Trends in central-line-associated bloodstream infections ...

https://www.cambridge.org/core/journals/infection... •


Similarly, average age of the patients with CAUTIs remained constant over the 4 years:  $62.8 \pm 17.3$  years in 2016,  $66.2 \pm 15.3$  years in 2017,  $65.4 \pm 15.6$  in 2018, and  $66.1 \pm 15.9$  in 2019. The number of infections per 1,000 device days progressively declined across the 11-hospital system (Table 1).

# Six-year multicenter study on short-term peripheral venous ... https://pubmed.ncbi.nlm.nih.gov/32295756

Six-year multicenter study on short-term peripheral venous catheters-related **bloodstream** infection rates in 246 **intensive** units of 83 hospitals in 52 cities of 14 countries of Middle East: **Bahrain**, Egypt, Iran, Jordan, **Kingdom** of Saudi Arabia, Kuwait, Lebanon, Morocco, Pakistan, Palestine, Sudan, Tunisia, Turkey, and United Arab Emirates-International Nosocomial Infection Control Consortium ...

Cited by: 4 Author: Víctor D. Rosenthal, Souad Belkebir, Fari...

Publish Year: 2020



Feedback

## Eurosurveillance | Prevention of hospital-acquired ...

https://www.eurosurveillance.org/content/10.2807/1560-7917.ES.2016.21.46.30400

We assessed the impact of 2% daily patient bathing with chlorhexidine gluconate (CHG) washcloths on the incidence of hospital-acquired (HA) and central line-associated (CLA) bloodstream infections (BSI) in intensive care units (ICUs). We searched randomised studies in Medline, EMBASE, Cochrane Library (CENTRAL) and Web of Science databases up to April 2015.

Cited by: 32 Author: Elsa Afonso, Koen Blot, Stijn Blot