Respiratory intermediate care units: a European survey ...

https://erj.ersjournals.com/content/20/5/1343 -

Nov 01, 2002 · Among these, 34 units, which admitted 6,803 patients, reported a rate of nosocomial pneumonia of 8%. The reported rate of nosocomial pneumonia in the three levels of care was 10%, 8%, and 5% in the intensive, intermediate, and monitoring units, respectively. Pseudomonas spp. and Staphylococcus aureus accounted for 39% and 29% of cases ...

Cited by: 157 Author: C. Roussos, N. Ambrosino, M. Confalonie...

Publish Year: 2002 Estimated Reading Time: 9 mins

British Thoracic Society guideline for diagnostic flexible ...


https://thorax.bmj.com/content/68/Suppl_1/i1 -

Aug 01, 2013 · All patients undergoing bronchoscopy should have heart rate, respiratory rate, blood pressure and oxygen saturation recorded repeatedly, including before, during and after the procedure. (Grade D) All bronchoscopy units should undertake periodic audit of bronchoscopic performance, including efficacy, complications and patient satisfaction surveys.

Cited by: 572 Author: I A Du Rand, J Blaikley, R Booton, N Chau...

Publish Year: 2013 Estimated Reading Time: 9 mins





Role of bronchoscopy in critically ill patients managed in intermedi







ALL

IMAGES

VIDEOS

56,900 Results

Any time *

Management of antithrombotic agents in patients ...

https://err.ersjournals.com/content/26/145/170001 -

Sep 30, 2017 - Bleeding is one of the most feared complications of flexible bronchoscopy. Although infrequent, it can be catastrophic and result in fatal outcomes. Compared to other endoscopic procedures, the risk of morbidity and mortality from the bleeding is increased, as even a small amount of blood can fill the tracheobronchial tree and lead to respiratory failure. Patients using antithrombotic agents ...

Cited by: 14 Author: Sami Abugayyas, Shine Raju, John R. B...

Publish Year: 2017


British Thoracic Society guideline for diagnostic flexible ...

https://thorax.bmj.com/content/68/Suppl 1/i1 -

Aug 01, 2013 - All patients undergoing bronchoscopy should have heart rate, respiratory rate, blood pressure and oxygen saturation recorded repeatedly, including before, during and after the procedure. (Grade D) All bronchoscopy units should undertake periodic audit of bronchoscopic performance, including efficacy, complications and patient satisfaction surveys.

Cited by: 576 Author: I A Du Rand, J Blaikley, R Booton, N Cha...

Publish Year: 2013 Estimated Reading Time: 9 mins



Feedback

Comparison between ultrasound- and bronchoscopy-guided ...

https://www.researchgate.net/publication/267749459...

Background To determine whether fibre optic bronchoscopy-guided percutaneous dilatational