

## Match Overview

1	<b>Internet</b> 27 words crawled on 30-Sep-2016 <a href="http://icm-experimental.springeropen.com">icm-experimental.springeropen.com</a>	1%
2	<b>Publications</b> 21 words "New Findings on Emergency Medicine from Sarawak General Hospital Summarized (Performance assessment)", Hea	1%
3	<b>Internet</b> 20 words crawled on 30-Mar-2015 <a href="http://jac.oxfordjournals.org">jac.oxfordjournals.org</a>	1%
4	<b>Internet</b> 18 words crawled on 22-Oct-2015 <a href="http://www.malariajournal.com">www.malariajournal.com</a>	1%
5	<b>Internet</b> 17 words crawled on 19-May-2016 <a href="http://journal.publications.chestnet.org">journal.publications.chestnet.org</a>	1%
6	<b>Internet</b> 16 words crawled on 19-May-2016 <a href="http://ccforum.biomedcentral.com">ccforum.biomedcentral.com</a>	1%

**Name of Journal:** *World Journal of Critical Care Medicine*

**ESPS Manuscript NO:** 29568

**Manuscript Type:** Original Article

*Retrospective Study*

**Risk factors for mortality in postoperative peritonitis in critically ill patients**

**Yoann Launey, Benjamin Duteurtre, Raphaëlle Larmet, Nicolas Nesseler, Audrey Tawa, Yannick Mallédant, Philippe Seguin**

### Abstract

**Aim:** To identify the risk factors for mortality in intensive care patients with postoperative peritonitis (POP).

**Methods:** This was a retrospective analysis using a prospective database that includes all patients hospitalized in a surgical intensive care unit for POP from

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 140,000 条结果 (用时 0.61 秒)

## Google 学术: Risk factors for mortality in postoperative peritonitis in critically ill patients.

... factors of mortality due to polymicrobial peritonitis with ... - Dupont - 被引用次数: 104

Candida as a risk factor for mortality in peritonitis - Montravers - 被引用次数: 180

Tertiary peritonitis: clinical features of a complex ... - Nathens - 被引用次数: 227

### Management of Peritonitis in the Critically Ill Patient

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > NCBI > Literature > PubMed Central (PMC) ▾ [翻译此页](#)

作者: CA Ordoñez - 2006 - 被引用次数: 69 - [相关文章](#)

The mortality of an intra-peritoneal infection in the early 1900s was close to 90%. .... The incidence of postoperative peritonitis occurs in 1% to 20% of patients ... to the primary reason for laparotomy and other risk factors that are discussed ...

### Factors associated with septic shock and mortality in generalized ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > NCBI > Literature > PubMed Central (PMC) ▾ [翻译此页](#)

作者: FC Riché - 2009 - 被引用次数: 71 - [相关文章](#)

2009年6月24日 - In secondary peritonitis [2], postoperative peritonitis is commonly .... Risk factors for mortality in patients with septic shock are presented in Table Table5.5. .... with Candida isolation in peritoneal fluid in critically ill patients.

### Predictors of in-hospital mortality in critically ill patients with secondary ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: JB Noguera - 2015 - [相关文章](#)

2015年10月1日 - Critically ill surgical patients remain at high risk of adverse outcomes as a result ... together with those factors associated with the development of tertiary peritonitis. ... between



[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 126,000 条结果 (用时 0.86 秒)

## Google 学术: Risk factors for mortality in postoperative peritonitis in critically ill patients

... factors of mortality due to polymicrobial peritonitis with ... - Dupont - 被引用次数: 109

Candida as a risk factor for mortality in peritonitis - Montravers - 被引用次数: 186

Tertiary peritonitis: clinical features of a complex ... - Nathens - 被引用次数: 227

## Perioperative risk factors for in-hospital mortality after emergency ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: JY Lee - 2016

2016年9月2日 - The findings of this study suggest that in critically ill patients ... Few studies have evaluated risk factors in critically ill surgical patients with peritonitis. ... Postoperative biochemical parameters were measured immediately after ...

## Factors associated with septic shock and mortality in ... - NCBI

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > NCBI > Literature > PubMed Central (PMC) ▾ [翻译此页](#)

作者: FC Riché - 2009 - 被引用次数: 70 - [相关文章](#)

2009年6月24日 - In secondary peritonitis [2], postoperative peritonitis is commonly thought to ... Septic shock was defined according to the criteria of the Critical Care ..., Risk factors for mortality in patients with septic shock are presented in Table Table5.5. .... with Candida isolation in peritoneal fluid in critically ill patients.

## Prognostic factors for mortality in (fecal) peritonitis: back to the roots ...

[link.springer.com/article/10.1007/s00134-013-3155-x](http://link.springer.com/article/10.1007/s00134-013-3155-x) - [翻译此页](#)

作者: C Eckmann - 2014 - 被引用次数: 2 - [相关文章](#)

The most striking prognostic risk factor in (fecal) peritonitis is increased age. ... of patients with

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 128,000 条结果 (用时 0.71 秒)

## Google 学术: Risk factors for mortality in postoperative peritonitis in critically ill patients

... factors of mortality due to polymicrobial peritonitis with ... - Dupont - 被引用次数: 109

Candida as a risk factor for mortality in peritonitis - Montravers - 被引用次数: 186

Tertiary peritonitis: clinical features of a complex ... - Nathens - 被引用次数: 229

### Management of Peritonitis in the Critically Ill Patient

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: CA Ordoñez - 2006 - 被引用次数: 68 - 相关文章

The mortality of an intra-peritoneal infection in the early 1900s was close to 90%. .... The incidence of postoperative peritonitis occurs in 1% to 20% of patients ... to the primary reason for laparotomy and other risk factors that are discussed ...

### Factors associated with septic shock and mortality in generalized ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: FC Riché - 2009 - 被引用次数: 74 - 相关文章

2009年6月24日 - In secondary peritonitis [2], postoperative peritonitis is commonly thought to ... Septic shock was defined according to the criteria of the Critical Care .... Risk factors for mortality in patients with septic shock are presented in Table Table5.5. .... with Candida isolation in peritoneal fluid in critically ill patients.

### Predictors of in-hospital mortality in critically ill patients with secondary ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > NCBI > Literature > PubMed Central (PMC) - 翻译此页

2015年10月1日 - Critically ill surgical patients remain at high risk of adverse outcomes as a result ...