

Name of Journal: *World Journal of Critical Care Medicine*

ESPS Manuscript NO: 29679

Manuscript Type: Original Article

Observational Study

Timing, method, and discontinuation of hydrocortisone administration for septic shock patients

Miguel A Ibarra-Estrada, Quetzalcóatl Chávez-Peña, Claudia I Reynoso-Estrella,
Jorge Rios-Zermeño, Pável E Aguilera-González, Miguel A García-Soto,
Guadalupe Aguirre-Avalos

Abstract

AIM

To characterize the prescribing patterns for hydrocortisone for patients with septic

Match Overview

1	Internet 44 words crawled on 14-Nov-2016 criticalultrasoundjournal.springeropen.com	1%
2	Crossref 23 words Green, Jeffrey P., Jason Adams, Edward A. Panacek, and Timothy A. Albertson. "The 2012 Surviving Sepsis Cam	1%
3	Crossref 15 words Bruno, Jeffrey J., Brian M. Dee, Brent A. Anderegg, Mike Hernandez, and S. Egbert Pravinkumar. "US practitioner	<1%
4	Internet 13 words crawled on 25-Dec-2016 pdfs.semanticscholar.org	<1%
5	Crossref 13 words &NA; "SOCIETY OF CRITICAL CARE MEDICINE 40TH CRITICAL CARE CONGRESS SAN DIEGO, CALIFORNI ...	<1%
6	Crossref 13 words "Sunday, 31 August 2008", <i>European Heart Journal</i> , 09/02/2008	<1%
7	Crossref 12 words	<1%



All News Images Shopping Videos More Search tools

About 221,000 results (0.74 seconds)

Scholarly articles for Timing, Method, and Discontinuation of Hydrocortisone Administration for Septic Shock Patients

... on mortality in **patients** with **septic shock** - Annane - Cited by 3120

... treatment of severe **sepsis** and **septic shock** in adults: a ... - Annane - Cited by 492

... for management of severe **sepsis** and **septic shock** - Dellinger - Cited by 4356

Prescribing patterns of hydrocortisone in septic shock: a single-center ...

www.ncbi.nlm.nih.gov/.../23787398 ▼ National Center for Biotechnology Information ▼
by KM Contrael - 2013 - Cited by 10 - Related articles

DESIGN: Retrospective chart review of **patients** with **septic shock** over 17 ... The **method** and **timing** of corticosteroid **discontinuation** also differed among providers. ... Humans; **Hydrocortisone/administration** & dosage*; Intensive Care Units* ...

Corticosteroids for severe sepsis: an evidence-based guide for ... - NCBI

www.ncbi.nlm.nih.gov/.../PMC32244... ▼ National Center for Biotechnology Information ▼
by D Annane - 2011 - Cited by 50 - Related articles

Apr 13, 2011 - Corticosteroids should be initiated only in **patients** with **sepsis** who require 0.5 within 1 hour after **hydrocortisone administration** and may likely result from line for **time** equal to 120 hours arguing for maintaining treatment at full ... it also may favor adrenal insufficiency after **withdrawal** of corticosteroids.

Early administration of hydrocortisone replacement after the advent of ...

www.ncbi.nlm.nih.gov/.../24674923 ▼ National Center for Biotechnology Information ▼
by CS Katsenos - 2014 - Cited by 17 - Related articles

Early **administration** of **hydrocortisone** replacement after the advent of **septic shock**: ... study in **patients** with **septic shock** treated with low doses of **hydrocortisone**. ... **Time of discontinuation** of vasopressors was earlier among **patients** with ...

[PDF] Withdrawal from glucocorticosteroid therapy - SciELO

www.scielo.br/pdf/jped/v84n3/en_v84n3a03.pdf ▼

找到约 347,000 条结果 (用时 0.67 秒)

Google 学术: Timing, method, and discontinuation of hydrocortisone administration for septic shock patients

... on mortality in patients with septic shock - Annane - 被引用次数: 3178

... treatment of severe sepsis and septic shock in adults: a ... - Annane - 被引用次数: 510

... for management of severe sepsis and septic shock - Dellinger - 被引用次数: 4433

Corticosteroids for severe sepsis: an evidence-based guide for ... - NCBI

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: D Annane - 2011 - 被引用次数: 55 - 相关文章

2011年4月13日 - Corticosteroids should be initiated only in patients with sepsis who require 0.5 ... within 1 hour after hydrocortisone administration and may likely result from ... line for time equal to 120 hours arguing for maintaining treatment at full ... it also may favor adrenal insufficiency after withdrawal of corticosteroids.

Prescribing patterns of hydrocortisone in septic shock: a single-center ...

<https://www.ncbi.nlm.nih.gov/pubmed/23787398> - 翻译此页

作者: KM Contrael - 2013 - 被引用次数: 11 - 相关文章

Prescribing patterns of hydrocortisone in septic shock: a single-center experience ... Sepsis Campaign suggests giving hydrocortisone to septic patients only if their ... The method and timing of corticosteroid discontinuation also differed among providers. ... Academic Medical Centers; Anti-Inflammatory Agents/administration ...

Stress doses of hydrocortisone reverse hyperdynamic septic shock: a ...

<https://www.ncbi.nlm.nih.gov/pubmed/10321661> - 翻译此页

作者: J Briegel - 1999 - 被引用次数: 1002 - 相关文章

Stress doses of hydrocortisone reverse hyperdynamic septic shock: a ... PATIENTS: Forty



全部

新闻

图片

视频

购物

更多

设置

工具

找到约 227,000 条结果 (用时 0.80 秒)

Google 学术: Timing, method, and discontinuation of hydrocortisone administration for septic shock patients

... on mortality in patients with septic shock - Annane - 被引用次数: 3184

... treatment of severe sepsis and septic shock in adults: a ... - Annane - 被引用次数: 510

... for management of severe sepsis and septic shock - Dellinger - 被引用次数: 4442

Corticosteroids for severe sepsis: an evidence-based guide for ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: D Annane - 2011 - 被引用次数: 55 - 相关文章

2011年4月13日 - Corticosteroids should be initiated only in patients with sepsis who require 0.5 ... within 1 hour after hydrocortisone administration and may likely result from ... line for time equal to 120 hours arguing for maintaining treatment at full ... it also may favor adrenal insufficiency after withdrawal of corticosteroids.

Stress doses of hydrocortisone reverse hyperdynamic septic shock: a ...

<https://www.ncbi.nlm.nih.gov/pubmed/10321661> - 翻译此页

作者: J Briegel - 1999 - 被引用次数: 1004 - 相关文章

Stress doses of hydrocortisone reverse hyperdynamic septic shock: a ... PATIENTS: Forty consecutive patients who met the ACCP/SCCM criteria for septic shock. ... The primary study end point was the time to cessation of vasopressor support ... Agents/therapeutic use; Double-Blind Method; Drug Administration Schedule ...

Prescribing patterns of hydrocortisone in septic shock: a single-center ...

<https://www.ncbi.nlm.nih.gov/pubmed/23787398> - 翻译此页