3191 words • 69 matches • 26 sources

FAQ

E- In

20515-Review

Quotes Excluded Bibliography Excluded

25%

iThenticate•

Name of Journal: World Journal of Hypertension

ESPS Manuscript NO: 20515

Manuscript Type: ORIGINAL ARTICLE

Prospective Study

Estimated net endogenous acid production and risk of prevalent and incident

hypertension in community-dwelling older people

Ruth Chan, Jason Leung, Jean Woo

Abstract

AIM: To investigate the associations of dietary acid-base load with prevalent and

CrossCheck 151 words 5% Chan, R., J. Leung, and J. Woo. "Association Between Esti mated Net Endogenous Acid Production and Subsequent CrossCheck 119 words 4% Chan, Ruth, Dicken Chan, and Jean Woo "A Prospective Cohort Study to Examine the Association between Dietar ... CrossCheck 111 words 3% Chan, Ruth, J. Leung, J. Woo, and T. Kwok. "Associations of dietary protein intake on subsequent decline in muscle CrossCheck 63 words 2% M. F. Engberink. "Dietary acid load and risk of hypertens... n: the Rotterdam Study", American Journal of Clinical Nutrit CrossCheck 61 words 2% Chan, Ruth, Jason Leung, and Jean Woo. "Dietary Pattern s and Risk of Frailty in Chinese Community-Dwelling Old ... Publications 50 words 2% Chan, R.S.M. Woo, J. Chan, D.C.C. Cheung, "Estimated n et endogenous acid production and intake of bone healt ...

Match Overview

incident hunartoncian in community living Chinago alder adults in Hang Vang

BIA IN



网页 图片 学术 高级搜索

Ruth Chan - Publications - ResearchGate

www.researchgate.net/profile/Ruth_Chan/publications

Article: Association Between Estimated Net Endogenous Acid Production and ... in 3,122 Chinese community-dwelling older people aged 65 and over in Hong Kong. ... Metabolic syndrome is the most important risk factor of incident NAFLD. ... public health strategies to prevent hypertension, stroke and renal disease.

[PDF] Estimated Dietary Acid Load Is Not Associated with ... - ResearchGate

www.researchgate.net/profile/Xiaoyan_Huang2/publication/272084616_Estimated_diet 17 Dec 2014 ... incidence of hypertension in older men taking into account each ... Hypertension is prevalent in one-third of the general population ... sium, and by the net endogenous acid production (NEAP) method ... community-dwelling cohort of elderly men. Also Their validity in older adults has been assumed (13).

Fracture Risk in Men With Congestive Heart Failure: Discussion www.medscape.com/viewarticle/576685 4

[10-12] Spironolactone is increasingly being used to treat hypertension and ... of spironolactone (.6 months) was inversely associated with total incident fractures. dietary estimates of net endogenous acid production are associated with low ... risk factors be used to predict fractures in community-dwelling older adults?

Fracture Risk in Men With Congestive Heart Failure: Risk Reduction ... www.sciencedirect.com/science/article/pii/S0735109708014174

We identified 167 cases with a single-incident fracture and matched those by ago and



Estimated net endogenous acid production and risk of prevalent and incident





岡页 新闻 图片 视频 更多▼ 搜索工具

找到约7,230条结果(用时0.75秒)

Ruth Chan - Publications - ResearchGate

www.researchgate.net/profile/Ruth_Chan/publications ▼ 翻译此页
No association between other dietary patterns and incident frailty was observed. ...
Article: A Healthier Lifestyle Pattern for Cardiovascular Risk Reduction Is Associated
Estimated net endogenous acid production (NEAP) was calculated using Chinese community-dwelling older people aged 65 and over in Hong Kong.

Estimated Dietary Acid Load Is Not Associated with Blood ... www.researchgate.net/.../272084616_Estimated_Dietary_Acid_...-翻译此页 2015年2月15日 - Previous research on dietary acid load and BP in the community has ... blood pressure or the incidence of hypertension in older men taking ..., acid load (PRAL) and net endogenous acid production (NEAP), ... excluded, PRAL was not a predictor of incident cases (P > 0.30). Although this is a common

Fracture Risk in Men With Congestive Heart Failure: Risk ...

www.sciencedirect.com/science/article/pii/S0735109708014174 - 翻译此页

作者: LD Carbone - 2008 - 被引用次数: 45 - 相关文章

We identified 167 cases with a single-incident fracture and matched these by age and ...

Age; race; height; weight; cigarette use; prevalent fractures of the hip, high dietary estimates of net endogenous acid production are associated with low ... risk factors be used to predict fractures in community-dwelling older adults?



Estimated net endogenous acid production and risk of prevalent and incident



岡页 新闻 图片 视频 更多▼ 搜索工具

找到约 6,860 条结果 (用时 0.59 秒)

Estimated Dietary Acid Load Is Not Associated with Blood ... www.researchgate.net/.../272084616_Estimated_Dietary_Acid_... - 翻译此页 2015年2月15日 - Previous research on dietary acid load and BP in the community has ... blood pressure or the incidence of hypertension in older men taking ... acid load (PRAL) and net endogenous acid production (NEAP), ... excluded, PRAL was not a predictor of incident cases (P > 0.30), ..., Although this is a common

Ruth Chan - Publications - ResearchGate

www.researchgate.net/profile/Ruth_Chan/publications ▼ 翻译此页
No association between other dietary patterns and incident frailty was observed. ...
Article: A Healthier Lifestyle Pattern for Cardiovascular Risk Reduction Is Associated
Estimated net endogenous acid production (NEAP) was calculated using Chinese community-dwelling older people aged 65 and over in Hong Kong.

Fracture Risk in Men With Congestive Heart Failure

content.onlinejacc.org/article.aspx?articleid=1139031 ▼ 翻译此页作者: LD Carbone - 2008 - 被引用次数: 46 - 相关文章 2008年7月8日 - Results We identified 167 cases with a single-incident fracture and matched Spironolactone is increasingly being used to treat hypertension and CHF dietary estimates of net endogenous acid production are associated with low ... factors be used to predict fractures in community-dwelling older adults?

Fracture Risk in Men With Congestive Heart Failure: Risk ...

www.sciencedirect.com/science/article/pii/S0735109708014174 - 翻译此页作者: LD Carbone - 2008 - 被引用次数: 45 - 相关文章

We identified 167 cases with a single-incident fracture and matched these by age and Age; race; height; weight; cigarette use; prevalent fractures of the hip, high dietary estimates of net endogenous acid production are associated with low ... risk factors be