

Name of Journal: *World Journal of Hypertension*

ESPS Manuscript NO: 29905

Manuscript Type: Minireviews

Reduction of diastolic blood pressure: should hypertension guidelines include a lower threshold target?

Steven Tringali, Jian Huang

## Match Overview

1	Crossref 30 words S. Bangalore. "J-curve revisited: an analysis of blood pressure and cardiovascular events in the Treating to New ...	1%
2	Internet 26 words crawled on 22-Oct-2014 <a href="http://circ.ahajournals.org">circ.ahajournals.org</a>	1%
3	Internet 21 words crawled on 13-Apr-2016 <a href="http://apiindia.org">apiindia.org</a>	1%
4	Internet 20 words crawled on 11-Feb-2014 <a href="http://www.kidneyonline.org">www.kidneyonline.org</a>	1%
5	Internet 17 words crawled on 15-Jun-2014 <a href="http://www.slideshare.net">www.slideshare.net</a>	1%
6	Internet 17 words crawled on 20-May-2016 <a href="http://www.bmj.com">www.bmj.com</a>	1%

学术搜索

找到约 28,500 条结果 (用时0.16秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用☐ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

## 2003 World Health Organization (WHO)/International Society of Hypertension (ISH) statement on management of hypertension

..., International Society of Hypertension Writing ... - ... of hypertension, 2003 - journals.lww.com  
... in 1999, more evidence has become available to support a systolic blood pressure threshold of 140 ... In order to achieve significant reductions in the avoidable CVD burden, a combination of ... Treating hypertension has been associated with about a 40% reduction in the risk of ...

被引用次数: 1742 相关文章 所有 8 个版本 引用 保存

## [HTML] 2014 evidence-based guideline for the management of high blood pressure adults: report from the panel members appointed to the Eighth Joint National ...

PA James, S Oparil, BL Carter, WC Cushman... - Jama, 2014 - jama.jamanetwork.com

Review from JAMA — 2014 Evidence-Based Guideline for the Management of High Blood Pressure in Adults — Report From the Panel Members Appointed to the Eighth Joint National Committee (JNC 8).

被引用次数: 2803 相关文章 所有 84 个版本 引用 保存

## [PDF] British Hypertension Society guidelines for hypertension management 2004 (BHS-IV): summary

B Williams, NR Poulter, MJ Brown, M Davis... - Bmj, 2004 - dablededucational.org

... for reduction of major cardiovascular events was 139/83 mm Hg and that reduction of blood ... the week • Consume at least five portions/day of fresh fruit and vegetables • Reduce the intake ... is no cost disadvantage to their use, they represent a sensible way of reducing the number ...

被引用次数: 867 相关文章 所有 23 个版本 引用 保存 更多

## Management guidelines in essential hypertension: report of the second working party of the British Hypertension Society.

D Sever, G Beevers, G Bullock, A Laver, J Peto... - Bmj, 1993 - bmi.com

学术搜索

找到约 27,300 条结果 (用时0.28秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National ...

PA James, S Oparil, BL Carter, WCushman... - Jama, 2014 - jama.jamanetwork.com

Review from JAMA — 2014 Evidence-Based Guideline for the Management of High Blood Pressure in Adults — Report From the Panel Members Appointed to the Eighth Joint National Committee (JNC 8).

被引用次数: 3099 相关文章 所有 84 个版本 引用 保存

2003 World Health Organization (WHO)/International Society of Hypertension (I statement on management of hypertension

..., International Society of Hypertension Writing ... - ... of hypertension, 2003 - journals.lww.com

... in 1999, more evidence has become available to support a systolic blood pressure threshold of 140 ... In order to achieve significant reductions in the avoidable CVD burden, a combination of ... Treating hypertension has been associated with about a 40% reduction in the risk of ...

被引用次数: 1784 相关文章 所有 8 个版本 引用 保存

Management guidelines in essential hypertension: report of the second working party of the British Hypertension Society.

P Sever, G Beevers, C Bulpitt, A Lever, L Ramsay... - Bmj, 1993 - bmj.com

... Non-pharmacological treatment \* Reduce energy intake \* Avoid excessive alcohol \* Reduce salt intake ... pressures achieved in those trials.\* Diastolic pressure should be reduced to less ... believe that the evidence is insufficient to recommend a limit on blood pressure reduction. ...

被引用次数: 369 相关文章 所有 13 个版本 引用 保存

[PDF] British Hypertension Society guidelines for hypertension management 200

[全部](#)[图片](#)[新闻](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 1,350,000 条结果 (用时 0.82 秒)

### Blood pressure targets for hypertension: What's the evidence? | BC ...

[www.bcmj.org](http://www.bcmj.org) > [Back to the October 2012 Issue](#) ▾ [翻译此页](#)

作者: NA Khan - [相关文章](#)

BP targets **can** differ from BP **thresholds**, which are the BP **criteria** used to decide ... This **includes** the pharmacological and nonpharmacological approaches ... The case for a **lower blood pressure target** for patients with this combination of ... with a **diastolic blood pressure** goal of less than 80 mm Hg had a 51% **reduction** in ...

### 2014 Guideline for Management of High Blood Pressure | Cardiology ...

[jamanetwork.com/journals/jama/fullarticle/1791497](http://jamanetwork.com/journals/jama/fullarticle/1791497) - [翻译此页](#)

作者: PA James - 2014 - 被引用次数: 3133 - [相关文章](#)

2013年12月18日 - Comparison of Current Recommendations With JNC 7 **Guidelines** ... **Target blood pressure** for treatment of isolated **systolic hypertension** in the ... Effects of intensive **blood-pressure** lowering and **low-dose** aspirin in patients ..... be assured that **blood pressure** (BP) treatment **will reduce** their disease burden, ...

### Perioperative hypertension management

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2515421/> - [翻译此页](#)

作者: J Varon - 2008 - 被引用次数: 94 - [相关文章](#)

In addition, there is a lack of consensus concerning treatment **thresholds** and ... **Blood pressure** should be **reduced** by 10%–15% (maximum of 20%) in a controlled ... In patients with chronic **hypertension**, as long as the **diastolic** BP is <110 mm Hg, .... Preferred agents **include** labetalol, esmolol, nicardipine, and fenoldopam.

### SIGN 97: Risk estimation and the prevention of cardiovascular disease ...

[www.sign.ac.uk/guidelines/fulltext/97/section10.html](http://www.sign.ac.uk/guidelines/fulltext/97/section10.html) ▾ [翻译此页](#)