

**Name of Journal:** *World Journal of Psychiatry*

**Manuscript NO:** 55614

**Manuscript Type:** ORIGINAL ARTICLE

*Observational Study*

Alcohol and drug use disorders in an adult attention-deficit/hyperactivity disorder clinical population: <sup>1</sup>Prevalence and associations with ADHD symptom severity and emotional dysregulation

Espen Anker, Jan Haavik, Trond Heir

## Match Overview

1 Internet 869 words  
crawled on 10-Apr-2020  
[www.researchsquare.com](http://www.researchsquare.com)

37%



Alcohol and drug use disorders in an adult attention-deficit/hyperac



ALL

IMAGES

VIDEOS

17,000 Results

Any time ▾

## Adult attention deficit hyperactivity disorder ...

<https://www.ncbi.nlm.nih.gov/pubmed/9040531>

**Adult ADHD** had high comorbidity with current depressive **disorder**, antisocial personality **disorder**, and **alcohol and drug** abuse/dependence. High correlations were found between patients' and independent observers' reports of **ADHD symptom severity**. Implications for further research are discussed. PMID: 9040531 [Indexed for MEDLINE] MeSH terms. **Adult**

Cited by: 329

Author: Karen K. Downey, Fred W. Stelson, Ovide F. ...

Publish Year: 1997

## [PDF] The Clinically Meaningful Link Between Alcohol Use and ...

<https://pubs.niaaa.nih.gov/publications/arh26-2/122-129.pdf>

Link Between **Alcohol Use** and **Attention Deficit Hyperactivity Disorder** Bradley H. Smith, Ph.D., Brooke S.G. Molina, Ph.D., and William E. Pelham, Jr., Ph.D. **Attention deficit hyperactivity disorder (ADHD)** is a childhood mental health **disorder** that can lead to **alcohol** and other **drug** (AOD)-related problems if it persists into adolescence and

## Substance-use disorders in adolescents and adults with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3480177>

**ADHD** is among the most prevalent neuro-behavioral **disorders** presenting for treatment in children and adolescents [1]. Between 6 and 9% of youths and 4–5% of adults have **ADHD** worldwide [2,3]. **ADHD** is associated with a persistent course, impairment in multiple domains, and co-occurring learning and psychiatric problems (for a review see [1]). ...

Cited by: 17

Author: Timothy E Wilens, Nicholas R Morrison

Publish Year: 2012

## Substance Abuse in Patients With Attention-Deficit ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2258479>

Jan 31, 2008 · Recognizing and Addressing SUD in **ADHD** Patients. Despite recommendations that every primary care practice should screen adolescent patients for **drug** and **alcohol use**, the American Academy of Pediatrics estimates that less than 50% of pediatricians actually perform such screening. Insufficient knowledge of treatment steps in the event of a positive screening and the age of the patient ...

Cited by: 25

Author: Oscar Bukstein





Alcohol and drug use disorders in adult attention-deficit/hyperactiv



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

16,200 Results

Any time ▾

## Smoking, alcohol and drug use in youth and adults with ...

<https://www.cambridge.org/core/journals/bjpsych...> ▾

Previous research suggests a relationship between **attention-deficit hyperactivity disorder** (ADHD) and smoking, **alcohol** and illicit **drug use**, however most studies have focused on adolescents or young adults, or clinically ascertained samples.

**Cited by:** 1

**Author:** Sydney Osland, Lauren Hirsch, Tamara P...

**Publish Year:** 2017

## Alcohol, drugs, and attention-deficit/hyperactivity ...

<https://www.researchgate.net/publication/7588593...>

Because of its early onset, high **prevalence** and known risk for SUD, **attention deficit hyperactivity disorder** (ADHD) is a model developmental **disorder** to evaluate in context to SUDs.

## Attention-Deficit/Hyperactivity Disorder in Relation to ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4403287>

Apr 20, 2015 - In a clinical population-based, birth-cohort study, it was found that childhood ADHD cases were 6.2 times more likely to have an **alcohol/drug use disorder** than non-ADHD controls from the same cohort, and that stimulant treatment tended to be a protective factor, but only in boys .

**Cited by:** 16

**Author:** Caroline Davis, Alina Cohen, Mark David...

**Publish Year:** 2015

## Adult attention-deficit/hyperactivity disorder and its ...

[https://www.researchgate.net/publication/277086155\\_Adult\\_attention-deficit...](https://www.researchgate.net/publication/277086155_Adult_attention-deficit...)

The worldwide **prevalence** of **adult** ADHD is estimated to be 2.8% in the general population (Fayyad et al., 2017). ADHD increases the risk for heavy substance **use** or ...

## Attention deficit hyperactivity disorder (ADHD) and ...

<https://www.researchgate.net/publication/11648190...>

Several studies have suggested a similar relationship for ADHD and substance **use disorder**, [1][2] [3] [4] such as that half of **drug** abusers had childhood ADHD, 4 or ADHD was associated with early ...

## Attention Deficit Hyperactivity Disorder

Medical condition

Attention-deficit/hyperactivity disorder (ADHD) is a brain disorder marked by an ongoing pattern of inattention and/or hyperactivity-impulsivity that interferes with functioning or development.

NIH

### Symptoms

There are three categories of ADHD symptoms in addition to three types of the disorder. Symptoms are classified as:

- inattention: getting distracted easily, having poor concentration and organizational skills
- impulsivity: interrupting, taking risks...

[Read more on Healthline](#)

### Causes

The cause of ADHD isn't known. Researchers say several things may lead to it, including:

- Heredity. ADHD tends to run in families.
- Chemical imbalance. Brain chemicals in people with ADHD may be out of balance.
- Brain changes. Areas of the brain that...

[Read more on WebMD](#)

### Treatments

Standard treatments for attention-deficit/hyperactivity disorder in children include medications, education, training and counseling. These treatments can relieve many of the symptoms of ADHD, but they don't cure it. It may take some time to...

[Read more on MayoClinic](#)