Factors causing a relapse of severe major depressive disorders folk







Q ALL

☑ IMAGES

■ I VIDEOS

11,900 Results

Any time ▼

Relapse following successful electroconvulsive therapy for ... europepmc.org/articles/PMC3799066 •

Jun 18, 2013 · INTRODUCTION. Electroconvulsive therapy (ECT) is a highly effective acute treatment for major depression (Eranti et al, 2007; Kellner et al, 2010; The UK ECT Review Group, 2003). Although remission rates exceed those seen with other somatic treatments, high rates of relapse, especially early relapse, are observed and acknowledged as a major clinical problem (Kellner et al, 2006; Sackeim et ...

Factors associated with relapse after a response to ...

https://www.sciencedirect.com/science/article/pii/S0165032716308059

Jan 15, 2017 · A number of factors that could lead to relapse following ECT for depression have been suggested, including symptom resistance to medication, residual symptoms, female gender, psychosis, and comorbid conditions (Medda et al., 2013).

Cited by: 17 Author: Kei

Author: Kei Itagaki, Minoru Takebayashi, Chiyo S...

Publish Year: 2017

| PEOPLE ALSO ASK | |
|---|---|
| Which is the best electroconvulsive therapy for depression? | ~ |
| What is the future of electroconvulsive therapy? | ~ |
| Who is most at risk for electroconvulsive therapy? | ~ |
| Who is the founder of electroconvulsive therapy? | ~ |

Feedback

Shared preventive factors associated with relapse after a ... https://onlinelibrary.wiley.com/doi/full/10.1111/pcn.12859

May 11, 2019 - Limitations in the current **study** include: (i) a **naturalistic** and retrospective design, rather than a prospective, randomized controlled trial; (ii) variation in the severity of the disorders; (iii) exclusion of **refractory** schizophrenia patients who received clozapine; (iv) inclusion of subjects who may have been 'refractory' (i.e., all patients in the current **study** received one or more trials of medication before ...

Cited by: 4

Author: Wataru Omori, Kei Itagaki, Naoto Kajitani...

26-Aug-2021 09:23PM

Name of Journal: World Journal of Psychiatry

Manuscript NO: 66444

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Factors causing a relapse of major depressive disorders following successful electroconvulsive therapy: A retrospective cohort study

Kurimoto N et al. Diagnostic change for MDD after ECT

Naoki Kurimoto, Takahiko Inagaki, Takashi Aoki, Hiroshi Kadotani, Fujiki Kurimoto, Kenichi Kuriyama, Naoto Yamada, Yuji Ozeki

| 6 | Match Overview | | E= late | |
|---|----------------|---|---------|--|
| | -4 | | Þ | |
| | 1 | Internet 34 words crawled on 01-May-2020 journals.sagepub.com | 1% | |
| | 2 | Crossref 17 words Hiroya lida, Masaji Tani, Fumitoshi Hirokawa, Masaki Ueno et a I. "Risk factors for incisional hernia according to different woun | <1% | |
| | 3 | Crossref 16 words M. R. M. Elghorori, I. Hassan, W. Dartey, E. Abdel-Aziz. "A w to lend objectivity to Bishop score", Journal of Obstetrics and | <1% | |
| | 4 | Internet 13 words crawled on 15-Apr-2016 www.dovepress.com | <1% | |
| | 5 | Internet 12 words crawled on 04-Feb-2020 www.science.gov | <1% | |
| | | | | |



Factors causing a relapse of major depressive disorders following s







ALL

IMAGES

VIDEOS

9,840 Results

Any time ▼

Relapse Following Successful Electroconvulsive Therapy for ...

https://www.nature.com/articles/npp2013149

Jun 18, 2013 · The impact of medication resistance and continuation pharmacotherapy on relapse following response to electroconvulsive therapy in major depression. J Clin Psychopharmacol 10:

. . .

Cited by: 181

Author: Ana Jelovac, Erik Kolshus, Declan M McL...

Publish Year: 2013

Factors associated with relapse after a response to ...

https://www.sciencedirect.com/science/article/pii/S0165032716308059

Jan 15, 2017 · Background. While **electroconvulsive therapy** (ECT) treatment for **depression** is highly effective, the high rate of **relapse** is a critical problem. The current **study** investigated **factors** associated with the risk of **relapse** in mood **disorders** in patients in which ECT was initially effective.

Cited by: 18 Author: Kei Itagaki, Minoru Takebayashi, Chiyo Shi...

Publish Year: 2017

People also ask

| How is electroconvulsive therapy used to treat depression? | ~ |
|--|---|
| Is there a link between REM latency post ECT and depression? | ~ |
| What is the impact of medication resistance on Relapse? | ~ |
| When did electroconvulsive therapy start in the US? | ~ |