

[Depression following hip fracture is associated with ...](#)

<https://bmgeriatr.biomedcentral.com/articles/10.1186/1471-2318-13-60>

Jun 17, 2013 - In the current study, depression post-hip fracture was associated with poorer ADL and slower walking speed six weeks and six months post-hip fracture. Depressed participants also had significantly poorer physical functioning, as reflected in lower Berg Balance scale scores at week 6.

Cited by: 60 Author: Anna C Phillips, Jane Upton, Niharika Arora...
Publish Year: 2013 Estimated Reading Time: 12 mins

[Predictors of Incident Depression After Hip Fracture Surgery](#)

<https://www.sciencedirect.com/science/article/pii/S1064748112605211>

Sep 01, 2007 - Depression after hip fracture surgery is increasingly recognized as an important and independent predictor of a less favorable functional recovery and mortality.7, 8 Prevalence figures of depression after a hip fracture in the elderly have been reported between 9% and 47% and largely exceed those of 2% and 10% reported for major and minor depressive disorder, respectively, in the general ...

Cited by: 61 Author: Richard C. Oude Voshaar, Sube Banerjee, ...
Publish Year: 2007

Search Tools

Turn off Hover Translation (关闭取词)

PEOPLE ALSO ASK

- What is the prevalence of depression after a hip fracture? ▾
- Are there any studies on depression after a fracture? ▾
- When is a broken hip more likely in the elderly? ▾

国内版 国际版

New-onset depression after hip fracture surgery among older patie



ALL IMAGES VIDEOS

67,300 Results

Any time ▾

Depressive symptoms post hip fracture in older adults are ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4307912>

Dec 16, 2014 · In the present study, we have reported that new onset depression results in both **phenotypic and functional alteration in peripheral T cells** in elderly hip fracture patients. Firstly, we report a decline in T cell frequency and absolute numbers in hip fracture patients with depressive symptoms compared with healthy controls, which is in agreement with previous reports of reduced ...

Cited by: 12

Author: Niharika Arora Duggal, Jane Upton, Anna ...

Publish Year: 2014

Depression following hip fracture is associated with ...

<https://bmccgeriatr.biomedcentral.com/articles/10.1186/1471-2318-13-60> ▾

Jun 17, 2013 · In the current study, depression post-hip fracture was associated with poorer ADL and **slower walking speed** six weeks and six months post-hip fracture. Depressed participants also had **significantly poorer physical functioning**, as reflected in lower Berg Balance scale scores at week 6.

Cited by: 60

Author: Anna C Phillips, Jane Upton, Niharika Aro...

Publish Year: 2013

Estimated Reading Time: 12 mins

PEOPLE ALSO ASK

What is the prevalence of depression after a hip fracture? ▾

Are there any studies on depression after a fracture? ▾

How many people die from a hip fracture? ▾

How is depression related to a medical event? ▾

Feedback

Predictors of Incident Depression After Hip Fracture Surgery

<https://www.sciencedirect.com/science/article/pii/S1064748112605211>

Sep 01, 2007 · Depression after hip fracture surgery is increasingly recognized as an important and independent predictor of a less favorable functional recovery and mortality.^{7, 8} Prevalence figures of depression after a hip fracture in the elderly have been reported between 9% and 47% and largely exceed