



27893-Review

Quotes Included
Bibliography Included17%
SIMILARName of Journal: *World Journal of Psychiatry*

ESPS Manuscript NO: 27893

Manuscript Type: ORIGINAL ARTICLE

Case Control Study

Voxel-based magnetic resonance imaging investigation of poor and preserved clinical insight in people with schizophrenia

Adegboyega Sapara, Dominic H ffytche, Michael A Cooke, Steven CR Williams, Veena Kumari

ABSTRACT

AIM: To define regional grey-matter abnormalities in schizophrenia patients with poor insight (Insight⁻), relative to patients with preserved clinical insight (Insight⁺), and healthy controls.

Match Overview

- | Rank | Source | Words | Similarity |
|------|--|-----------|------------|
| 1 | CrossCheck Premkumar, P... "Structural magnetic resonance imaging ... edictors of responsiveness to cognitive behaviour therapy | 114 words | 3% |
| 2 | CrossCheck Sapara, Adegboyega, Dominic H ffytche, Michael A. Cooke, Steven C.R. Williams, and Veena Kumari. "Is it me? Ve... | 101 words | 3% |
| 3 | CrossCheck "Abstracts of the 11th International Congress on Schizophrenia Research", Schizophrenia Bulletin, 01/29/2007 | 69 words | 2% |
| 4 | CrossCheck Premkumar, P... "Emotional decision-making and its dissociable components in schizophrenia and schizoaffective di... | 67 words | 2% |
| 5 | Internet crawled on 06-Aug-2015 www.ncbi.nlm.nih.gov | 54 words | 1% |
| 6 | CrossCheck Sapara, A... "Prefrontal cortex and insight in schizophrenia: A volumetric MRI study", Schizophrenia Research, 200701 | 49 words | 1% |



全部 图片 新闻 视频 更多 ▾ 搜索工具

找到约 81,600 条结果 (用时 0.96 秒)

Google 学术: A Voxel-based MRI Investigation of Poor and Preserved Clinical Insight in Schizophrenia

... bipolar disorder investigated with MRI and voxel-based ... - Nugent - 被引用次数: 171

... cortex and insight in schizophrenia: a volumetric MRI ... - Sapara - 被引用次数: 88

Mapping grey matter reductions in schizophrenia: an ... - Fornito - 被引用次数: 186

Schizophrenia with preserved insight is associated with ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: C Faget-Agius - 2012 - 被引用次数: 18 - 相关文章

In this study, we investigated the functional substrate underlying preserved

Sociodemographic and clinical characteristics of patients with schizophrenia with

Neurological basis of poor insight in psychosis: a voxel-based MRI study.

The neuroanatomical correlates of cognitive insight in ...

scan.oxfordjournals.org/content/8/4/418.full ▾ 翻译此页

作者: MD Orfei - 2013 - 被引用次数: 20 - 相关文章

2012年1月28日 - Thus, cognitive insight complements clinical insight, because it ... of schizophrenia, a lower cognitive insight index indicates a poor ... Over the past 20 years, the neuroanatomical correlates of clinical insight were investigated, Neurological

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 69,400 条结果 (用时 0.68 秒)

Google 学术: Voxel-based magnetic resonance imaging investigation of poor and preserved clinical insight in people with schizophrenia

... in schizophrenia: a quantitative magnetic resonance ... - Shenton - 被引用次数: 1174

... network of schizophrenia patients with poor insight - Liemburg - 被引用次数: 25

Diffusion indices on magnetic resonance imaging and ... - Rose - 被引用次数: 173

Schizophrenia with preserved insight is associated with ... - NCBI

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ [翻译此页](#)

作者: C Faget-Agius - 2012 - 被引用次数: 19 - 相关文章

In this study, we investigated the functional substrate underlying preserved Sociodemographic and clinical characteristics of patients with schizophrenia with Neurological basis of poor insight in psychosis: a voxel-based MRI study.

Person - OpenAIRE

https://www.openaire.eu/search/person?personId=dedup_wf_001... ▾ [翻译此页](#)

It involved 40 stable schizophrenia outpatients, 20 with preserved and 20 wit... Neurological basis of poor insight in psychosis: A voxel-based MRI study ... to cognitive behaviour therapy in schizophrenia: a functional MRI investigation. ... poor insight, people with schizophrenia often disagree with carers and clinicians about ...

Insight in schizophrenia: From conceptualization to neuroscience ...

onlinelibrary.wiley.com > ... > Vol 66 Issue 3 > Abstract ▾ [翻译此页](#)

作者: M Ouzir - 2012 - 被引用次数: 19 - 相关文章

2012年3月23日 - To improve the treatment of people with schizophrenia, it is thus crucial ...

Furthermore, poor insight in schizophrenia has been proposed to result Engh et al. investigated the same question in a cross-sectional study of 143 patients. some studies have used magnetic resonance imaging-voxel-based

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 68,000 条结果 (用时 0.72 秒)

Google 学术: Voxel-based magnetic resonance imaging investigation of poor and preserved clinical insight in people with schizophrenia

... in schizophrenia: a quantitative magnetic resonance ... - Shenton - 被引用次数: 1177

... network of schizophrenia patients with poor insight - Liemburg - 被引用次数: 26

Diffusion indices on magnetic resonance imaging and ... - Rose - 被引用次数: 173

Schizophrenia with preserved insight is associated with increased ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ [翻译此页](#)

作者: C Faget-Agius - 2012 - 被引用次数: 19 - 相关文章

In this study, we investigated the functional substrate underlying preserved Sociodemographic and clinical characteristics of patients with schizophrenia with Neurological basis of poor insight in psychosis: a voxel-based MRI study.

Insight in schizophrenia: From conceptualization to neuroscience ...

onlinelibrary.wiley.com > ... > Vol 66 Issue 3 > Abstract ▾ [翻译此页](#)

作者: M Ouzir - 2012 - 被引用次数: 20 - 相关文章

2012年3月23日 - To improve the treatment of people with schizophrenia, it is thus crucial ... Furthermore, poor insight in schizophrenia has been proposed to result Engh et al. investigated the same question in a cross-sectional study of 143 patients. some studies have used magnetic resonance imaging, voxel-based ...

Neurobiological Basis of Insight in Schizophrenia: A Systema ...

journals.lww.com > Home > Current Issue - [翻译此页](#)

Background: Insight in schizophrenia is defined as awareness into illness, ... of depression in first-episode psychosis (), whereas poor or impaired insight is are commonly associated with clinical insight and its dimensions. using MRI ... of the precuneus bilaterally in the preserved insight patients