

Name of Journal: World Journal of Transplantation

ESPS Manuscript NO: 21115

Manuscript Type: REVIEW

Reducing transfusion requirements in liver transplantation

Ciara I Donohue, Susan V Mallett

Abstract

Liver transplantation was historically associated with massive blood loss and transfusion. Over the past two decades transfusion requirements have reduced dramatically and increasingly transfusion-free transplantation is a reality. Both bleeding and transfusion are associated with adverse outcomes in liver transplantation. Minimising bleeding and reducing unnecessary transfusions are therefore key goals in the perioperative period. As the understanding of the causes of bleeding has evolved so too have techniques

Match Overview

1	CrossCheck 53 words &NA; - "Ceftriaxone injection more costly than cefotaxime infusion", Inpharma Weekly, 1997.	1%
2	CrossCheck 43 words Ilona T. A. Pereboom. "Platelet Transfusion During Liver Transplantation Is Associated with Increased Postoperative ...	1%
3	CrossCheck 33 words Mallett, Susan. "Clinical Utility of Viscoelastic Tests of Coagulation (TEG/ROTEM) in Patients with Liver Disease an ...	1%
4	CrossCheck 22 words Camus, Marine. "Epistaxis in end stage liver disease may be a heralding as severe upper gastrointestinal hemorrhage", W	<1%
5	CrossCheck 19 words "Acute Liver Failure", Encyclopedia of Trauma Care, 2015.	<1%
6	CrossCheck 19 words Susan S. Yoo. "Coagulation Abnormalities Made Easy :". A SA Refresher Courses in Anesthesiology, 2010	<1%

[网页](#) [图片](#) [学术](#) [高级搜索](#)

Transfusion practice in orthotopic liver transplantation

www.ncbi.nlm.nih.gov/pmc/articles/PMC2823092/

Liver transplant procedures require the most blood components, despite the as short as possible to reduce the intraoperative RBC transfusion requirement as....

Recombinant factor VIIa reduces transfusion requirements in liver ...

www.ncbi.nlm.nih.gov/pubmed/16623915

Transfus Med. 2006 Apr.16(2):93-100. Recombinant factor VIIa reduces transfusion requirements in liver transplant patients with high MELD scores. Niemann....

Reduced transfusion requirements by recombinant factor VIIa in ...

www.ncbi.nlm.nih.gov/pubmed/11233901

Reduced transfusion requirements by recombinant factor VIIa in orthotopic liver transplantation: a pilot study. Hendriks HG(1), Meijer K, de Wolf JT, Klompmaker....

Transfusion Requirements in Liver Transplantation: Overview ...

emedicine.medscape.com/article/431573-overview

25 Nov 2013 ... Liver transplantation has emerged as an increasingly successful treatment for patients with end-stage liver disease (ESLD). Orthotopic liver....

[PDF] Blood loss, predictors of bleeding, transfusion practice and ...

office.wjgnet.com/1948-5182/pdf/v5/i1/1.pdf

27 Jan 2013 ... how urgently the patient needs a liver transplant within the next three control blood loss and reduce transfusion requirements should be....



[网页](#) [新闻](#) [图片](#) [视频](#) [购物](#) [更多](#) [搜索工具](#)

找到约 621,000 条结果 (用时 0.49 秒)

Google 学术: Reducing transfusion requirements in liver transplantation

Reduced transfusion requirements by recombinant ... - Hendriks - 被引用次数: 262

... acid reduces blood loss, transfusion requirements, and ... - Boylan - 被引用次数: 165

... red blood cell transfusion in liver transplantation: ... - Ramos - 被引用次数: 167

Transfusion practice in orthotopic liver transplantation

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: AS Devi - 2009 - 被引用次数: 9 - [相关文章](#)

Keywords: Blood loss, factors, liver transplantation, transfusion. Go to: [Reduced transfusion requirements](#) by recombinant factor VIIa in orthotopic liver ...

Reduced transfusion requirements by recombinant factor VIIa

www.ncbi.nlm.nih.gov/pubmed/11233901 - [翻译此页](#)

作者: HG Hendriks - 2001 - 被引用次数: 262 - [相关文章](#)

[Reduced transfusion requirements](#) by recombinant factor VIIa in orthotopic liver transplantation: a pilot study. Hendriks HG(1), Meijer K, de Wolf JT, Klompmaker ...

Conjugated estrogen reduces transfusion and coagulation ...



[网页](#) [新闻](#) [图片](#) [视频](#) [地图](#) [更多 ▾](#) [搜索工具](#)

找到约 1,120,000 条结果 (用时 0.44 秒)

Google 学术: Reducing transfusion requirements in liver transplantation

[Reduced transfusion requirements by recombinant ...](#) - Hendriks - 被引用次数: 263

[... acid reduces blood loss, transfusion requirements, and ...](#) - Boylan - 被引用次数: 166

[... red blood cell transfusion in liver transplantation: ...](#) - Ramos - 被引用次数: 169

Transfusion practice in orthotopic liver transplantation

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: AS Devi - 2009 - 被引用次数: 9 - [相关文章](#)

[Liver transplant](#) procedures require the most blood components, despite the ... as short as possible [to reduce](#) the intraoperative RBC [transfusion requirement](#) as ...

Transfusion and coagulation management in liver ...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: B Clevenger - 2014 - 被引用次数: 14 - [相关文章](#)

2014年5月28日 - Keywords: [Liver](#) disease, [Transplantation](#), Coagulation, Transfusion, Patient ... Universal leukoreduction of blood has helped [to reduce](#) this risk. ... In general increased [transfusion requirements](#) for bleeding lead to higher ...

Transfusion Requirements in Liver Transplantation ...

emedicine.medscape.com/article/431573-overview ▾ [翻译此页](#)

2013年11月25日 - Overview; History and Background; Bleeding During [Liver Transplantation](#); [Transfusion Requirements](#); Operative Techniques That Minimize ...

Impact of blood salvage during liver transplantation on ...

www.scielo.br/scielo.php?script=sci_arttext&pid=S0102... ▾ [翻译此页](#)

作者: ... - 2000 - 被引用次数: 10 - [相关文章](#)