



**Name of Journal:** *World Journal of Transplantation*

**ESPS Manuscript NO:** 26461

**Manuscript Type:** Retrospective Cohort Study

**Thromboelastographic reference ranges for a cirrhotic patient population undergoing liver transplantation**

**Lesley De Pietri, Marcello Bianchini, Gianluca Rompianesi, Elisabetta Bertellini, Bruno Begliomini.**

## Match Overview

1	Internet 108 words crawled on 07-Mar-2016 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	2%
2	Internet 97 words crawled on 31-May-2016 <a href="http://www.scielo.br">www.scielo.br</a>	2%
3	Internet 68 words crawled on 29-Sep-2014 <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	1%
4	Internet 58 words crawled on 21-Jan-2010 <a href="http://www.anesthesia-analgesia.org">www.anesthesia-analgesia.org</a>	1%
5	CrossCheck 49 words Reichert, Benedikt, Alexander Kaltenborn, Thomas Beck er, Mario Schiffer, Jürgen Klempnauer, and Harald Schre	1%
6	CrossCheck 32 words De Pietri, Lesley, Marcello Bianchini, Roberto Montalti, Ni cola De Maria, Tommaso Di Maira, Bruno Begliomini, Gi	1%



学术搜索

找到约 811 条结果 (用时0.07秒)

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☐ 包括专利☐ 包含引用

[HTML] Intraoperative hypercoagulability during liver transplantation as demonstrated by thromboelastography

D Krzanicki, A Sugavanam, S Mallett - Liver Transplantation, 2013 - Wiley Online Library

... 9] Coagulation monitoring using viscoelastic coagulation tests [eg, thromboelastography (TEG) and ... patients with high G traces had platelet counts above the normal reference range. ... liver transplantation present with or develop hypercoagulable thromboelastograms during the ...

被引用次数: 39 相关文章 所有 4 个版本 引用

The coagulopathy of cirrhosis assessed by thromboelastometry and its correlation with conventional coagulation parameters

A Tripodi, M Primignani, V Chantarangkul, Y Viscardi... - Thrombosis research, 2009 - Elsevier

... None. References. ... Multi-centre investigation on reference ranges for ROTEM thromboelastometry. Blood Coagul Fibrinolysis, 16 (2005), pp. ... Evaluation of the contribution of platelets to clot strength by thromboelastography in rabbits: the role of tissue factor and cytochalasin D. ...

被引用次数: 78 相关文章 所有 7 个版本 引用

[HTML] Efficacy and safety of repeated perioperative doses of recombinant factor VIIa in liver transplantation

JPA Lodge, S Jonas, RM Jones... - Transplantation, 2005 - Wiley Online Library

... plasma concentrate, cryoprecipitate, and fibrinogen concentrate transfusions could also be administered as indicated by thromboelastogram tracings. ... The coagulation values observed in the study population were outside the range for healthy volunteers; such ... References. ...

被引用次数: 218 相关文章 所有 12 个版本 引用

Transfusion medicine service policies for recombinant factor VIIa administration

LT Goodnough, DM Lublin, L Zhang, G Despotis... - ..., 2004 - Wiley Online Library

... trial, 10 cirrhotic patients whose PT did not correct to within 2 seconds above the control reference value were given ... REFERENCES. ... VIIa is used in a Jehovah's witness with liver cirrhosis to correct



Your search - **"THROMBOELASTOGRAPHY REFERENCE RANGES FOR A CIRRHOTIC PATIENT POPULATION UNDERGOING LIVER TRANSPLANTATION"** - did not match any articles.

Suggestions:

Make sure all words are spelled correctly.

Try different keywords.

Try more general keywords.

Try fewer keywords.

[Try your query on the entire web.](#)

[About Google Scholar](#)

[Privacy](#)

[Terms](#)

[Provide feedback](#)

[全部](#)
[图片](#)
[新闻](#)
[视频](#)
[更多 ▾](#)
[搜索工具](#)

找到约 4,940 条结果 (用时 0.79 秒)

您是不是要找: **Thromboelastography** reference ranges for a cirrhotic patient population undergoing liver transplantation

### Intraoperative hypercoagulability during liver ... - Wiley Online Library

[onlinelibrary.wiley.com/doi/10.1002/lt.23668/pdf](https://onlinelibrary.wiley.com/doi/10.1002/lt.23668/pdf) ▾ 翻译此页

作者: D Krzanicki - 2013 - 被引用次数: 41 - 相关文章

**Thromboelastography** ... hypercoagulation is associated with complications in many **populations**, but the incidence in **liver transplant** ... Six of the **patients** developed early **hepatic** artery thrombosis (5%); 3 of these **patients** .... We examined a cohort of **patients undergoing** ... TEG tracing and **reference ranges** for TEG parame-

### Intraoperative hypercoagulability during liver transplantation as ...

[onlinelibrary.wiley.com/doi/10.1002/lt.23668/full](https://onlinelibrary.wiley.com/doi/10.1002/lt.23668/full) - 翻译此页

作者: D Krzanicki - 2013 - 被引用次数: 41 - 相关文章

2013年7月8日 - **thromboelastography**. α ... [2] reported venous thromboembolism events in 0.5% of **cirrhotic** ... We examined a cohort of **patients undergoing liver transplantation** to quantify and describe this issue. ... TEG tracing and **reference ranges** for TEG parameters. ... Etiologies of Liver Disease in the Study **Population** ...

### Potential Applications of Thromboelastography in Patients with Acute ...

[www.ncbi.nlm.nih.gov](https://www.ncbi.nlm.nih.gov) ▸ NCBI ▸ Literature ▸ PubMed Central (PMC) ▾ 翻译此页

作者: RT Stravitz - 2012 - 被引用次数: 30 - 相关文章

Studies using TEG in **patients with cirrhosis** and acute liver failure have .... within 30 minutes of reaching maximum amplitude; this **value** is a measure of fibrinolysis. ... in **patients undergoing liver transplantation** (LT). 19 During the preanhepatic .... sicker **population** (Table 3); the a-angle was also depressed in **patients with** ...