

Name of Journal: *World Journal of Transplantation*

Manuscript NO: 38412

Manuscript Type: Original Article

Observational Study

Cumulative ²³ positive fluid balance is a risk factor for acute kidney injury and requirement for renal replacement therapy after liver transplantation

Codes L *et al.* Fluid overload and liver transplantation

Liana Codes, Ygor Gomes de Souza, Ricardo Azevedo Cruz D'Oliveira, Jorge Luiz Andrade Bastos, Paulo Lisboa Bittencourt

符合内容总览

1	Crossref 57 个字 Francoz, Claire, Mitra K. Nadim, and Fran e Durand. "Kidney Biomarkers in Cirrhosis"	2%
2	互联网 23 个字 于 2017 年 12 月 29 日 缓慢进行 www.hsd-fmsb.org	1%
3	互联网 22 个字 于 2014 年 02 月 19 日 缓慢进行 www.shifa.com.pk	1%
4	互联网 20 个字 于 2016 年 06 月 10 日 缓慢进行 onlinelibrary.wiley.com	1%
5	互联网 20 个字 于 2017 年 11 月 24 日 缓慢进行 journals.plos.org	1%
6	互联网 19 个字 于 2017 年 06 月 02 日 缓慢进行 e-sciencecentral.org	1%
7	互联网 17 个字 于 2006 年 01 月 09 日 缓慢进行 clincancerres.aacrjournals.org	1%
8	Crossref 17 个字 "Acute Cardiovascular Care 2014", European Heart Journal: Acute Cardiovascular Care	1%
9	Crossref 15 个字 Sebagh, M., "Place of the liver biopsy in liver transplantation" Journal of Hepatology 2004	<1%
10	互联网 15 个字 于 2016 年 10 月 15 日 缓慢进行 www.elsevier.es	<1%
11	Crossref 14 个字 "Supplement 2 September 2011", Intensive C are, Medicine 09/2011	<1%



All

News

Images

Videos

Shopping

More

Settings

Tools

About 241,000 results (0.79 seconds)

Fluid overload is an independent risk factor for acute kidney injury in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5286805/> ▼

by N Salahuddin - 2017 - Cited by 4 - Related articles

Feb 1, 2017 - Fluid overload is an independent **risk factor** for **acute kidney injury** in critically ill patients: results of a cohort study that patients admitted to the ICU may be at a higher risk of **AKI** if they had a net **positive fluid balance** or received chloride— liberal fluids for both resuscitation and as maintenance **therapy**.

Chloride-liberal fluids are associated with acute kidney injury after ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4258383/> ▼

by A Nadeem - 2014 - Cited by 35 - Related articles

Nov 19, 2014 - We hypothesized that **AKI** occurring in the early postoperative period **after liver transplant** may be associated with the use of chloride-liberal fluids and overall In the first 72 hours of ICU stay, transplant recipients were in a **cumulative positive fluid balance** of 12.7 ± 7.8 L with 9 ± 6 L crystalloids, 6.4 ± 3.5 L ...

Influence of Daily Fluid Balance prior to Continuous Renal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4951567/> ▼

by MJ Han - 2016 - Cited by 5 - Related articles

May 23, 2016 - **Positive fluid balance** is a **risk factor** for mortality in critically ill patients, especially those **requiring** continuous renal replacement therapy (CRRT). ... **Acute kidney injury (AKI)** frequently occurs in critically ill patients (6,7,8), and fluid therapy is performed to maintain adequate renal perfusion in patients with or ...

Fluid overload at initiation of renal replacement therapy is associated ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3471547/>

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)[已启用安全](#)

找到约 145,000 条结果 (用时 0.62 秒)

Fluid accumulation, survival and recovery of kidney function in ...

<https://www.sciencedirect.com/science/article/pii/S0085253815539756> - [翻译此页](#)作者: J Bouchard - 2009 - [被引用次数: 672](#) - [相关文章](#)

2009年8月2日 - The Program to Improve Care in Acute Renal Disease (PICARD) began as an observational study of 618 critically ill patients with AKI from five academic ... Patients on continuous renal replacement therapy were more likely to reduce the percentage of fluid accumulation compared with patients treated with ...

Prevention and Therapy of Acute Kidney Injury in the Developing ...

<https://www.sciencedirect.com/science/article/pii/S2468024917300980> - [翻译此页](#)作者: V Kher - 2017 - [相关文章](#)

Volume depletion is one of the major risk factors for AKI. In contrast, some suspected, confirmed, and/or persistent AKI can present with volume overload and worsening renal function, such as congestive heart failure, or cardiorenal syndrome. Fluid overload was defined as the difference between cumulative fluid intake and ...

Influence of Daily Fluid Balance prior to Continuous Renal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4951567/> - [翻译此页](#)作者: MJ Han - 2016 - [被引用次数: 6](#) - [相关文章](#)

2016年5月23日 - Positive fluid balance is a risk factor for mortality in critically ill patients, especially those requiring continuous renal replacement therapy (CRRT). ... Acute kidney injury (AKI) frequently occurs in critically ill patients (6,7,8), and fluid therapy is performed to maintain adequate renal perfusion in patients with or ...

Fluid overload is an independent risk factor for acute kidney injury in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5286805/> - [翻译此页](#)作者: N Salahuddin - 2017 - [被引用次数: 5](#) - [相关文章](#)

2017年2月1日 - In patients with established AKI, further fluid overloading has been linked with lower

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 222,000 条结果 (用时 0.54 秒)

Fluid Balance in Patients with Acute Kidney Injury: Emerging Concepts ...

<https://www.karger.com/Article/FullText/354713> ▾ 翻译此页

作者: M Godin - 2013 - 被引用次数: 30 - 相关文章

2013年9月4日 - Intensive care unit and surgical populations are at increased risk for acute kidney injury (AKI) and oliguria, which often lead to fluid accumulation. ... Hu Y, Song PX, Ojo AO: Fluid overload at initiation of renal replacement therapy is associated with lack of renal recovery in patients with acute kidney injury.

Influence of Daily Fluid Balance prior to Continuous Renal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4951567/> ▾ 翻译此页

作者: MJ Han - 2016 - 被引用次数: 6 - 相关文章

2016年5月23日 - Positive fluid balance is a risk factor for mortality in critically ill patients, especially those requiring continuous renal replacement therapy (CRRT). ... Acute kidney injury (AKI) frequently occurs in critically ill patients (6,7,8), and fluid therapy is performed to maintain adequate renal perfusion in patients with or ...

Outcomes from a cohort of patients with acute kidney injury subjected ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5398654/> - 翻译此页

作者: TQC dos Santos - 2017 - 相关文章

2017年4月20日 - Several observational studies have linked positive fluid balance with AKI occurrence and mortality, especially before RRT initiation [8, 14–16]. However, only a small number of studies have evaluated the association between fluid overload and mortality during renal replacement therapy [11,15,17, 18].

Chloride-liberal fluids are associated with acute kidney injury after ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4258383/> ▾ 翻译此页

作者: A Nadeem - 2014 - 被引用次数: 36 - 相关文章

2014年11月19日 - Acute kidney injury (AKI) occurs frequently after liver transplantation and is associated with