

Name of Journal: *World Journal of Virology*

ESPS Manuscript NO: 22968

Manuscript Type: MINIREVIEWS

Hepatitis C virus / human T lymphotropic virus/2 co-infection: regional burden and virological outcomes in people who inject drugs

Erika Castro, Elena Roger

Erika Castro, Addiction Medicine Centre, Service of Community Psychiatry,
Department of Psychiatry, Centre Hospitalier Universitaire Vaudois, CH-1003
Lausanne, Switzerland

Match Overview

- 1** **CrossCheck** 25 words 1%
Leal, Fabio E., Lishomwa C. Ndhlovu, Aaron M. Hasenkrug, Fernanda R. Bruno, Karina I. Carvalho, Harry Wynn-William
- 2** **CrossCheck** 25 words 1%
Araya, Natsumi, Tomoo Sato, Naoko Yagishita, Hitoshi Ando, Atae Utsunomiya, Steven Jacobson, and Yoshihisa Yam



Hepatitis C virus / human T lymphotropic virus/2 co-infection: regional bur

[全部](#)[新闻](#)[图片](#)[视频](#)[更多](#)[搜索工具](#)

找到约 3,850 条结果 (用时 0.75 秒)

The Impact of Human T-Cell Lymphotropic Virus I Infection ...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: F Bahia - 2011 - 被引用次数: 9 - 相关文章

HIV, hepatitis C (HCV), and human T-cell lymphotropic virus I (HTLV-1) are ... of triply infected patients (Group 2) reported any history of injection drug use ... HIV/HTLV-1 and HIV/HCV coinfections may worsen clinical related outcomes, but virologic ... HIV and HTLV-1 coinfection has been found in Bahia with worse morbidity ...

[PDF] Prevention and control of infectious diseases among peo...

ecdc.europa.eu/en/.../111012_Guidance_ECDC-EMCDDA.pdf ▾ [翻译此页](#)

interventions to prevent infections among people who inject drugs. a large burden on European health systems, significant individual suffering, with more than one type of hepatitis, or co-infection with HIV and hepatitis C ... In addition to HIV and hepatitis B and C, infections caused by human T-lymphotropic virus type II ...

Burden of hepatitis C virus disease and access to hepatitis ...

www.lancet.com/journals/laninf/article/PIIS1473...X/fulltext ▾ [翻译此页](#)

2014年12月2日 - 90% of individuals infected with hepatitis C virus (HCV) worldwide reside in ... Weighted HCV prevalence was 5777 (37.2%) of 14 447; HIV/HCV co-infection ... Of the 5777 people who inject drugs that were HCV antibody positive, 440

human immunodeficiency, and human T-lymphotropic viruses



Hepatitis C virus / human T lymphotropic virus/2 co-infection: regional bur

[全部](#)[新闻](#)[图片](#)[视频](#)[更多](#)[搜索工具](#)

找到约 3,850 条结果 (用时 0.75 秒)

The Impact of Human T-Cell Lymphotropic Virus I Infection ...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: F Bahia - 2011 - 被引用次数: 9 - 相关文章

HIV, hepatitis C (HCV), and human T-cell lymphotropic virus I (HTLV-1) are ... of triply infected patients (Group 2) reported any history of injection drug use ... HIV/HTLV-1 and HIV/HCV coinfections may worsen clinical related outcomes, but virologic ... HIV and HTLV-1 coinfection has been found in Bahia with worse morbidity ...

[PDF] Prevention and control of infectious diseases among peo...

ecdc.europa.eu/en/.../111012_Guidance_ECDC-EMCDDA.pdf ▾ [翻译此页](#)

interventions to prevent infections among people who inject drugs. a large burden on European health systems, significant individual suffering, with more than one type of hepatitis, or co-infection with HIV and hepatitis C ... In addition to HIV and hepatitis B and C, infections caused by human T-lymphotropic virus type II ...

Burden of hepatitis C virus disease and access to hepatitis ...

www.lancet.com/journals/laninf/article/PIIS1473...X/fulltext ▾ [翻译此页](#)

2014年12月2日 - 90% of individuals infected with hepatitis C virus (HCV) worldwide reside in ... Weighted HCV prevalence was 5777 (37.2%) of 14 447; HIV/HCV co-infection ... Of the 5777 people who inject drugs that were HCV antibody positive, 440

human immunodeficiency, and human T-lymphotropic viruses

[全部](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 3,680 条结果 (用时 1.32 秒)

The Impact of Human T-Cell Lymphotropic Virus I Infection ...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: F Bahia - 2011 - 被引用次数: 9 - 相关文章

HIV, hepatitis C (HCV), and human T-cell lymphotropic virus I (HTLV-1) are ... of triply infected patients (Group 2) reported any history of injection drug use ... HIV/HTLV-1 and HIV/HCV coinfections may worsen clinical related outcomes, but virologic ... HIV and HTLV-1 coinfection has been found in Bahia with worse morbidity ...

Burden of hepatitis C virus disease and access to hepatitis ...

www.lancet.com/journals/laninf/article/PIIS1473...X/fulltext ▾ [翻译此页](#)

2014年12月2日 - 90% of individuals infected with hepatitis C virus (HCV) worldwide reside in ... HCV care continuum outcomes were self-reported except for viral clearance in ... Weighted HCV prevalence was 5777 (37.2%) of 14 447; HIV/HCV co-infection ... Of the 5777 people who inject drugs that were HCV antibody ...

^[PDF] Prevention and control of infectious diseases among peo...

ecdc.europa.eu/en/.../111012_Guidance_ECDC-EMCDDA.pdf ▾ [翻译此页](#)

interventions to prevent infections among people who inject drugs. a large burden on European health systems, significant individual suffering, with more than one type of hepatitis, or co-infection with HIV and hepatitis C ... In addition to HIV and hepatitis B and C, infections caused by human T-lymphotropic virus type II ...

Prevalence of HIV, Hepatitis B and C Infections and an ...

journals.plos.org/plosone/article?id=10.1371/journal.pone... ▾ [翻译此页](#)

作者: HT Kinkel - 2015

2015年8月11日 - The prevalence rates of HBsAg, anti-HIV antibodies and HCV-RNA were 3.5%, ... duration of injecting drug use, history of imprisonment and HIV co-infection.