

Name of Journal: *World Journal of Nephrology*

ESPS Manuscript NO: 22385

Manuscript Type: Original Article

Observational Study

Matrix metalloproteinase-2 as a superior biomarker for peritoneal deterioration in
peritoneal dialysis

Ichiro Hirahara, Eiji Kusano, Yoshiyuki Morishita, Makoto Inoue, Tetsu Akimoto,
Osamu Saito, Shigeaki Muto, Daisuke Nagata

Match Overview

1	CrossCheck 222 words Hirahara, I., M. Inoue, T. Umino, O. Saito, S. Muto, and E. Kusano. "Matrix metalloproteinase levels in the drained dia	5%
2	CrossCheck 130 words E. Kusano. "The potential of matrix metalloproteinase-2 as a marker of peritoneal injury, increased solute transport, o	3%
3	CrossCheck 127 words "2009 Japanese Society for Dialysis Therapy Guidelines r Peritoneal Dialysis - 2009 JSOT Guidelines for Peritoneal	3%
4	Internet 47 words crawled on 22-May-2009 pdconnect.com	1%
5	CrossCheck 45 words T. Fujita. "Methylglyoxal induces peritoneal thickening by mesenchymal-like mesothelial cells in rats". <i>Nephrology Di</i>	1%
6	CrossCheck 31 words Suwandil, , Seishi Akino, and Nana Kondo. "Common Spea r Rot of Oil Palm in Indonesia". <i>Plant Disease</i> , 2012.	1%
7	Publications 26 words "New Kidney Disease Findings from D.L. Barreto and Co authors Described (Peritoneal Effluent MMP-2 an". <i>Health &</i>	1%
8	CrossCheck 25 words Hirahara, Ichiro, Tetsu Akimoto, Yoshiyuki Morishita, Makot o Inoue, Osamu Saito, Shigeaki Muto, and Eiji Kusano. "	1%

学术搜索

获得 4 条结果 (用时0.06秒)

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[PDF] ... C levels reflect peritoneal deterioration in peritoneal dialysis: MAJOR IN PD (Multicenter Analysis in Japan, ORiginal Indicator of Peritoneal Deterioration) study I Hirahara, E Kusano, T Imai, Y Morishita, M Inoue... - downloads.hindawi.com
... We suggest that MMP-2 should serve as a superior indicator of general peritoneal deterioration. ...
9. DL Barreto, AM Coester, DG Struijk, RT Krediet RT, "Can effluent matrix metalloproteinase 2 and plasminogen activator inhibitor 1 be used as biomarkers of peritoneal ...
引用 保存 更多

Effluent Tenascin-C Levels Reflect Peritoneal Deterioration in Peritoneal Dialysis: MAJOR IN PD Study
I Hirahara, E Kusano, T Imai, Y Morishita... - BioMed Research ..., 2015 - hindawi.com
... We suggest that MMP-2 should serve as a superior indicator of general peritoneal ... Therefore, effluent TN-C has a possibility of biomarker for EMT-like change in ... DL Barreto, AM Coester, DG Struijk, and RT Krediet, "Can effluent matrix metalloproteinase 2 and plasminogen ...
引用 保存 更多

[HTML] Peritoneal dialysis-2
T Kusumoto, K Fukami, S Yamagishi, S Ueda... - NDT ..., 2009 - ckj.oxfordjournals.org
We use cookies to enhance your experience on our website. By continuing to use our website, you are agreeing to our use of cookies. You can change your cookie settings at any time. Find out more. Skip Navigation. ...
所有 2 个版本 引用 保存

Methods, kits, reagents and devices for detecting mesothelin and/or megakaryocyte potentiating factor in peritoneal fluids
A Rump, Y Fujii - US Patent 8,110,369, 2012 - Google Patents
... As a consequence, PD has a superior survival rate during the first two years of ... and consistent evidence over the years indicating that mesothelin and MPF are distinct biomarkers with different ... of occurrence of MSLN or MPF, or fragments thereof, in a patient's peritoneal fluid can ...
所有 5 个版本 引用 保存

☒ 包括专利

☒ 包含引用

☐ 创建快讯

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 32,400 条结果 (用时 0.58 秒)

Effluent Tenascin-C Levels Reflect Peritoneal Deterioration ...

www.hindawi.com/journals/bmri/2015/241098/ ▾ [翻译此页](#)

2015年10月29日 - Peritoneal dialysis (PD) is a treatment for patients with severely reduced or ... Some effluent biomarkers, such as matrix metalloproteinase-2 (MMP-2), should serve as a superior indicator of general peritoneal deterioration.

Peritoneal dialysis - 2

ckj.oxfordjournals.org/content/2/suppl_2/ii1937.abstract ▾ [翻译此页](#)

作者: T Kusumoto - 2009

Further, although matrix metalloproteinase-2 (MMP-2) has been shown to chronic peritoneal dialysis could prevent the deterioration of the peritoneal CYSTATIN C: A GOOD BIOMARKER FOR EVALUATING RESIDUAL RENAL been designed to determine whether Medihoney™ will be superior to nasal mupirocin.

[PDF] PROGRESS IN PERITONEAL DIALYSIS

www.zums.ac.ir/files/research/site/.../Progress_in_Peritoneal_Dialysis.pdf ▾

Progress in Peritoneal Dialysis. 2 semipermeable membrane that represents various Functional deterioration of the peritoneum as a dialyzing organ is a leading cause of injury biomarkers (e.g. hyaluronic acid, IL-6, MMP-2) in the peritoneal This is considered superior to other currently available techniques.

Cyclooxygenase-2 Mediates Dialysate-Induced Alterations ...

<https://jasn.asnjournals.org/content/20/3/582.full> - [翻译此页](#)

作者: LS Aroeira - 2009 - 被引用次数: 41 - 相关文章

During peritoneal dialysis (PD), exposure of the peritoneal membrane to ... COX-2 inhibition as a potential strategy to ameliorate peritoneal deterioration of MCs, the accumulation of extracellular matrix (ECM), and angiogenesis. inducing membrane-



全部

图片

新闻

视频

更多 ▾

搜索工具

找到约 32,100 条结果 (用时 0.84 秒)

Effluent Tenascin-C Levels Reflect Peritoneal Deterioration ...

www.hindawi.com/journals/bmri/2015/241098/ ▾ 翻译此页

2015年10月29日 - Peritoneal dialysis (PD) is a treatment for patients with severely reduced or ... Some effluent biomarkers, such as matrix metalloproteinase-2 (MMP-2), should serve as a superior indicator of general peritoneal deterioration.

Peritoneal dialysis - 2

ckj.oxfordjournals.org/content/2/suppl_2/ii1937.abstract ▾ 翻译此页

作者: T Kusumoto - 2009

Further, although matrix metalloproteinase-2 (MMP-2) has been shown to chronic peritoneal dialysis could prevent the deterioration of the peritoneal CYSTATIN C: A GOOD BIOMARKER FOR EVALUATING RESIDUAL RENAL been designed to determine whether Medihoney™ will be superior to nasal mupirocin.

BioMed Research International - JournalTOCs

www.journaltoocs.ac.uk > Browse by Subject ▾ 翻译此页

Abstract: Peritoneal deterioration causing structural changes and functional decline is a major complication of peritoneal dialysis (PD). The aim of this study was to explore effluent biomarkers reflecting peritoneal deterioration. ... Effluent matrix metalloproteinase-2 (MMP-2) levels increased in both the MGO- and FA-treated ...

[PDF] PROGRESS IN PERITONEAL DIALYSIS

www.zums.ac.ir/files/research/site/.../Progress_in_Peritoneal_Dialysis.pdf ▾

Progress in Peritoneal Dialysis. 2 semipermeable membrane that represents various Functional deterioration of the peritoneum as a dialyzing organ is a leading cause of injury biomarkers (e.g. hyaluronic acid, IL-6, MMP-2) in the peritoneal This is considered superior to other currently available techniques.